Czechoslovak Mathematical Journal

Sahib Ram Mandan

Polarity for a simplex

Czechoslovak Mathematical Journal, Vol. 16 (1966), No. 2, 307–313

Persistent URL: http://dml.cz/dmlcz/100732

Terms of use:

© Institute of Mathematics AS CR, 1966

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Чехословацкий математический журнал, т. 16 (91) 1966, Прага

POLARITY FOR A SIMPLEX[1])

SAHIB RAM MANDAN, Kharagpur (India)

(Received June 26, 1965)

The main purpose of this paper is to relate the first polar, for a simplex S in an n-space, of a point with a locus (Theorems 10, 11, 14, 15) and that of a hyperplane with an envelope (Theorems 8, 9, 13) associated with a pair of harmonic systems of quadrics respectively inscribed to an S-configuration ((S−C) and circumscribed to its dual or reciprocal (R.S−C), S being self polar for quadrics of both systems (Theorem 12) as the common diagonal simplex of the (S−C) and (R.S−C). Their association with isotomic and isogonal transformations too is observed. The relation of S with the polar quadric of a point as well as of a hyperplane for S is also considered (Theorems 2, 3).

1. INTRODUCTION

Let S be a simplex in a projective space of n dimensions, or briefly in an n-space, denoted by $[n]$.

We may consider S to be a *degenerate primal of order* $n + 1$ represented by the joint equation $x_0 \ldots x_n = 0$ of its $n + 1$ primes $a_i \equiv x_i = 0$ or *of class* $n + 1$ represented tangentially by the joint equation $u_0 \ldots u_n = 0$ of its $n + 1$ vertices $A_i \equiv u_i = 0$. The *first polar* ([18], p. 11) of a point $Z(Z_0, \ldots, Z_n)$ for S is then found to be the primal $\sum(Z_i/x_i) = 0$, of order n, circumscribed to S; ...; *the* $(n - 1)$*th polar* is the quadric $(Z) \equiv \sum(x_i x_j/Z_i Z_j) = 0$ $(i \neq j)$, circumscribed to S, called the *polar quadric of* Z *for* S; the *nth polar* is the hyperplane $z \equiv \sum(x_i/Z_i) = 0$, called the *polar of* Z *for* S. The $n + 1$ *tangential coordinates* of z are then $u_i = 1/Z_i$.

Tangentially, the first polar (cf. [17], p. 118) of z for S may be defined to be similarly the primal $\sum(z_i/u_i) = 0$, of class n, inscribed to S; ...; the $(n - 1)$th polar as its polar quadric $(z) \equiv \sum(u_i u_j/z_i z_j) = 0$, inscribed to S; the nth polar as its *pole* $Z \equiv \sum(u_i/z_i) = 0$.

As an immediate consequence we have the following

[1]) Proceedings of the 48th Session of the Indian Science Congress Association held at Roorkee in the first week of January 1961, p. 8.

Theorem 1. *The polar z of Z or the pole Z of z for a simplex S is the same for its polar primals, w.r.t. S, of all orders, in particular for its first polar as well as for its polar quadric* (cf. [17], p. 51).

2. POLAR QUADRIC

The tangent hyperplane of the polar quadric (Z), of the point Z (§ 1) for the simplex S, at its vertex A_j is given by

$$\sum(x_i/Z_i) - x_j/Z_j = 0 .$$

It is evidently coaxal with the prime $x_j = 0$ and the polar prime $\sum(x_i/Z_i) = 0$ of Z for S or (Z).

Again the point of contact of the polar quadric (z), of the hyperplane z (§ 1) for the simplex S, with its prime a_j is given by

$$\sum(u_i/z_i) - u_j/z_j = 0 .$$

It is obviously collinear with the vertex $u_j = 0$ of S and the pole $\sum(u_i/z_i) = 0$ of z for S, or (z). Thus we have the following

Theorem 2. *The polar reciprocal S'' of a simplex S in $[n]$ w.r.t. the polar quadric (Z) of a point Z for S is the "anticevian simplex" of S for Z, and that, say S', w.r.t. the polar quadric (z) of a hyperplane z for S is the "cevian simplex" of S for z. If Z, z be the pole and polar for S, they are also pole and polar for both (Z) and (z) as the centre and hyperplane of perspectivity of the 3 simplexes S, S', S''. Hence the vertices of S and Z form a "self-conjugate" $(n + 2)$ad of points for both (Z) and (z) such that the join of any two of them is conjugate to the hyperplane of the other n points for both (Z) and (z). Dually the primes of S and z form a self-conjugate $(n + 2)$ad of hyperplanes for both (z) and (Z) such that the polar hyperplanes for (z) and (Z), of the point common to any n of them pass through the common $[n - 2]$ of the other two hyperplanes of the $(n - 2)$ad.* (cf. [6], p. 97, ex. 5; 11; 13).

3. ISODYNAMIC AND ISOGONIC SIMPLEXES

We may define an *isodynamic simplex* S as one whose solid faces all form *isodynamic tetrahedra* such that the tangent hyperplanes of its circumhypersphere (S) at its vertices form a simplex S'' perspective with S. Consequently S'' is said to be *isogonic* such that the simplex S formed of the points of contact of its inscribed hypersphere (S) is perspective with it. The centre L of perspectivity of S, S'' may be then referred to as the *Lemoine point* of S, *Gergonne point* following COXETER ([7])

or *Fermat point* ([15]) of S'', and the hyperplane l of their perspectivity as the *Lemoine prime* of S in analogy with those of such tetrahedra ([5]; [14]). Now follows from Theorem 2 the following

Theorem 3. *The polar quadric of a point L (hyperplane l) for a simplex S is a hypersphere (S), if and only if S is isodynamic (isogonic) such that L, l are pole and polar for both S and (S).*

4. POLARITY FOR A QUADRIC

Following COURT ([1]) we can prove the following

Theorem 4. *If Y, z be pole and polar for a quadric Q, and S, S'' be polar reciprocal simplexes for Q, the pole Z of z and the polar y of Y for S and S'' respectively are also pole and polar for Q.*

Theorem 5. *If z is a tangent hyperplane for a quadric Q, at a point Y on it, the pole Z of z and the polar y of Y for a simplex S, self-polar for Q, are the pole and polar for Q.*

5. AN S-CONFIGURATION

a. If P_{ij} be the trace of a hyperplane z (§ 1) on an edge A_iA_j of a simplex S, and Q_{ij} be its harmonic conjugate on this edge w.r.t. A_i, A_j, the points P_{ij}, Q_{ij} then lie by $\binom{n+1}{2}$s in the 2^n hyperplanes, like z, of an *S-configuration* (S−C) as the $\binom{n+1}{2}$ pairs of its opposite vertices, S being the diagonal simplex ([8]). The equations of these hyperplanes referred to S are

$$x_0/Z_0 \pm x_1/Z_1 \pm \ldots \pm x_n/Z_n = 0$$

b. The 2^n points $(Z_0, \pm Z_1, \ldots, \pm Z_n)$ form an *associated* [(10)] or *closed set* ([8]; [10]) as the vertices of the *dual* or *reciprocal* (R.S−C) of the (S−C) such that all the quadrics, for which their common diagonal simplex S is self-polar, passing through one of them, pass through all of them. Conversely we can show the following

Theorem 6. *The quadrics circumscribed to an* (R.S−C) *form a system s such that its diagonal simplex S is self-polar for all of them.*

They are therefore represented by the equations

$$\sum k_i x_i^2 = 0 = \sum k_i Z_i^2 \,, \quad \text{referred to } S \,.$$

Dually we shall have the following

Theorem 7. *The quadrics inscribed to an* (S−C) *form a tangential system* s' *such that its diagonal simplex* S *is self-polar for all of them.*

They are therefore represented tangentially by the equations referred to S,

$$\sum p_i u_i^2 = 0 = \sum p_i z_i^2 = 0\,,$$

the tangential coordinates of the hyperplanes of (S−C) being $(z_0, \pm z_1, \ldots, \pm z_n)$.

6. SYSTEMS OF QUADRICS

a. The point of contact Y of a quadric Q' of the tangential system s' (Theorem 7) inscribed in the (S−C) with its hyperplane z (§§ 1, 5) is given by $\sum p_i u_i z_i = 0 = \sum p_i z_i^2$. The tangential coordinates of the polar y of Y for the diagonal simplex S of (S−C) are then $u_i = 1/p_i z_i$ where $\sum p_i z_i^2 = 0$. Thus the envelope of y, as Q' varies is the primal $\sum z_i/u_i = 0$ (§ 1).

Again by the Theorem 4 or otherwise by § 1, y is the polar prime for Q', of the pole Z of z for S. Hence we have the following

Theorem 8. *The polars of the point of contact of the quadrics* Q' *of a tangential system* s' *inscribed in an* (S−C) *with one of its* 2^n *hyperplanes, say* z, *for its diagonal simplex* S *or the polars, for* Q', *of the pole* Z *of* z *for* S *envelope a primal of class* n *no other than the first polar of* z *for* S.

Theorem 9. *The envelope of the polar of a variable point of a given hyperplane* z *for a simplex* S *coincides with the first polar of* z *for* S. (cf. [6], p. 97, ex. 3)

b. Dually thus we have the following

Theorem 10. *The poles of the tangent hyperplanes of the quadrics* Q *of a system* s *circumscribed to an* (R.S−C) *at one of its vertices, say* Z, *for its diagonal simplex* S *or the poles, for* Q, *of the polar of* Z *for* S *describe a primal of order* n *no other than the first polar of* Z *for* S.

Theorem 11. *The locus of the pole of a variable hyperplane through a given point* Z *for a simplex* S *coincides with the first polar of* Z *for* S (cf. [6], p. 97).

c. The 2 systems s, s' (Theorems 6−8, 10) of quadrics are evidently related dually and may be said to be *harmonically associated* (cf. [1]), if the vertices of the (R.S−C) inscribed to s are the poles of the hyperplanes of the (S−C) circumscribed to s' w.r.t. their common diagonal simplex S. In fact, there exists a quadric W for which

s, s' are polar reciprocal, viz., $W \equiv \sum z_i^2 x_i^2 = 0$ (which is the same as $\sum Z_i^2 u_i^2 = 0$ tangentially, where $Z_i z_i = 1$ (§ 1)). For in this polarity, the quadric $Q \equiv \sum k_i x_i^2 = \sum k_i Z_i^2$ of s corresponds to $Q' \equiv \sum p_i u_i^2 = 0 = \sum p_i z_i^2$ of s' for $p_i = k_i Z_i^4$. Thus we have

Theorem 12. *The harmonically associated systems of quadrics have a common self-polar simplex S, and their relation is reciprocal such that one reciprocates into other w.r.t. a quadric W for which too S is self-polar.*

7. ISOTOMIC TRANSFORMATION

a. A pair of points Z_{ij}, Z'_{ij} on an edge A_iA_j of a simplex S equidistant from its midpoint M_{ij} are said to be *isotomic conjugates* w.r.t. A_iA_j ([4]; [19]). If Z_{ij} be the feet of the *bicevians* of S through a point Z (secants through Z to its edges and the respectively opposite $[n-2]$s) on its edges, it is shown in [13] that their isotomic conjugates Z'_{ij} thereat are also the feet of the bicevians of S through another point Z'. The pair of points like Z, Z' are said to be *isotomic conjugates for S*, and their polars z, z' for S are consequently called *isotomically conjugate hyperplanes* for S. Thus: *The 2^n vertices of the* (R.S−C) *with one vertex at the centroid G of its diagonal simplex S and their polar hyperplanes for S, that of G being at infinity, are all isotomically self-conjugate for S.*

b. It can be shown that if G be taken as unit point $(1, 1, \ldots, 1)$ of S, and the coordinates of Z be as before (§ 1), those of its isotomic conjugate point Z' are proportional to their reciprocals respectively. That is, $Z_iZ'_i = k$ (say). Similarly, therefore, are related the tangential coordinates of z, z' too. That is, $z_iz'_i = 1/k = k'$ (say), for $Z_iz_i = 1 = Z'_iz'_i$ (§ 1). Thus $Z_i = kz'_i$, $z_i = k'Z'_i$.

c. Consider a variable hyperplane u' $(u'_0, \ldots, u'_n)$ through the given point Z', and its isotomic conjugate hyperplane u $(u_0, \ldots, u_n)$ for the simplex S. Then $\sum Z'_iu'_i = 0$, $u_iu'_i = k'$. Hence, as u' varies through Z', u envelope the primal $\sum Z'_i/u_i = 0$ which is no other than the first polar (§ 1) of the hyperplane z for S. For its tangential coordinates are $z_i = k'Z'_i$. Thus we have the following

Theorem 13. *The envelope of the isotomic conjugates of the hyperplanes through a given point Z′ for a simplex S in an* $[n]$ *is a primal of class n no other than the first polar, for S, of the isotomic conjugate hyperplane z of the polar z′ of Z′ for S.*

d. Dually thus we have the following

Theorem 14. *The locus of the isotomic conjugate of a variable point in a given hyperplane z′ for a simplex S in an* $[n]$ *is a primal of order n no other than the first polar, for S, of the isotomic conjugate point Z of the pole Z′ of z′ for S* (cf. [3]).

8. ISOGONAL TRANSFORMATION

The coordinates of a pair of *isogonal conjugate points* (cf. [6], [9], [16], [19]) Z, Z' for a simplex S with unit point at its incentre I are also seen to be related reciprocally. That is, $Z_i Z'_i$ is a constant. Hence following the argument of the preceding section we have

Theorem 15. *The locus of the isogonal conjugate of a variable point in a given hyperplane z' for a simplex S in an $[n]$ is a primal of order n other than the first polar, for S, of the isogonal conjugate point Z of the pole Z' of z' for S.*

9. WHEN $n = 2$

The first polar and the polar quadric of a point Z in the plane of a triangle t for t is the polar conic ([4]) of Z for t. The harmonically associated system (§ 5) s, s' of quadrics become respectively a pencil and a range of conics for which t is self-polar. For detail of these particular cases of evident interest reference be made to Court ([1]; [2]; [3]) who has also discussed the isotomic conics of the 4 isotomically self-conjugate transversals of t one being the line at infinity in the plane of t.

Thanks are due to Prof. B. B. SETH for his generous, kind and constant encouragement in my pure pursuits.

References

[1] *N. A. Court:* Pencils of conics. Bol. Mat., *28* (1955), 7—15.

[2] *N. A. Court:* Three hyperbolas associated with a triangle. Am. Math. Mon., *64* (1957), 241—247.

[3] *N. A. Court:* Sur la transformation isotomique. Mathesis, *66* (1957), 291—297.

[4] *N. A. Court:* Sur la cubique gauche, application de la transformation harmonique. Mathesis, *68* (1959), 110—127.

[5] *N. A. Court:* Modern Pure Solid Geometry. New York, 1964.

[6] *H. S. M. Coxeter:* The real projective plane. Cambridge, (1955).

[7] *H. S. M. Coxeter:* Review A 1253. Math. Reviews *23* (1962), 224.

[8] *S. R. Mandan:* An *S*-configuration in Euclidean and elliptic *n*-space. Can. J. Math., *10* (1958), 489—501.

[9] *S. R. Mandan:* Semi-orthocentric and orthogonal simplexes in a 4-space. Bul. Cal. Math. Soc. *50* (Supp. 1958), 21—29.

[10] *S. R. Mandan:* Harmonic Inversion. Math. Mag. *32* (1959), 13—20.

[11] *S. R. Mandan:* Polarity for a quadric in an *n*-space. Rev. Fac. Soc. Uni. Istanbul, A *24* (1959), 21—40.

[12] *S. R. Mandan:* A closed set of 2^n points in an *n*-space. Math. Sem. *1* (1960), 43—44.

[13] *S. R. Mandan:* Cevian simplexes. Proc. Amer. Math. Soc. *11* (1960), 837—845.

[14] *S. R. Mandan:* Semi-isodynamic and isogonic tetrahedra. Rend. Math. (3—4), *19* (1960), 401—415.

[15] *S. R. Mandan:* Isodynamic and isogonal simplexes. Ann. Mat. (4) *53* (1961), 45—55, (To Enricho Bompiani on his Scientific Jubilee).
[16] *S. R. Mandan:* Altitudes of a simplex in an *n*-space. Jour. Australian Math. Soc., *2* (1962), 403—424.
[17] *G. Salmon:* Higher plane curves. (New York, 1934).
[18] *J. G. Semple* and *L. Roth:* Algebraic Geometry. (Oxford 1949).
[19] *D. M. Y. Sommerville:* Analytical conics. (London, 1949).

Author's address: Kharagpur, India (Indian Institute of Technology).

Резюме

ПОЛЯРНЫЕ СООТНОШЕНИЯ ДЛЯ СИМПЛЕКСА

САГИБ РАМ МАНДАН (Sahib Ram Mandan), Харагпур (Индия)

Изучаются полярные соотношения относительно симплекса и их связ с изотомическим и изогональним соотношениями.