

P. L. Ganguli; Benoy Kumar Lahiri
Some results on certain sets of series

Czechoslovak Mathematical Journal, Vol. 18 (1968), No. 4, 589–594

Persistent URL: <http://dml.cz/dmlcz/100857>

Terms of use:

© Institute of Mathematics AS CR, 1968

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SOME RESULTS ON CERTAIN SETS OF SERIES

P. L. GANGULI, B. K. LAHIRI, Calcutta

(Received February 27, 1967)

Introduction. Let $\sum_1^{\infty} u_n$ be conditionally convergent series. Then it is known that

(i)
$$u_n \rightarrow 0 \text{ as } n \rightarrow \infty$$

and (ii) the series of positive terms of $\sum_1^{\infty} u_n$ and the series of absolute values of negative terms of $\sum_1^{\infty} u_n$ both diverge to $+\infty$.

Any series $\sum_1^{\infty} u_n$ which satisfies conditions (i) and (ii) will be said, in what follows, to be of type (A).

The authors of the present note have proved [4] that the results of Riemann's theorem [5] on conditionally convergent series all hold for any series of the type (A).

Let $\sum_1^{\infty} u_n$, which will sometimes be written as $\sum_1^{\infty} u(n)$, denote any infinite series. To each rearrangement of the series $\sum_1^{\infty} u(n)$ there corresponds a complex (x_1, x_2, x_3, \dots) and conversely, where x_1, x_2, x_3, \dots is a permutation of the set of all positive integers. The set of all such complexes form a metric space E [1] with the metric

$$d(x, y) = \sum_{n=1}^{\infty} \frac{1}{2^n} \cdot \frac{|x_n - y_n|}{1 + |x_n - y_n|}$$

where $x = (x_1, x_2, \dots)$ and $y = (y_1, y_2, \dots)$.

Let $\sum_1^{\infty} c(n)$ be any conditionally convergent series and let A denote the proper subset of E consisting of all points of E which correspond to convergent rearrangements of $\sum_1^{\infty} c(n)$. Further, let $A(\alpha)$ denote the proper subset of A consisting of those points of A which correspond to rearrangements of $\sum_1^{\infty} c(n)$ converging to the given

real number α . Then it has been shown by H. M. SEN GUPTA [7] that the sets A and $A(\alpha)$ are dense boundary subsets of E . It immediately follows that the above result holds for any series of the type (A).

The purpose of the present paper is to extend certain results of Sengupta and to determine the cardinal number of sets of certain types of series.

I

The set $A(\lambda, \mu)$. Let $\sum_1^\infty u_n$ be any series of the type (A) and let λ and μ ($\lambda \geq \mu$) be any two preassigned real numbers, finite or infinite. We shall denote by $A(\lambda, \mu)$ the set of all points of E which correspond to those rearrangements of $\sum_1^\infty u_n$ for which (S_n , as usual, denoting the n -th partial sum)

$$\overline{\lim} S_n = \lambda \quad \text{and} \quad \underline{\lim} S_n = \mu.$$

It is obvious that $A(\lambda, \lambda) = A(\lambda)$.

Theorem 1. *The set $A(\lambda, \mu)$ is a dense boundary subset of E .*

Proof. Let $\varepsilon > 0$ be given. We can then choose a positive integer N such that $2^{-N} < \varepsilon$. Let $p \equiv (a_1, a_2, \dots)$ be any point of E . If $q \equiv (b_1, b_2, \dots) \neq p$ be any point of the sphere $S(p, 2^{-N-1})$, then $b_i = a_i$ for $i = 1, 2, \dots, N$. Let M be the greatest positive integer such that $b_i = a_i$ for $i = 1, 2, \dots, M$. Then, clearly $M \geq N$.

We now consider the series

$$u(b_{M+2}) + u(b_{M+3}) + \dots$$

which is clearly of the type (A). Hence there exists a rearrangement of this series, say the series

$$u(c_{M+2}) + u(c_{M+3}) + \dots$$

for which $\overline{\lim} S_n = \lambda - \alpha$ and $\underline{\lim} S_n = \mu - \alpha$ where $\alpha = u(a_1) + u(a_2) + \dots + u(a_M) + u(b_{M+1})$.

It follows that for the series

$$u(a_1) + \dots + u(a_M) + u(b_{M+1}) + u(c_{M+2}) + u(c_{M+3}) + \dots,$$

$$\overline{\lim} S_n = \lambda \quad \text{and} \quad \underline{\lim} S_n = \mu.$$

Also the point $(a_1, a_2, \dots, a_M, b_{M+1}, c_{M+2}, c_{M+3}, \dots)$ lies in $S(p, \varepsilon)$ and is distinct from p . This shows that the set $A(\lambda, \mu)$ is everywhere dense in E . It is easy to show that $A(\lambda, \mu)$ is a boundary set. This completes the proof of Theorem 1.

Taking $\lambda = \mu = \alpha$ (finite), we get Sengupta's Theorem [7].

It easily follows that each of the sets $A(\infty, \mu)$, $A(\lambda, -\infty)$, $A(\infty, -\infty)$, $A(\infty, \infty)$, $A(-\infty, -\infty)$ is a dense boundary subset of E .

Let $\bar{A}(\lambda)$ denote the set of all points of E which correspond to those rearrangements of $\sum_1^\infty u_n$ for which $\overline{\lim} S_n = \lambda$ (whatever be $\underline{\lim} S_n$). Then clearly $\bar{A}(\lambda) \supset A(\lambda, \mu)$ and it follows that $\bar{A}(\lambda)$ is also a dense boundary subset of E . Similarly, the set $\underline{A}(\mu)$ consisting of those points of E for which $\underline{\lim} S_n = \mu$ is a dense boundary subset of E .

Mapping f .: Let $\sum_1^\infty u_n$ be a series of the type (A) and let B denote the subset of E which consists of those points of E for which $\lambda \equiv \overline{\lim} S_n$ is finite. We define a mapping f , of the subspace B onto the real number space E_1 , by the following rule

$$f(p) = \overline{\lim} S_n \quad \text{for } p \in B.$$

The following properties of the mapping f (proofs of which are similar to those of Sengupta's [7] corresponding theorems) may be stated.

1. The function f is everywhere discontinuous over the set B .
2. The map f from B to E_1 is open in the sense that each non-empty subset of B which is open in B is mapped into a non-empty open set in E_1 .
3. The map f from B to E_1 is not closed in the sense that there is at least one closed set in B whose image is not closed in E_1 .

II

In this section we shall prove certain theorems on the Cardinal number of sets of certain types of series.

Theorem 2. Let $\sum_1^\infty a_n$ be any series of the type (A) and let λ and μ ($\lambda \geq \mu$) be any two preassigned numbers. Then the set of all series obtainable from $\sum_1^\infty a_n$ by rearrangement of its terms, for each of which the sequence of partial sums has λ and μ as its upper and lower limits respectively has the power of the continuum.

Proof. Let $\sum_1^\infty a_n$ be any series of the type (A). Let p_1, p_2, p_3, \dots denote the non-negative terms and q_1, q_2, q_3, \dots the negative terms of $\sum_1^\infty a_n$, the terms being taken in the order in which they appear in the original series. It is then possible to decompose

$\sum_1^{\infty} p_n$ into two subseries $\sum_1^{\infty} p'_n$ and $\sum_1^{\infty} p''_n$ and $\sum_1^{\infty} q_n$ into $\sum_1^{\infty} q'_n$ and $\sum_1^{\infty} q''_n$ such that the series $\sum_1^{\infty} p'_n, \sum_1^{\infty} p''_n, \sum_1^{\infty} q'_n, \sum_1^{\infty} q''_n$ are all properly divergent.

Now, form the series $\sum_1^{\infty} a'_n$, taking all the terms of $\sum_1^{\infty} p'_n$ and $\sum_1^{\infty} q'_n$ (in the order in which they appear in $\sum_1^{\infty} a_n$). Similarly, we construct the series $\sum_1^{\infty} a''_n$ with p''_n and q''_n as its terms. From the very manner of construction of the two series $\sum_1^{\infty} a'_n$ and $\sum_1^{\infty} a''_n$, it is clear that each is a series of the type (A). Also every term of $\sum_1^{\infty} a_n$ occurs once and only once in one of $\sum_1^{\infty} a'_n$ and $\sum_1^{\infty} a''_n$.

Now, for every real number α , there exist rearrangements $\sum_1^{\infty} l'_n$ and $\sum_1^{\infty} l''_n$ of $\sum_1^{\infty} a'_n$ and $\sum_1^{\infty} a''_n$ respectively such that

$$\overline{\lim} B'_n = \lambda - \alpha, \quad \underline{\lim} B'_n = \mu - \alpha \quad \text{and} \quad \lim B''_n = \alpha$$

where $B'_n = l'_1 + l'_2 + \dots + l'_n, B''_n = l''_1 + l''_2 + \dots + l''_n$.

Now, consider the series

$$l'_1 + l''_1 + l'_2 + l''_2 + \dots$$

which is a rearrangement of the original series $\sum_1^{\infty} a_n$. If Q_n denotes the n -th partial sum of this series, it easily follows that

$$\overline{\lim} Q_n = \lambda \quad \text{and} \quad \underline{\lim} Q_n = \mu.$$

This shows that the power of our set is $\geq c$, the power of the continuum. But it is known that the power of the set E is c [6]. This proves the theorem.

Taking $\lambda = \mu = \alpha$ we get the following theorem.

Theorem 3. Let $\sum_1^{\infty} a_n$ be any series of the type (A) and let α be any preassigned number. Then the set of all series, obtainable from $\sum_1^{\infty} a_n$ by rearrangement of its terms, which converge to α has the power of the continuum.

It is easy to see that Theorem 2 remains valid in each of the following cases also.

- I. $\lambda = +\infty, \mu$ is finite,
- II. λ is finite, $\mu = -\infty$,
- III. $\lambda = +\infty, \mu = -\infty$,
- IV. $\lambda = \mu = \pm\infty$.

Theorem 4. Let $\sum_1^{\infty} a_n$ be any series of the type (A) and let α be any number given in advance. Then the set of all subseries of $\sum_1^{\infty} a_n$ which converges to α has the power of the continuum.

Proof. Let us suppose that $\alpha > 0$. Let x denote any number in $0 < x \leq 1$. Then, using the same notation as in Theorem 2, there exists [3] a subseries $\sum_1^{\infty} p_{n_k}$ of $\sum_1^{\infty} p_n$ and a subseries $\sum_1^{\infty} q_{m_k}$ of $\sum_1^{\infty} q_n$ such that

$$\sum_1^{\infty} p_{n_k} = \alpha + x \quad \text{and} \quad \sum_1^{\infty} q_{m_k} = -x.$$

The series

$$p_{n_1} + q_{m_1} + p_{n_2} + q_{m_2} + \dots$$

clearly converges absolutely to α and by means of a suitable rearrangement of this series we obtain a subseries of $\sum_1^{\infty} a_n$, the sum of which is α . It follows that the set of all such series has power $\geq c$, but it is easy to see that the set of all subseries of a given series has the power c . Similar proof applies to the case when $\alpha \leq 0$. The theorem is thus proved.

Theorem 5. Let $\sum_1^{\infty} a_n$ be a divergent series of positive terms where $a_n \rightarrow 0$ as $n \rightarrow \infty$ and let s be any positive number chosen in advance. Then the set of all subseries of $\sum_1^{\infty} a_n$ which converges to s has the power c .

Proof. We decompose $\sum_1^{\infty} a_n$ into two subseries $\sum_1^{\infty} b_n$ and $\sum_1^{\infty} c_n$ such that the series $\sum_1^{\infty} b_n$ and $\sum_1^{\infty} c_n$ are both divergent.

Let x denote any number in $0 < x < \frac{1}{2}s$. Then it is possible to find [3] subseries $\sum_1^{\infty} b_{n_k}$ and $\sum_1^{\infty} c_{m_k}$ of $\sum_1^{\infty} b_n$ and $\sum_1^{\infty} c_n$ respectively such that

$$\sum_1^{\infty} b_{n_k} = \frac{1}{2}s + x \quad \text{and} \quad \sum_1^{\infty} c_{m_k} = \frac{1}{2}s - x.$$

The series of positive terms

$$b_{n_1} + c_{m_1} + b_{n_2} + c_{m_2} + \dots$$

converges to s and by means of a suitable rearrangement of this series we obtain a subseries of $\sum_1^{\infty} a_n$, the sum of which is s . It follows that the required power is c .

It easily follows that the set of all subseries of a divergent series $\sum_1^{\infty} a_n$ of positive terms (where $a_n \rightarrow 0$) which themselves are divergent has the power c . It also follows that the set of all divergent subseries $\sum_1^{\infty} a_{n_k}$ of $\sum_1^{\infty} a_n$, for which $n_{k+1} - n_k \rightarrow \infty$ as $k \rightarrow \infty$ [2] has the power c .

References

- [1] *Agnew, R. P.*: On rearrangement of series, *Bull. Amer. Math. Soc.* Vol. 46 (1940), 797–99.
- [2] *Agnew, R. P.*: Subseries of series which are not absolutely convergent, *Bull. Amer. Math. Soc.* Vol. 53, Part I, 1947, P. 118.
- [3] *Banerjee C. R., Lahiri, B. K.*: On subseries of divergent series, *Amer. Math. Monthly*, Vol. 71, No. 7, 1964, 767–768.
- [4] *Ganguli, P. L., Lahiri, B. K.*: A note on oscillatory series. To appear in *Bull. Cal. Math. Soc.*
- [5] *Knopp, K.*: *Theory and application of infinite Series*, 1947, P 318.
- [6] *Sengupta, H. M.*: On rearrangement of series, *Proc. Amer. Math. Soc. Vol. 1*, No. 1, 1950, 71–75.
- [7] *Sengupta, H. M.*: Rearrangement of series, *Proc. Amer. Math. Soc. Vol. 7*, No. 3, 1956, 347–350.

Author's addresses: B. K. Lahiri, Kalyani University, West Bengal, India; P. L. Ganguli, Dept. of Pure Mathematics, Calcutta University, India.