Czechoslovak Mathematical Journal

Paul R. Meyer

The Sorgenfrey topology is a join of orderable topologies

Czechoslovak Mathematical Journal, Vol. 23 (1973), No. 3, 402–403

Persistent URL: http://dml.cz/dmlcz/101181

Terms of use:

© Institute of Mathematics AS CR, 1973

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 23 (98) 1973, Praha

THE SORGENFREY TOPOLOGY IS A JOIN OF ORDERABLE TOPOLOGIES

PAUL R. MEYER, New York[1])

(Received February 17, 1972)

The Sorgenfrey half open interval topology on the real line R is shown to be the join (in the lattice of all topologies on R) of two orderable topologies. This is in contrast with a result of Lutzer [3] who showed that this topology is not orderable.

To prove the result in the title, let $X = (]0, 1[\times \{0, 1\}) \cup \{(1, 0)\}$. We construct order topologies t_1 and t_2 on X such that (X, t) is homeomorphic to the Sorgenfrey line, where $t = t_1 \vee t_2$. Let t_1 be the lexicographic order topology and t_2 the usual euclidean topology, which is orderable. Since t-neighborhoods are formed by intersecting t_i-neighborhoods, we see that, locally, t is the Sorgenfrey topology. Since X contains the point $(1, 0)$ but not $(1, 1)$, the two intervals can be fitted together to form a single interval with the desired topology.

The Sorgenfrey topology is known to be a generalized order topology (GO topology) in the sense of Čech [1, page 245]. A join of orderable topologies need not be a GO topology; in fact an example in [4] shows that it need not even be a chain net topology. It is still an open question whether or not there exists a GO topology which is not a join of orderable topologies.

For connected spaces, it is known that the concepts of orderable and GO topologies are equivalent; the following proposition shows that the notion of a join of orderable topologies is also equivalent. Furthermore, this proposition extends the uniqueness criterion for possible orderings of connected orderable spaces [1, Corollary c, page 361]. (This uniqueness criterion can be used [1, page 844] to show that certain (GO) subspaces of the real line such as $Y =]0, 1[\cup \{2\}$ are not orderable. However, it is easy to construct two orderable topologies on Y whose join is the usual topology.)

Proposition. *If (X, t) is connected and t is the join of orderable topologies t_1 and t_2, then $t = t_1 = t_2$.*

[1]) The author is grateful to Westfield College, University of London, for their hospitality during the preparation of this note.

Proof. For $i = 1, 2$, let $\leqq_i$ denote an ordering of X which induces t_i. Since $t \supset t_i$, t_i is connected. It suffices to show $\leqq_1 = \leqq_2$ or $\leqq_1 = \leqq_2^{-1}$. If not, we may assume there are distinct points x, y, and z with $x \leqq_1 y \leqq_1 z$ and $x \leqq_2 z \leqq_2 y$. Since each t_i-separation is a t-separation, we can t-separate these three distinct points in more than one way. This is a contradiction [2, Lemma 3].

Note added in proof. The dual question has also been answered: the Sorgenfrey topology is not a finite intersection of orderable topologies, although it is the intersection of infinitely many orderable topologies. These follow from more general results to appear in: *Lattice operations on metric and order topologies,* Proceedings of 1973 topology conference, Blacksburg, Virginia.

References

[1] *E. Čech:* Topological spaces, ČSAV, Prague, 1966. MR *35* # 2254.

[2] *H. Kok:* On conditions equivalent to the orderability of a connected space, Nieuw Arch. Wiskunde (3), *18* (1970), 250—270.

[3] *D. J. Lutzer:* A metrization theorem for linearly orderable spaces, Proc. Amer. Math. Soc. 22 (1969), 557—558. MR *40* # 2012.

[4] *P. R. Meyer:* On total orderings in topology, Third Prague Symp. on general topology, 1971, 301—396.

Author's address: Lehman College of the City University of New York, Department of Mathematics, Bronx, New York 10 468, U.S.A.