Czechoslovak Mathematical Journal

Ján Jakubík

Sublattices with saturated chains

Czechoslovak Mathematical Journal, Vol. 25 (1975), No. 3, 442–444

Persistent URL: http://dml.cz/dmlcz/101338

Terms of use:

© Institute of Mathematics AS CR, 1975

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 25 (100) 1975, Praha

SUBLATTICES WITH SATURATED CHAINS[1])

JÁN JAKUBÍK, Košice

(Received May 8, 1974)

Let L be a lattice. A chain $R \subseteq L$ will be called saturated in L if $r_1, r_2 \in R$, $x \in L$, $r_1 < x < r_2$ implies $x \in R$ whenever x is comparable with each element of R. A sublattice L_1 of L is said to be a c-sublattice of L if each chain R of L_1 which is saturated in L_1 is also saturated in L. Let K be a lattice of locally finite length. A sublattice S of K is a c-sublattice of K if and only if, whenever $s_1, s_2 \in S$ and s_1 covers s_2 in S, then s_1 covers s_2 in K.

Some classes of lattices of locally finite length can be characterized by the requirement of not containing c-sublattices of certain types.



Let B be the lattice in Fig. 1 and let B' be the lattice dual to B. Let C be the nondistributive modular lattice with five elements. For positive integers $m \geqq 3$, $n \geqq 3$ we denote by $L(m, n)$ the lattice consisting of elements $x_1 < x_2 < \ldots < x_m$, $y_1 < y_2 < \ldots < y_n$ such that $x_1 = y_1$, $x_m = y_n$ and x_i is incomparable with y_j whenever $1 < i < m$, $1 < j < n$.

The following results are known:

(A) (GRÄTZER [3], p. 151.) *Let L be a finite modular lattice. Then L is nondistributive if and only if it contains a c-sublattice isomorphic to C.*

(B) (Cf. [4].) *Let L be a lattice of locally finite length. Then L is distributive if and only if it does not contain a c-sublattice isomorphic to some of the following lattices*: B, B', $L(m, n)$ $(m \geqq 3,\ n \geqq 4)$, C.

(C) (VILHELM [5], Thm. 2.2.) *Let L be a lattice of locally finite length. Then L is semimodular if and only if it does not contain a c-sublattice isomorphic to B' or $L(m, n)$ $(m \geqq 3,\ n \geqq 4)$.*

In view of these results the following question seems to be quite natural: which classes of lattices can be defined by the requirement of not containing c-sublattices of certain types? Let us formulate the corresponding notions more precisely.

Let K be a class of lattices and let K_1 be a subclass of K. The class K_1 will be said to be characterizable in K by c-sublattices if there exists a set C of lattices belonging to $K \setminus K_1$ such that, for each lattice $L \in K$, the following conditions are equivalent:

(i) L belongs to K_1;

(ii) if L_1 is a c-sublattice of L, then no lattice $L' \in C$ is isomorphic to L_1.

Let K_0 be the class of all lattices. Further let K_d and K_m be the class of all distributive lattices and the class of all modular lattices, respectively. A variety K of lattices will be called nontrivial if there exists a lattice $L \in K$ with card $L > 1$.

We have the following negative result:

Theorem. *Let K be a nontrivial variety of lattices such that $K \subseteqq K_m$. Then K is not characterizable in K_0 by c-sublattices.*

Proof. Suppose that K is characterizable in K_0 by c-sublattices. Then there exists a set C of lattices of the class $K_0 \setminus K$ such that a lattice L belongs to K if and only if it does not contain any c-sublattice isomorphic to some of the lattices of C. Let $C = \{L_j\}$ $(j \in J)$, $\alpha = \sup \{\text{card } L_j\}$ and let β be a cardinal with $\beta > \alpha$. Let ω_β be the first ordinal with the property that the power of the set of all lower ordinals equals β. Let A and B be dually isomorphic to ω_β, $C_1 = A \times B$. Let x, y, u be three distinct elements not belonging to C_1, $L = C_1 \cup \{x, y, u\}$.

We define in L a partial order $\leqq$ as follows. If $c_1, c_2 \in C_1$, then $c_1 \leqq c_2$ in L has the same meaning as $c_1 \leqq c_2$ in C_1. The element u is the least element in L. Further we put $x < y$. For $c \in C_1$, $z \in \{x, y\}$ we set $z < c$ if and only if $c = (a, 0)$, where a is any element of A and 0 is the greatest element of B; if c is not of this form, then z and c are incomparable.

It is well-known that the variety K_d of all distributive lattices is the unique atom in the lattice of all varieties of lattices (cf., e.g., [2], p. 182). Hence $K_d \subseteqq K$.

The lattice L is not modular, hence it does not belong to K. Thus according to the assumption there exists a c-sublattice L_0 of L isomorphic to a lattice of the class C.

---

[1]) The results of this paper have been presented at the Conference on lattice theory held at Szeged (August, 1974).

Put $L_0 \cap C_1 = C_0$. If each $c \in C_0$ has the form $(a, 0)$ with $a \in A$, then L_0 is a chain and hence $L_0 \in K_d \subseteqq K$, which is a contradiction. If $\{x, y\} \cap L_0 = \emptyset$, then L_0 is distributive, thus $L_0 \in K$, which is impossible. Hence there is $z \in \{x, y\}$ and $c = (a, b)$, $b \neq 0$, such that $\{z, c\} \subseteqq L_0$. Then

$$c_1 = c \vee z = (a, 0) \in L_0 .$$

Let A_1 be a saturated chain in L_0 having the least element z and the greatest element c_1. Then A_1 is a saturated chain in L between z and c_1. Thus the set $A_2 = \{(a_2, 0) : a_2 \in A,\ a_2 \leqq a\}$ is a subset of A_1. Obviously card $A_2 = \beta$ and hence card $L_0 \geqq \beta$. But card $L_j \leqq \alpha < \beta$ for each $L_j \in C$ and hence L_0 cannot be isomorphic to L_j.

The following questions remain open:

(a) Is the class K_d characterizable in K_m by c-sublattices?

(b) Does there exist a nontrivial variety $K \neq K_0$ that is characterizable in K_0 by c-sublattices?

References

[1] *G. Birkhoff:* Lattice theory, third edition, Providence 1967.

[2] *P. Crawley, R. P. Dilworth:* Algebraic theory of lattices, New York—London 1973.

[3] *G. Grätzer:* Lectures on lattice theory, Vol. *I*, San Francisco 1971.

[4] *J. Jakubík:* Modular lattices of locally finite length, Acta scient. mathem. (to appear).

[5] *B. Вильгельм:* Двойственное себе ядро условий Биркгофа в структурах с конечными цепами, Czech. Math. J. *5* (1975), 439—450.

Author's address: 040 01 Košice, Švermova 5, ČSSR (VŠT v Košiciach).