Czechoslovak Mathematical Journal

David Lowell Lovelady

On the asymptotic classes of solutions of a superlinear differential equation

Czechoslovak Mathematical Journal, Vol. 27 (1977), No. 2, 242–245

Persistent URL: http://dml.cz/dmlcz/101463

Terms of use:

© Institute of Mathematics AS CR, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 27 (102) 1977, Praha

ON THE ASYMPTOTIC CLASSES OF SOLUTIONS OF A SUPERLINEAR DIFFERENTIAL EQUATION

DAVID LOWELL LOVELADY, Tallahassee

(Received May 8, 1975)

Let n be an integer, $n \geqq 2$, let q be a positive continuous function on $[0, \infty)$, and let α be a real number, $\alpha > 1$. It is known (see I. LIČKO and M. ŠVEC [3] and G. H. RYDER and D. V. V. WEND [6]) that

$$\int_0^\infty t^{2n-1}\, q(t)\, \mathrm{d}t < \infty \tag{1}$$

is a necessary and sufficient condition for the existence of a nonoscillatory solution of

$$u^{(2n)}(t) + q(t)\, |u(t)|^\alpha \operatorname{sgn}(u(t)) = 0\,. \tag{2}$$

The sufficiency of (1) is usually shown by showing the existence of a solution u of (2) with

$$\lim_{t \to \infty} u(t) \tag{3}$$

existing and not being zero. On the other hand, I. T. KIGURADZE [2] has shown that the nonoscillatory solutions of (2) fall into n distinct classes, and one of these classes contains all solutions u for which (3) exists. We shall obtain separate necessary and sufficient conditions for each of the n classes.

Suppose u is an eventually positive solution of (2). Now there is $c \geqq 0$ such that u is defined on $[c, \infty)$ and none of $u, u', \ldots, u^{(2n-1)}$ has a zero in $[c, \infty)$. Let j_u be the largest integer such that $u^{(i)} > 0$ on $[c, \infty)$ if $i \leqq j_u$ (where we write $u = u^{(0)}$). Now j_u is odd and $u^{(k)} u^{(k+1)} < 0$ on $[c, \infty)$ if $j_u \leqq k \leqq 2n - 1$. Since j_u is odd, we see that there are n possible values for the function described by $u \to j_u$, and that the eventually positive solutions of (2) fall into n classes. If u is an eventually negative solution of (2), then $-u$ is an eventually positive solution, so similar analyses apply. (See Kiguradze [2], Ryder and Wend [6], and the present author [4], [5] for details on the above construction).

Theorem. *Suppose k is an odd integer in $[0, 2n]$. Then*

$$\int_0^\infty t^{2n-k+\alpha(k-1)}\, q(t)\, \mathrm{d}t < \infty \tag{4}$$

if and only if there is an eventually positive solution u of (2) with $k = j_u$.

Note that our Theorem is an improvement of [5, Corollary 2], in which it was shown that if $k > 1$, $\alpha > k/(k-1)$, and

$$\int_0^\infty t^{2n-k+\alpha(k-1)-1}\, q(t)\, \mathrm{d}t = \infty$$

then (2) has no eventually positive solution u with $k = j_u$. If m is an odd integer in $[0, 2n]$ and $m < k$ then (4) implies

$$\int_0^\infty t^{2n-m+\alpha(m-1)}\, q(t)\, \mathrm{d}t < \infty\,,$$

since $\alpha > 1$. This gives the following result.

Corollary 1. *If m and k are odd integers in $[0, 2n]$, if $m < k$, and if there is an eventually positive solution u of (2) with $j_u = k$, then there is an eventually positive solution u of (2) with $j_u = m$.*

If u is an eventually positive solution of (2) and $k = j_u$, then $u^{(k)} > 0$ and $u^{(k+1)} < 0$, eventually, so $\lim_{t\to\infty} u^{(k)}(t)$ exists. This and k applications of L'Hôpital's Rule say that $\lim_{t\to\infty} u(t)/t^k$ exists. Thus we have another corollary.

Corollary 2. *Suppose k is an odd integer in $[2, 2n]$. Then*

$$\int_0^\infty t^{2n-k+\alpha(k-1)}\, q(t)\, \mathrm{d}t = \infty \tag{5}$$

if and only if

$$\lim_{t\to\infty} \frac{u(t)}{t^{k-2}} \tag{6}$$

exists and is finite whenever u is a nonoscillatory solution of (2).

Proof of the theorem. Suppose there is an eventually positive solution u of (2) with $k = j_u$. Find $c \geqq 0$ such that u is defined on $[c, \infty)$ and none of $u, u', \ldots, u^{(2n-1)}$ has a zero on $[c, \infty)$. Now

$$u^{(k)}(t) \geqq \frac{1}{(2n-k-1)!} \int_0^\infty (s-t)^{2n-k-1}\, q(s)\, u(s)^\alpha\, \mathrm{d}s \tag{7}$$

if $t \geqq c$. Also, if $s \geqq t \geqq c$,

$$(8)\quad u(s) \geqq \frac{1}{(k-2)!}\int_c^s (s-\xi)^{k-2}\, u^{(k-1)}(\xi)\,\mathrm{d}\xi \geqq \frac{1}{(k-2)!}\int_t^s (s-\xi)^{k-2}\, u^{(k-1)}(\xi)\,\mathrm{d}\xi \geqq$$
$$\geqq \frac{u^{(k-1)}(t)}{(k-2)!}\int_t^s (s-\xi)^{k-2}\,\mathrm{d}\xi = \frac{(s-t)^{k-1}}{(k-1)!}\, u^{(k-1)}(t)\,,$$

since $u^{(k-1)}$ is increasing (recall $u^{(k)} > 0$). If $v = u^{(k-1)}$ and $\beta = ((2n-k-1)!)^{-1}\,.$ $.\,((k-1)!)^{-\alpha}$ then (7) and (8) say

$$v'(t) \geqq \beta\, v(t)^{\alpha} \int_t^{\infty} (s-t)^{2n-k-1+\alpha(k-1)}\, q(s)\,\mathrm{d}s\,,$$

$$v'(t)\, v(t)^{-\alpha} \geqq \beta \int_t^{\infty} (s-t)^{2n-k-1+\alpha(k-1)}\, q(s)\,\mathrm{d}s\,,$$

$$\frac{1}{\alpha-1}\left(v(c)^{1-\alpha} - v(t)^{1-\alpha}\right) \geqq \beta \int_c^t \left(\int_s^{\infty} (\xi-s)^{2n-k-1+\alpha(k-1)}\, q(\xi)\,\mathrm{d}\xi\right)\mathrm{d}s$$

if $t \geqq c$. Since $\lim\limits_{t\to\infty} v(t)^{1-\alpha}$ exists, because $\alpha > 1$, this says

$$(9)\qquad \int_c^{\infty}\left(\int_s^{\infty} (\xi-s)^{2n-k-1+\alpha(k-1)}\, q(\xi)\,\mathrm{d}\xi\right)\mathrm{d}s < \infty\,.$$

But (9) implies (4), so the first part of the proof is complete.

Now suppose (4) holds, and let

$$\gamma = \int_0^{\infty} t^{2n-k+\alpha(k-1)}\, q(t)\,\mathrm{d}t\,.$$

Find positive numbers β and b such that

$$(10)\qquad \beta + \frac{b^{\alpha}\gamma}{(k-1)!\,(2n-k-1)!} \leqq b\,.$$

(Clearly there are such numbers, since $\alpha > 1$.) Let $\mathscr{F}$ be the set to which f belongs if and only if f is a continuous function from $[0, \infty)$ to $[0, \infty)$, and $f(t) \leqq bt^{k-1}$ if $t \geqq 0$. If f is in $\mathscr{F}$ then (4) says

$$\int_0^{\infty} t^{2n-k-1}\, q(t)\, f(t)^{\alpha}\,\mathrm{d}t < \infty\,.$$

Let T be that function on $\mathscr{F}$, each value of which is a continuous function from $[0, \infty)$ to $[0, \infty)$, such that $g = T(f)$ if and only if

$$g(t) = \beta t^{k-1} + \frac{1}{(k-1)!\,(2n-k-1)!}\int_0^t (t-s)^{k-1}\left(\int_s^{\infty} (\xi-s)^{2n-k-1}\, q(\xi)\, f(\xi)^{\alpha}\,\mathrm{d}\xi\right)\mathrm{d}s$$

whenever $t \geqq 0$. Suppose f is in $\mathscr{F}$ and $g = (Tf)$. If $t \geqq 0$,

$$g(t) \leqq \beta t^{k-1} + \frac{b^\alpha}{(k-1)!\,(2n-k-1)!} \int_0^t t^{k-1} \left(\int_s^\infty \xi^{2n-k-1+\alpha(k-1)} q(\xi)\, \mathrm{d}\xi \right) \mathrm{d}s =$$

$$= t^{k-1} \left(\beta + \frac{b^\alpha}{(k-1)!\,(2n-k-1)!} \int_0^t \left(\int_s^\infty \xi^{2n-k-1+\alpha(k-1)} q(\xi)\, \mathrm{d}\xi \right) \mathrm{d}s \right) \leqq$$

$$\leqq \left(\beta + \frac{b^\alpha \gamma}{(k-1)!\,(2n-k-1)!} \right) t^{k-1} \leqq b t^{k-1},$$

from (10), of g is in $\mathscr{F}$, and T maps $\mathscr{F}$ into $\mathscr{F}$. Now routine computations show that T is continuous with respect to the topology of locally uniform convergence, and that the range of T is locally equicontinuous. Thus the fixed point theorem of J. SCHAUDER [7] (see also W. A. COPPEL [1, p. 9]) says that there is u in $\mathscr{F}$ with $u = T(u)$, i.e.,

$$u(t) = \beta t^{k-1} + \tag{11}$$

$$+ \frac{1}{(k-1)!\,(2n-k-1)!} \int_0^t (t-s)^{k-1} \left(\int_s^\infty (\xi - s)^{2n-k-1} q(\xi)\, u(\xi)^\alpha \, \mathrm{d}\xi \right) \mathrm{d}s$$

if $t \geqq 0$. Now (11) says $u(t) \geqq \beta t^{k-1}$ if $t \geqq 0$, so $u(t)$ is positive if $t \geqq 0$. Also, it is easily seen that u is a solution of (2), and $j_u = k$. The proof is complete.

Corollaries 1 and 2 are now obvious.

References

[1] *W. A. Coppel:* Stability and asymptotic behavior of differential equations, D. C. Heath & Co., Boston, 1965.

[2] *I. T. Kiguradze:* Oscillatory properties of certain ordinary differential equations, Soviet Math. Dokl., *3* (1962), 649—652 (Russian).

[3] *I. Ličko* and *M. Švec:* Le caractère oscillatoire des solutions de l'équation $y^{(n)} + f(x)\, y^\alpha = 0$, $n > 1$, Czechoslovak Math. J. *13*, (*88*), (1963), 481—491.

[4] *D. L. Lovelady:* Oscillation and even order linear differential equations, Rocky Mountain Math. J., *6* (1976), 299—304,

[5] *D. L. Lovelady:* Nonlinear even order oscillation criteria according to Kiguradze classes, (unpublished).

[6] *G. H. Ryder* and *D. V. V. Wend:* Oscillation of solutions of certain ordinary differential equations of nth order, Proc. Amer. Math. Soc., *25* (1970), 463—469.

[7] *J. Schauder:* Der Fixpunktsatz in Funktionalräumen, Studia Math., *2* (1930), 171—180.

Author's address: Department of Mathematics, The Florida State University, Tallahassee, Florida 32306, U. S. A.