Czechoslovak Mathematical Journal

Ivan Lončar

Relative continuity of the functor β

Czechoslovak Mathematical Journal, Vol. 37 (1987), No. 4, 517–521

Persistent URL: http://dml.cz/dmlcz/102179

Terms of use:

© Institute of Mathematics AS CR, 1987

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 37 (112) 1987, Praha

RELATIVE CONTINUITY OF THE FUNCTOR β

IVAN LONČAR, Varaždin

(Received May 11, 1984)

0. INTRODUCTION AND PRELIMINARY RESULTS

0.1. The set of positive integers is denoted by N. The symbol ω_m denotes the set of all ordinals of the cardinality $\aleph_{m-1}$, $m \geqq 1$.

0.2. We use the notion of the inverse systems as in [2]. The inverse system is denoted by $\boldsymbol{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ and its limit by $X = \lim \boldsymbol{X}$.

0.3. We say that a covariant functor $F: \mathscr{A} \to \mathscr{B}$ is continuous [3; p. 258] if for every inverse system $\boldsymbol{X} = \{X_\alpha, f_{\alpha\beta}, A\}$, the object $F(\varprojlim \boldsymbol{X})$ is a limit of the inverse system $F\boldsymbol{X} = \{F(X_\alpha). F(f_{\alpha\beta}), A\}$.

It is well known that the Čech-Stone functor β is not continuous. It suffices to consider an inverse system with the empty limit space.

A covariant functor $F: \mathscr{A} \to \mathscr{B}$ is said to be *relatively continuous* with respect to $\boldsymbol{X} = \{X_\alpha, f_{\alpha\beta}, A\}$, or $\boldsymbol{X}$-continuous, if the object $F(\varprojlim \boldsymbol{X})$ is a limit of the inverse system $F\boldsymbol{X} = \{F(X_\alpha), F(f_{\alpha\beta}), A\}$.

The following two theorems were proved by Nagata [10].

0.4. Theorem. *Let $\boldsymbol{X} = \{X_n, f_{nm}, N\}$ be an inverse sequence with a countably compact limit X. If X_n, $n \in N$, are normal firstcountable spaces, then $\beta X = \varprojlim \{\beta X_n, \beta f_{nm}, N\}$.*

0.5. Theorem. *Let $\boldsymbol{X} = \{X_n, f_{nm}, N\}$ be an inverse sequence of normal spaces X_n and open mappings f_{nm}; $n, m \in N$. If $X = \varprojlim \boldsymbol{X}$ is pseudocompact, then $\beta X = \varprojlim \{\beta X_n, \beta f_{nm}, N\}$.*

1. THEOREMS

We start with the following key lemma.

1.1. Lemma. *Let $\boldsymbol{X} = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system of completely regular spaces X_α. Then:* a) *$\beta\boldsymbol{X} = \{\beta X_\alpha, \beta f_{\alpha\beta}, A\}$ is an inverse system;* b) *if the projections $f_\alpha: \varprojlim \boldsymbol{X} \to X_\alpha$, $\alpha \in A$, are onto mappings, then $\varprojlim \beta\boldsymbol{X}$ is the compactification of*

Supported by SIZ for Scientific Work of SRH.

the space $\varprojlim X$. *Moreover, the spaces* $\varprojlim \beta X$ *and* $\beta(\varprojlim X)$ *are homeomorphic iff for every pair* F_1, F_2 *of completely separated subsets of* $\varprojlim X$ *there exists* $\alpha \in A$ *such that* $f_\alpha(F_1)$ *and* $f_\alpha(F_2)$ *are completely separated subsets of* X_α.

Proof. The proof of a) is trivial and can be omitted. For the first assertion of b) it suffices to prove that $\varprojlim \beta X$ is an extension of $\varprojlim X$ (since $\varprojlim \beta X$ is compact). For every nonempty open set $U \subseteq \varprojlim \beta X$ there is a nonempty open $U_\alpha \subseteq \beta X_\alpha$ such that $f_\alpha^{-1}(U_\alpha) \subseteq U$. The set $V_\alpha = U_\alpha \cap X_\alpha$ is nonempty since X_α is dense in βX_α. The surjectivity of f_α implies that $f_\alpha^{-1}(V_\alpha)$ is a nonempty subset of $\varprojlim X$. Clearly, $f_\alpha^{-1}(V_\alpha) \subseteq U$. This means that $\varprojlim X$ is dense in $\varprojlim \beta X$.

We prove now the second part of b).

The "if" part. From the assumption $f_\alpha(F_1) \cap f_\alpha(F_2) = \emptyset$ it follows that $f_\alpha(F_1)^{\beta X_\alpha} \cap \cap f_\alpha(F_2)^{\beta X_\alpha} = \emptyset$ [4; p. 226]. It follows that $\bar{F}_1^{X'} \cap \bar{F}_2^{X'} = \emptyset$, $X' = \varprojlim \beta X$ since $\operatorname{cl} F_i^{X'} \subseteq (\beta f_\alpha)^{-1} \operatorname{cl}(f_\alpha(F_i))^{\beta X_\alpha}$, $i = 1, 2$. From [4: p. 226] it follows that X' is homeomorphic to $\beta(\varprojlim X)$.

The "only if" part. Now, from $\operatorname{cl} F_1^{X'} \cap \operatorname{cl} F_2^{X'} = \emptyset$ it follows that there exists $\alpha \in A$ such that $\operatorname{cl}(f_\alpha(F_1)^{\beta X_\alpha} \cap \operatorname{cl}(f_\alpha(F_2))^{\beta X_\alpha} = \emptyset$ since βX is an inverse system of compact spaces βX_α. From the normality of βX_α it follows that $\operatorname{cl}(f_\alpha(F_1)^{\beta X_\alpha}$ and $\operatorname{cl}(f(F_2)^{\beta X_\alpha}$ are completely separated. This means that $f_\alpha(F_1)$ and $f_\alpha(F_2)$ are completely separated. The proof is complete.

1.2. Remark. *If* X_α, $\alpha \in A$, *are normal spaces, then* $\beta(\varprojlim X)$ *and* $\varprojlim \beta X$ *are homeomorphic iff for every pair* F_1, F_2 *of completely separated subsets of* $\lim X$ *there exists* $\alpha \in A$ *such that* $\operatorname{cl}(f_\alpha(F_1))^{X_\alpha'} \cap \operatorname{cl}(f_\alpha(F_2))^{X_\alpha'} = \emptyset$.

If $\varprojlim X$ is normal, then we can assume that F_1 and F_2 are closed subsets of $\varprojlim X$.

Applying Lemma 1.1 and Remark 1.2 we prove the following theorems.

1.3. Theorem. *Let* $X = \{X_n, f_{nm}, N\}$ *be an inverse sequence such that the mappings* f_{nm} *are onto and the spaces* X_n *are normal. If* $X = \varprojlim X$ *is a countably compact space, then* $\beta X = \varprojlim \beta X$.

Proof. Let us prove that X is normal. Consider two disjoint closed subsets F_1, F_2 of X. If we assume that $Y_n = \operatorname{cl}(f_n(F_1))^{X_n} \cap \operatorname{cl}(f_n(F_2))^{X_n}$ is nonempty for every $n \in N$, then we obtain a contradiction: $\emptyset \neq \bigcap\{f_n^{-1}(Y_n): n \in N\} \subseteq F_1 \cap F_2 = \emptyset$ since the space X is countably compact. It follows that there exists $n_0 \in N$ such that $\operatorname{cl}(f_{n_0}(F_1))^{X_{n_0}} \cap \cap \operatorname{cl}(f_{n_0}(F_2))^{X_{n_0}} = \emptyset$. Since X_{n_0} is normal, it follows that there exist disjoint open sets $U \supseteq \operatorname{cl}(f_{n_0}(F_1))^{X_{n_0}}$ and $V \supseteq \operatorname{cl}(f_{n_0}(F_2))^{X_{n_0}}$. Clearly, $f_{n_0}^{-1}(U) \supseteq F_1$ and $f_{n_0}^{-1}(V) \supseteq \supseteq F_2$. The normality of X is proved.

Let $X' = \varprojlim \beta X$. For two disjoint closed subsets $F_1, F_2 \subseteq X$ we have $n_0 \in N$ such that $\operatorname{cl}(f_{n_0}(F_1))^{X_{n_0}} \cap \operatorname{cl}(f_{n_0}(F_2))^{X_{n_0}} = \emptyset$. Lemma 1.1 and Remark 1.2 imply that $\beta X = \varprojlim \beta X$. The proof is complete.

1.4. Corollary. *Let* $X = \{X_n, f_{nm}, N\}$ *be an inverse sequence of normal countably compact spaces* X_n. *If* f_{nm}, $n, m \in N$, *are closed onto mappings, then* $\beta(\varprojlim X) = = \varprojlim \beta X$.

Proof. In [8] it is proved that $\underline{\lim}\, X$ is a normal countably compact space. Now, apply Theorem 1.3.

1.5. Remark. Corollary 1.4 implies Theorem 0.4 since a continuous mapping $f: X \to Y$ is closed if X is countably compact, Y is a regular first-countable space, and for every inverse system there exists another one which has onto projections f'_α.

We say that a mapping $f: X \to Y$ is fully closed [6] if for every point $y \in Y$ and every finite cover $\{U_1, \ldots, U_s\}$ of $f^{-1}(y)$ by open sets $U_i \subseteq X$, $i = 1, \ldots, s$, the set $\{y\} \cup \cup (f^\# U_1 \cup \ldots f^\# U_s\}$ is an open set in Y. The set $f^\# U_i$ is defined by $f^\# U_i = \{y: f^{-1}(y) \subseteq \subseteq U_i\}$.

1.6. Theorem. *Let $X = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse system such that $f_{\alpha\beta}$ are perfect fully closed onto mappings. If the spaces X_α are normal countably compact spaces, then $X = \lim X$ is normal and $\beta X = \underline{\lim}\, \beta X$.*

Proof. The projections $f_\alpha: X \to X_\alpha$, $\alpha \in A$, are perfect fully closed onto mappings [6; Lemma 3]. It is readily seen that X is countably compact. On the other hand, for every pair F_1, F_2 of disjoint closed subsets of X, the sets $Y_\alpha = f_\alpha(F_1) \cap f_\alpha(F_2)$, $\alpha \in A$, are discrete [6; Lemma 1(b)]. This means that Y_α, $\alpha \in A$, are finite sets since X_α, $\alpha \in A$, are countably compact. It follows that $Y = \{Y_\alpha, f_{\alpha\beta}/Y_\beta, A\}$ has a non-empty limit Y, if all Y_α are nonempty. The contradiction $\emptyset \neq Y \subseteq F_1 \cap F_2 = \emptyset$ implies that there exists $\alpha \in A$ such that $Y_\alpha = f_\alpha(F_1) \cap f_\alpha(F_2) = \emptyset$. This fact yields that X is normal. Moreover, the conditions of Lemma 1.1 and Remark 1.2 are satisfied. The proof is complete.

For a sequentially compact (strongly countably compact, D-compact) spaces [8; p. 158] we prove

1.7. Theorem. *If $X = \{X_n, f_{nm}, N\}$ is an inverse system of normal sequentially compact (strongly countably compact, D-compact) spaces X_n, then $X = \lim X$ is a normal space and $\beta X = \underline{\lim}\, \beta X$.*

Proof. X is a countably compact space [4; p. 268] ([8; p. 164]). Applying Theorem 1.3 we complete the proof.

We say that an inverse system $X = \{X_\alpha, f_{\alpha\beta}, A\}$ is a factorizable, or f-system [12], if for every real-valued function $f: \lim X \to R$ there exist $\alpha \in A$ and a real-valued function $g_\alpha: X_\alpha \to X$ such that $f = g_\alpha f_\alpha$, where $f_\alpha: \lim X \to X_\alpha$ is the projection.

1.8. Remark. If X is a limit space of an inverse f-system $X = \{X_\alpha, f_{\alpha\beta}, A\}$ of pseudocompact spaces X_α, then X is pseudocompact.

1.9. Theorem. *If X is a limit space of an inverse f-system with surjective projections $f_\alpha: X \to X_\alpha$, $\alpha \in A$, then $\beta X = \underline{\lim}\, \beta X$.*

Proof. Let $X = \{X_\alpha, f_{\alpha\beta}, A\}$ be an inverse f-system and $f: X \to I$ a continuous real-valued function. There exist $\alpha \in A$ and $g_\alpha: X_\alpha \to I$ such that $f = g_\alpha f_\alpha$. Let

f'_α: $\varprojlim \beta X \to \beta X_\alpha$ be the projection. The function $f' = (\beta g_\alpha) \cdot f'_\alpha$ is an extension of f. By virtue of [4; 3.6.3 Corollary] it follows that $\varprojlim \beta X$ is equivalent to the compactification βX.

1.10. Corollary. *Let $X = \{X_\alpha, f_{\alpha\beta}, A\}$ be a σ-directed inverse system such that the projections f_α: $\varprojlim X \to X_\alpha$, $\alpha \in A$, are onto mappings. If $\lim X$ is a Lindelöf space, then $\beta(\varprojlim X) = \varprojlim \beta X$.*

Proof. X is an f-system [12; pp. 28]. Theorem 1.9 completes the proof.

1.11. Corollary. *Let X be a limit space $X = \{X_\alpha, f_{\alpha\beta}, A\}$ of an m-directed inverse system, where the Souslin number $c(X_\alpha) \leqq m$. If the projections f_α: $\lim X \to X_\alpha$, $\alpha \in A$, are onto mappings, then $\beta(\lim X) = \varprojlim \beta X$.*

Proof. X is an f-system [12; pp. 28, Predloženije 1.8]. Now, apply Theorem 1.9.

1.12. Corollary. *Let $X = \{X_\alpha, f_{\alpha\beta}, \Omega\}$ be a well-ordered inverse system of hereditarily Lindelöf spaces X_α such hat $\mathrm{cf}(\Omega) > \omega_1$. If the projections f_α: $\varprojlim X \to X_\alpha$, $\alpha \in A$, are onto mappings, then $\beta(\varprojlim X) = \varprojlim \beta X$.*

Proof. By virtue of [13] it follows that $\lim X$ is a hereditarily Lindelöf space. Corollary 1.10 implies that $\beta(\varprojlim X) = \varprojlim \beta X$. The proof is complete.

We close this paper with the following theorem.

1.13. Theorem. *Let $X = \{X_\alpha, f_{\alpha\beta}, \Omega\}$ be an inverse system, where the spaces X_α are normal and such that the hereditary Lindelöf number $\mathrm{hl}(X_\alpha) < \aleph_m$ and $\mathrm{cf}(\Omega) > \aleph_{m+1}$. If the projections f_α: $\varprojlim X \to X_\alpha$, $\alpha \in A$, are onto mappings, then $\beta(\varprojlim X) = \varprojlim \beta X$.*

Proof. From [9] and [13] it follows that for every pair F_1, F_2 of disjoint closed subsets of $\lim X$ there exists $\alpha \in \Omega$ such that $F_i = f_\alpha^{-1}\,\mathrm{cl}(f_\alpha(F))$, $i = 1, 2$. This means that $\varprojlim X$ is a normal space and that the conditions of Lemma 1.1 and Remark 1.2 are satisfied. The proof is complete.

References

[1] *Arhangelskij A. V.* and *Ponomarev V. I.:* Osnovy obščej topologii v zadačah i upražnenijah, Nauka, Moskva, 1974.

[2] *Bucur I.* and *Deleanu A.:* Vvedenije v teoriju kategorij i funktorov, Mir, Moskva, 1972.

[3] *Eilenberg S.* and *Steenrod N.:* Foundations of algebraic topology, Princeton, 1952.

[4] *Engelking R.:* General Topology, PWN, Warszawa, 1977.

[5] *Engelking R.:* Remarks on real-compact spaces, Fund. Math. *55* (1964), 303—308.

[6] *Fedorčuk V. V.:* Metod razvertyvaemyh spektrov i vpolne zamknutyh otobraženij v obščej topologii, UMN *35* (1980), 112—121.

[7] *Gillman L.* and *Jerison M.:* Rings of continuous functions, D. van Nostrand co., New York, 1960.

[8] *Lončar I.:* Inverse limits which generalize compact spaces, Glasnik matematički *17* (*37*) (1982), 155—173.

[9] *Lonačar I.:* Lindelöfov broj i inverzni limesi, Zbornik radova Fakulteta organizacije i informatike Varaždin *6* (1982), 155—165.
[10] *Nagota Jun-iti:* Some aspects of extension theory in general topology, Contribs. Extens. Theory. Topol. Struct. Proc. Sympos. Berlin 1967, Berlin 1969, 157—161.
[11] *Semadeni Z.:* Banach spaces of continuous functions I, PWN, Warszawa, 1971.
[12] *Ščepin E. V.:* Funktory i nesčetnye stepeni kompaktov, UMN *36* (1981), 3—62.
[13] *Tkačenko M. G.:* Cepy i kardinaly, DAN SSSR *239* (1978), 546—549.

Author's address: Fakultet organizacije i informatike, Varaždin, Yugoslavia.