Czechoslovak Mathematical Journal

Francesco Catino

On complete α-ideals in semigroups

Czechoslovak Mathematical Journal, Vol. 40 (1990), No. 1, 155–158

Persistent URL: http://dml.cz/dmlcz/102366

Terms of use:

© Institute of Mathematics AS CR, 1990

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 40 (115) 1990, Praha

ON COMPLETE α-IDEALS IN SEMIGROUPS

FRANCESCO CATINO, Lecce*)

(Received May 30, 1988)

By X^* we denote the free monoid over an alphabet X. Let S be a semigroup. By $\mathscr{P}(S)$ we denote the semigroup of all subset of S under set product with the unity $\emptyset$. For $\alpha \in \{0, 1\}^*$ we shall define $f_\alpha^S: \mathscr{P}(S) \to \mathscr{P}(S)$ as follows: $f_\alpha^S(A) = A_1 A_2 \dots A_n$ if $\alpha = \alpha_1 \alpha_2 \dots \alpha_n$, $\alpha_i \in \{0, 1\}$, where

$$A_i = \begin{cases} A & \text{for } \alpha_i = 1 \\ S & \text{for } \alpha_i = 0 \end{cases}$$

We put $\Lambda = \{0, 1\}^* \setminus \{1\}^*$ and $f_\alpha := f_\alpha^S$. Let $\alpha \in \Lambda$. A subsemigroup M of a semigroup S is called an *α-ideal* of S if $f_\alpha(M) \subseteq M$ (see [6]). If, especially $f_\alpha(M) = M$, then we say that M is a *complete* α-ideal of S.

In this note we shall characterize the semigroups in which every α-ideal is complete.

For the undefined notions and notations we refer to [1].

It is well known (see [6]) the following

Lemma 1. *Let $\alpha, \beta \in \Lambda$ and A, B non-empty subsets of a semigroup S.*

(i) *If $A \subseteq B$ then $f_\alpha(A) \subseteq f_\alpha(B)$;*

(ii) $f_{\alpha\beta}(A) = f_\alpha(A) f_\beta(A)$;

(iii) $A f_\alpha(A) \subseteq f_\alpha(A)$ *and* $f_\alpha(A) A \subseteq f_\alpha(A)$;

(iv) $f_\alpha(A) f_\alpha(A) \subseteq f_\alpha(A)$;

(v) $f_\alpha(A \cup f_\alpha(A)) \subseteq f_\alpha(A)$.

The following lemma is an extension of Lemma 1 (v).

Lemma 2. *Let $\alpha \in \Lambda$ and k a positive integer. If A is a non-empty subset of a semigroup S, then*

$$f_\alpha(A \cup A^2 \cup \dots \cup A^k \cup f_\alpha(A)) \subseteq f_\alpha(A) .$$

Proof. At first we prove that

$$f_\alpha(A \cup A^2 \cup \dots \cup A^k) \subseteq f_\alpha(A) \tag{1}$$

by induction on the length $l(\alpha)$ of α. It clear that the result is true for $l(\alpha) = 1$. Assume now that $l(\alpha) \geqq 2$ and the result holds for every word β such that $l(\beta) =$

*) Research supported by a grant from M.P.I. (Italy)

$= l(\alpha) - 1$.

Case 1: $\alpha = 1\beta$. Then $\beta \in \Lambda$ and $f_\beta(A \cup \ldots \cup A^k) \subseteq f_\beta(A)$. Using Lemma 1 (ii), (iii)

$$\begin{aligned} f_\alpha(A \cup A^2 \cup \ldots \cup A^k) &= (A \cup A^2 \cup \ldots \cup A^k) f_\beta(A \cup A^2 \cup \ldots \cup A^k) \subseteq \\ &\subseteq (A \cup A^2 \cup \ldots \cup A^k) f_\beta(A) \subseteq A f_\beta(A) \cup A^2 f_\beta(A) \cup \ldots \cup A^k f_\beta(A) = \\ &= A f_\beta(A) \cup A(A f_\beta(A)) \cup \ldots \cup A^{k-1}(A f_\beta(A)) \subseteq \ldots \subseteq A f_\beta(A) = \\ &= f_{1\beta}(A) = f_\alpha(A)\,. \end{aligned}$$

Case 2: $\alpha = 0\beta$. If $\beta \in \Lambda$,

$$f_\alpha(A \cup A^2 \cup \ldots \cup A^k) = S f_\beta(A \cup A^2 \cup \ldots \cup A^k) \subseteq S f_\alpha(A) = f_\alpha(A)\,.$$

If $\beta \notin \Lambda$, $\alpha = \gamma 1$ with $\gamma \in \Lambda$. This is dual to case 1.

Now, by Lemma 1(i), (v) and by (1),

$$\begin{aligned} &f_\alpha(A \cup A^2 \cup \ldots \cup A^k \cup f_\alpha(A)) \subseteq \\ &\subseteq f_\alpha(A \cup A^2 \cup \ldots \cup A^k \cup f_\alpha(A \cup A^2 \cup \ldots \cup A^k)) \subseteq \\ &\subseteq f_\alpha(A \cup A^2 \cup \ldots \cup A^k) \subseteq f(A)\,. \end{aligned}$$

Theorem 3. *Let A be a non-empty subset of a semigroup S. Then $A \cup A^2 \cup \ldots \cup A^{l(\alpha)-1} \cup f_\alpha(A)$ is the smallest α-ideal of S containing A, where $l(\alpha)$ is the length of α.*

Proof. Since $(A \cup A^2 \cup \ldots \cup A^{l(\alpha)-1} \cup f_\alpha(A))\ (A \cup A^2 \cup \ldots \cup A^{l(\alpha)-1} f_\alpha(A)) \subseteq$
$\subseteq A^2 \cup A^3 \cup \ldots \cup A^{l(\alpha)-1} \cup A^{l(\alpha)} \cup A f_\alpha(A) \cup (A^{l(\alpha)+1} \cup A^2 f(A)) \cup \ldots$
$\ldots \cup (A^{2l(\alpha)-2} \cup A^{l(\alpha)-1} f_\alpha(A)) \cup (f_\alpha(A) A \cup \ldots \cup f_\alpha(A) A^{l(\alpha)-1}) \cup f_\alpha(A) f_\alpha(A) \subseteq$
$\subseteq A^2 \cup A^3 \cup \ldots \cup A^{l(\alpha)-1} \cup \ldots \cup A^{2l(\alpha)-2} \cup f_\alpha(A) \subseteq A^2 \cup \ldots \cup A^{l(\alpha)-1} \cup f_\alpha(A)$
we see that $A \cup A^2 \cup \ldots \cup A^{l(\alpha)-1} \cup f_\alpha(A)$ is a subsemigroup of S containing A. By Lemma 2 it is an α-ideal of S. Moreover, let B be an α-ideal of S containing A. Then $A \cup \ldots \cup A^{l(\alpha)-1} \cup f_\alpha(A) \subseteq B \cup f_\alpha(B) \subseteq B$.

If A is a non-empty subset of a semigroup S then $A \cup A^2 \cup \ldots \cup A^{l(\alpha)-1} \cup f_\alpha(A)$ is called the *α-ideal generated by A* and denoted by $(A)_\alpha$, where $l(\alpha)$ is the length of α.

Let $\alpha \in \Lambda$. A semigroup S is called *α-semigroup* if every α-ideal of S is complete.

Theorem 4. *Let $\alpha \in \Lambda$. A semigroup S is an α-semigroup if and only if $a \in f_\alpha(\{a\})$ for all $a \in S$.*

Proof. Suppose that every α-ideal of S is complete. Let $a \in S$. Then, by Theorem 3 and Lemma 2, $a \in \{a\} \cup \ldots \cup \{a^{l(\alpha)-1}\} \cup f_\alpha(\{a\}) = f_\alpha(\{a\} \cup \ldots \cup \{a^{l(\alpha)-1}\} \cup f_\alpha(\{a\})) \subseteq$
$\subseteq f_\alpha(\{a\})$.

Conversely, if A is an α-ideal of S. Then, by Lemma 1(i), $a \in f_\alpha(\{a\}) \subseteq f_\alpha(A)$ for all $a \in A$. Hence A is a complete α-ideal of S.

Remarks. 1) If $\alpha = 10$ then any α-ideal of S is a left ideal of S. By Theorem 4, every left ideal of S is complete iff $a \in aS$ for all $a \in S$ (see [2]).

2) If $\alpha = 101$ then any α-ideal of S is a bi-ideal of S. By Theorem 4, every bi-ideal of S is complete iff S is a regular semigroup (see [7]).

3) If $\alpha = 1011$ then any α-ideal of S is a (1,2)-ideal of S (see [3]). By Theorem 4, every (1,2)-ideal of S is complete iff S is a completely regular semigroup (see [4]).

Now, we remark that in general the intersection of complete α-ideals of a semigroup S need not be complete (see [2]).

Theorem 5. *Let $\alpha \in \Lambda$. If A, B are complete α-ideals of a semigroup S, then $(A \cup B)_\alpha$ is a complete α-ideal of S.*

Proof. By Lemma 1(i), $A \cup B = f_\alpha(A) \cup f_\beta(B) \subseteq f_\alpha(A \cup B)$. Thus, by Lemma 1(iii), $(A \cup B)^k \subseteq f_\alpha(A \cup B)$ for all positive integers k. Hence $(A \cup B)_\alpha = (A \cup B) \cup \cup (A \cup B)^2 \cup \ldots \cup (A \cup B)^{l(\alpha)-1} \cup f_\alpha(A \cup B) \subseteq f_\alpha(A \cup B) \subseteq f_\alpha((A \cup B)_\alpha)$.

It is clear that the product of two α-ideals of a semigroup, in general, will not be an α-ideal (see [6]). We will prove that the product of two complete α-ideals of a semigroup is an α-ideal when $\alpha \in \Lambda$ and α containing *a* 1. But we first need some auxiliary statements.

Let $\alpha \in \Lambda$. If $\alpha = 0^n$ for some n positive integer, we put $\alpha^* = 0$. If $\alpha = \alpha_1\alpha_2 \ldots \alpha_n$ and there exists $j > 1$ such that $\alpha_j \neq \alpha_1$ we put

$$\alpha^* = \begin{cases} \alpha_1\alpha_j\alpha_n & \text{if} \quad \alpha_n \neq \alpha_j \\ \alpha_1\alpha_j & \text{if} \quad \alpha_n = \alpha_j \end{cases}$$

We remark that $\alpha^* \in \{0, 10, 01, 101, 010\}$.

Lemma 6. *Let $\alpha \in \Lambda$ and let A, B be complete α-ideals of a semigroup S. Then $f_\alpha(AB) \subseteq f_{\alpha^*}(AB)$.*

Proof. Let $\alpha = \alpha_1\alpha_2 \ldots \alpha_n$. If $\alpha_1 = 1$ then there exists $j > 1$ such that $\alpha_j = 0$. Thus $f_\alpha(AB) \subseteq ABf_{\alpha_2\ldots\alpha_{j-1}}(AB)\, S f_{\alpha_{j+1}\ldots\alpha_n}(AB) \subseteq AB\, S f_{\alpha_n}(AB)$.

If $\alpha_n = 0$ then $f_\alpha(AB) \subseteq ABS = f_{\alpha^*}(AB)$. If $\alpha_n = 1$ then $f_\alpha(AB) \subseteq ABSAB = = f_{\alpha^*}(AB)$.

Now, we suppose that $\alpha_1 = 0$. If $\alpha_j = 0$ for all $j \in \{2, \ldots, n\}$ then evidently $f_\alpha(AB) \subseteq S = f_{\alpha^*}(AB)$. If $\alpha_n = 1$ for some $h > 1$, then $f_\alpha(AB) \subseteq SABf_{\alpha_{h+1}\ldots\alpha_n}(AB) \subseteq \subseteq SABSf_{\alpha_n}(AB)$ if there is r such that $h + 1 \leqq r < n$ and $\alpha_r = 0$ or $f_\alpha(AB) \subseteq \subseteq SABf_{\alpha_n}(AB)$. Thus $f_\alpha(AB) \subseteq SABS = f_{\alpha^*}(AB)$ if $\alpha_n = 0$ and $f_\alpha(AB) \subseteq SABAB \subseteq \subseteq SAB = f_{\alpha^*}(AB)$ if $\alpha_n = 1$.

Lemma 7. *Let $\alpha \in \Lambda$, $\alpha^* \neq 0$ and let A, B be complete α-ideals of a semigroup S. Then $f_{\alpha^*}(AB) \subseteq AB$.*

Proof. If $\alpha^* = 01$ then $\alpha = 0\alpha'$ with $\alpha' \in \{0, 1\}^*$. Thus $f_{\alpha^*}(AB) = SAB = = S f_\alpha(A)\, B = SS f_{\alpha'}(A)\, B \subseteq S f_{\alpha'}(A)\, B \subseteq f_\alpha(A)\, B = AB$. The case $\alpha^* = 10$ is dual to the preceding case. If $\alpha^* = 010$, then $f_{\alpha^*}(AB) = SABS = S f_\alpha(A) f_\alpha(B)\, S \subseteq \subseteq f_\alpha(A) f_\alpha(B) = AB$. If $\alpha^* = 101$, then $f_{\alpha^*}(AB) = ABSAB \subseteq ABSB = = A f_\alpha(B)\, S f_\alpha(B) \subseteq A f_{\alpha 0 \alpha}(B) \subseteq A f_{\alpha_1}(B) f_{\alpha_2}(B) \ldots f_{\alpha_j}(B) \ldots f_{\alpha_n}(B)\, S f_{\alpha_1}(B) \ldots \ldots f_{\alpha_j}(B) \ldots f_{\alpha_n}(B) \subseteq A f_\alpha(B) = AB$, where $\alpha = \alpha_1\alpha_2 \ldots \alpha_n$.

Lemma 8. *Let $\alpha \in \Lambda$, $\alpha^* \neq 0$ and let A be a complete α-ideal of a semigroup S. Then $f_{\alpha^*}(A) = A$.*

Proof. If $\alpha^* = 010$ then $f_{\alpha^*}(A) = SAS = S f_\alpha(A) S \subseteq f_\alpha(A)$. The cases $\alpha^* \in \{01, 10\}$ are similar. If $\alpha^* = 101$ then $f_{\alpha^*}(A) = ASA = f_\alpha(A) S f_\alpha(A) = f_{\alpha 0 \alpha}(A) \subseteq f_\alpha(A)$.

Moreover, by the proof of Lemma 6, $f_\alpha(A) \subseteq f_{\alpha^*}(A)$. Thus $A = f_{\alpha^*}(A)$.

Theorem 9. *Let $\alpha \in \Lambda$, $\alpha^* \neq 0$. The product of two complete α-ideals A, B of a semigroup S is an α-ideal.*

Proof. We prove that AB is a subsemigroup of S. If $\alpha^* = 010$ then, by Lemma 8, $ABAB = ABASBS \subseteq ASASBS \subseteq ASBS \subseteq AB$. We proceed similarly if $\alpha^* \in \{01, 10, 101\}$.

Furthermore, by Lemma 6 and Lemma 7, $f_\alpha(AB) \subseteq AB$. Thus AB is an α-ideal of S.

We remark that if $\alpha \in \Lambda$ and $\alpha^* \neq 0$, in general, the product of two complete α-ideals will not be complete and if $\alpha^* = 0$, the product of two complete α-ideals need not be an α-ideal.

Let $\alpha \in \Lambda$, $\alpha^* \neq 0$ and let S be an α-semigroup. Let F_α be the set of all α-ideals of S. By Theorem 9 F_α is a semigroup respect to the set product.

Theorem 10. *Let $\alpha \in \Lambda$, $\alpha^* \neq 0$. A semigroup S is an α-semigroup if and only if F_α is an α-semigroup.*

Proof. Let S be an α-semigroup and let A be an element of F_α. Then $A = f_\alpha(A) \in f_\alpha^{F_\alpha}(\{A\})$. Thus, by Theorem 4, F_α is an α-semigroup. Conversely, if $A \in F_\alpha$, by Theorem 4, $A \in f_\alpha^{F_\alpha}(\{A\})$. Hence $A \subseteq f_\alpha(A)$.

Remarks. 1) If $\alpha = 101$, then by Theorem 10, S is regular iff the semigroup of all bi-ideal of S is regular (see [7]).

2) If $\alpha = 1011$, then by Theorem 10, S is completely regular iff the semigroup of all (1,2)-ideals of S is completely regular (see [5]).

References

[1] *Clifford, A. H.* and *G. B. Preston:* The algebraic theory of semigroups, vol. I, Amer. Math. Soc., Providence, 1961.

[2] *Fabrici, I.:* On complete ideals in semigroups, Mat. Časopis Sloven. Akad. Vied *18* (1968), 34—39.

[3] *Lajos, S.:* Generalized ideals in semigroups, Acta Sci. Math., *22* (1961), 217—222.

[4] *Lajos, S.:* On the characterization of completely regular semigroups, Math. Japonicae (Special Issue) *20* (1975), 33—35.

[5] *Lajos, S.:* Some characterizations of completely regular semigroups, Kyungpook Math. J. *19* (1979), 213—214.

[6] *Miccoli M. M.* and *B. Pondělíček:* On α-ideals and generalized α-ideals in semigroups (to appear).

[7] *Steinfeld, O.:* Quasi-ideals in rings and semigroups, Akadémiai Kiadó, Budapest, 1978.

Author's address: Dipartimento di Matematica, Universita' di Lecce, 72100 Lecce, Italy.