Applications of Mathematics

Maia Koicheva

A characterization of the gamma distribution in terms of conditional moment

Applications of Mathematics, Vol. 38 (1993), No. 1, 19--22

Persistent URL: http://dml.cz/dmlcz/104531

Terms of use:

© Institute of Mathematics AS CR, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://project.dml.cz

A CHARACTERIZATION OF THE GAMMA DISTRIBUTION IN TERMS OF CONDITIONAL MOMENT

MAIA KOICHEVA, Brussels

(Received December 21, 1991)

Summary. A characterization of the Gamma distribution in terms of the k-th conditional moment presented in this paper extends the result of Shunji Osaki and Xin-xiang Li (1988).

Keywords: characterization of distribution, Gamma distribution, conditional moment

AMS classification: 60E99

1. INTRODUCTION

The problem of characterization of statistical distributions is of great interest because of its applications in reliability theory and statistics.

The books by Kagan, Linnik, Rao [3] and by Galambos and Kotz [2] were the basis of an increasing interest for research in this area. Most of the results deal with the exponential distribution (see e.g. [1], [2], [3]).

As the Gamma distribution is often used to model a wide variety of positive valued random unities in reliability theory and queueing system analysis, the characterization of this distribution is an appealing and interesting problem with a great number of applications. There are several characterizations of the Gamma distributions (see e.g. [3], [4], [5] and [6]). In this paper, we give a new one based on a relationship between the k-th conditional moment and the failure rate (Section 2).

2. Characterization of the Gamma distribution

Let X be a non-negative, absolutely continuous r.v. with distribution function (df) $F(x)$. Then $\overline{F}(x) = 1 - F(x)$, $f(x) = F'(x)$ and $r(x) = f(x)/\overline{F}(x)$ are respectively the survival function, the density function (pdf) and the failure rate of X. In the following, it is said that the r.v. X has the Gamma distribution with parameters (α, β) with $\alpha > 0$ and $\beta > 0$ (denoted by $X \sim \Gamma(\alpha, \beta)$) if $f(x) = \beta^\alpha \left[\Gamma(\alpha)\right]^{-1} \cdot \mathrm{e}^{-\beta x} x^{\alpha-1} \cdot I\{x > 0\}$.

Without loss of generality the characterization below is obtained for the Gamma distribution with parameters $(\lambda, 1)$. Let us denote, for any positive integer k and for any number $a \neq 0$, by $a^{(k)}$ the product $a^{(k)} = a(a-1)\dots(a-k+1)$ and put $a^{(0)} = 1$.

Theorem. *Let X be a non-negative r.v. with* df $F(x)$, pdf $f(x)$ *such that $f'(x)$ is continuous on $(0, \infty)$ and with failure rate $r(x)$. Then $X \sim \Gamma(\lambda, 1)$ if and only if for an integer $k \geqslant 1$, the representation*

$$E\{X^k \mid X \geqslant y\} = \left[\sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i}\right] r(y) + (\lambda + k - 1)^{(k)} \tag{1}$$

holds for all $y \geqslant 0$.

Proof. (*Necessity*): Suppose $X \sim \Gamma(\lambda, 1)$, i.e. $f(x) = x^{\lambda-1}\mathrm{e}^{-x}/\Gamma(\lambda)$. For any given integer j, it is easy to obtain that the following recurrence relation holds:

$$\int_y^\infty t^j f(t)\,\mathrm{d}t = y^j f(y) + (\lambda + j - 1) \int_y^\infty t^{j-1} f(t)\,\mathrm{d}t \quad \text{for all } y \geqslant 0. \tag{2}$$

In the case of $y = 0$, (2) is the well-known relation between the moments $\mathrm{E}X^j$ and $\mathrm{E}X^{j-1}$ of a Gamma distributed random variable X, namely $\mathrm{E}X^j = (\lambda + j - 1)\mathrm{E}X^{j-1}$.

Dividing (2) by y^j and denoting $I_j = \int_y^\infty \left(\frac{t}{y}\right)^j f(t)\,\mathrm{d}t$ we rewrite (2) as a recurrence relations between the terms of the sequence $\{I_j\}$, i.e. $I_j = f(y) + \frac{(\lambda+j-1)}{y} I_{j-1}$. Therefore, noting that $I_0 = \overline{F}(y)$, we obtain the representation

$$I_j = \sum_{i=0}^{j-1} \frac{(\lambda + j - 1)^{(i)}}{y^i} f(y) + \frac{(\lambda + j - 1)^{(j)}}{y^j} \overline{F}(y).$$

Consequently, for any given integer $j \geqslant 1$ we have

$$\int_y^\infty t^j f(t)\,\mathrm{d}t = \sum_{i=0}^{j-1} (\lambda + j - 1)^{(i)} y^{j-i} f(y) + (\lambda + j - 1)^{(j)} \overline{F}(y).$$

Now, setting $j = k$, dividing by $\overline{F}(y)$ and taking into account that $\mathrm{E}\{X^k \mid X \geqslant y\} = \frac{1}{\overline{F}(y)} \int_y^\infty t^k f(t)\, \mathrm{d}t$ we complete the proof of necessity.

(*Sufficiency*): Let us assume that (1) is true, then we have to prove that $X \sim \Gamma(\lambda, 1)$. We write (1) in the form of an equation with an unknown function

$$\int_y^\infty t^k f(t)\, \mathrm{d}t = \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i} f(y) + (\lambda + k - 1)^{(k)} \int_y^\infty f(t)\, \mathrm{d}t.$$

Differentiating both sides of the last equation with respect to y gives

$$-y^k f(y) = \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} \left[y^{k-i} f'(y) + (k-i) k^{k-i-1} f(y)\right] - (\lambda + k - 1)^{(k)} f,$$

which is a differential equation of order one with respect to the unknown function $f(y)$. The last equation can be written in the form

$$\begin{aligned} (3) \qquad & \left\{ y \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i-1} \right\} f'(y) \\ & = \left\{ (\lambda + k - 1)^{(k)} - y^k - \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} (k-i) y^{k-i-1} \right\} f(y). \end{aligned}$$

Now applying the auxiliary identity $(\lambda - 1 + k - i)(\lambda + k - 1)^{(i)} = (\lambda + k - 1)^{(i+1)}$ it can easily be verified that

$$(\lambda + k - 1)^{(k)} - y^k - \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} (k-i) y^{k-i-1} = (\lambda - 1 - y) \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i-1}.$$

To prove this relation we proceed as follows: We write

$$\begin{aligned} (*) = (\lambda - 1 - y) \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i-1} + \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} (k-i) y^{k-i-1} \\ = \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i+1)} y^{k-i-1} - \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i}. \end{aligned}$$

The change of the summation index i by $j = i + 1$ in the first term yields to

$$\begin{aligned} (*) &= \sum_{j=1}^{k} (\lambda + k - 1)^{(j)} y^{k-j} - \sum_{i=0}^{k-1} (\lambda + k - 1)^{(i)} y^{k-i} \\ &= (\lambda + k - 1)^{(k)} - y^k, \end{aligned}$$

which we wanted to verify.

Then the equation (3) is equivalent to the equation

$$\left\{\sum_{i=0}^{k-1}(\lambda+k-1)^{(i)}y^{k-i-1}\right\}yf'(y)=\left\{\sum_{i=0}^{k-1}(\lambda+k-1)^{(i)}y^{k-i-1}\right\}(\lambda-1-y)f(y).$$

Hence for $y \geqslant 0$ we derive

$$yf'(y)=(\lambda-1-y)f(y).$$

Consequently, $f(y)=Ae^{-y}y^{\lambda-1}$ for some constant $A>0$. Therefore, the solution of (3) is $f(y)=Ae^{-y}y^{\lambda-1}$. The constant A can be determined from $\int_0^\infty f(y)\,\mathrm{d}y=1$. Then we have $f(y)=y^{\lambda-1}e^{-y}/\Gamma(\lambda)$, which is the gamma pdf with parameters $(\lambda, 1)$. □

References

[1] *T. A. Azlarov, N. A. Volodin*: Characterization problems associated with the exponential distribution., "FAN", Tashkent, (1982).

[2] *J. Galamos, S. Kotz*: Characterizations of probability distributions, Lect. Notes Math. *675* (1978).

[3] *A. M. Kagan, Y. V. Linnik, C. R. Rao*: Characterization problems in mathematical statistics, "Nauka", Moskva, 1972.

[4] *J. Letac*: A characterization of the gamma distribution, Adv. Apl. Prob. *17* (1985), 911–912.

[5] *E. Lukacs*: A characterization of the gamma distribution, Ann. Math. Statist. *26* (1965), 319–324.

[6] *Shunji Osaki, Xin-xiang Li*: Characterizations of gamma and negative binomial distributions, IEEE Transactions on reliability *37 (4)* (1988), 379–382.

Author's address: *Maia Koicheva*, EHSAL, Stormstraat 2, 1 000 Brussels, Belgium.