Applications of Mathematics

Wiktor Oktaba; Andrzej Kieloch

Wishart distributions in the multivariate Gauss-Markoff model with singular covariance matrix

Applications of Mathematics, Vol. 38 (1993), No. 1, 61--66

Persistent URL: http://dml.cz/dmlcz/104534

Terms of use:

© Institute of Mathematics AS CR, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://project.dml.cz

WISHART DISTRIBUTIONS IN THE MULTIVARIATE GAUSS-MARKOFF MODEL WITH SINGULAR COVARIANCE MATRIX

WIKTOR OKTABA, ANDRZEJ KIELOCH, Lublin

(Received January 30, 1992)

Summary. This paper concerns generalized quadratic forms for the multivariate case. These forms are used to test linear hypotheses of parameters for the multivariate Gauss-Markoff model with singular covariance matrix. Distributions and independence of these forms are proved.

Keywords: multivariate general linear Gauss-Markoff model, Wishart distribution, multinormal distribution, set of linear estimable parametric functions, test, quadratic form, singular covariance matrix

AMS classification: 62H10

1. INTRODUCTION

Let $(U, XB, \Sigma \otimes \sigma^2 V)$ denote multivariate Gauss-Markoff model, with known $V \geqslant 0$, where $\otimes$ is the symbol of Kronecker product of matrices, B is a matrix of unknown fixed parameters, X is a given known design matrix, U is a random matrix of observations with the expected value $\varepsilon(U) = XB$ and with a fixed nonsingular matrix Σ, σ^2 is unknown positive scalar. Let us state that $T = V + XMX'$, where $M = M'$ is an arbitrary matrix such that $R(X) \subset R(T)$. The symbol $R(X)$ is used to denote the vector space spanned on the columns of the matrix X.

Problem of testing the hypothesis $\mathrm{L}B = 0$ in the model $(U, XB, \Sigma \otimes V)$, $V \geqslant 0$ and in the model $(U, XBA, \Sigma \otimes I)$ is presented by Srivastava and Khatri ([7], pp. 170–193), G.A.F. Seber ([6], p. 423) considers testing hypothesis $\mathrm{L}BA = 0$ in $(U, XB, \Sigma \otimes I)$ or equivalent by testing hypothesis $\mathrm{L}B^* = 0$ in $(U^*, XB^*, \Sigma \otimes I)$.

K.V. Mardia, J.T. Kent and J.M. Bibby [1] transform testing hypothesis $\mathrm{L}BA = D$ in $(U, XB, \Sigma \otimes I)$ into testing hypothesis $\mathrm{L}B = D$ in $(UM, XBM, M'\Sigma M \otimes I)$.

The case when V is allowed to be singular is discussed in full by Rao and Mitra [4], but in the univariate case only. The case of a singular V has not received much explicit attention in literature, so three theorems concerning testing a set of hypotheses of the form $\mathrm{L}BA = 0$ are presented and proved in this paper. Wilk's, Hotelling's, Pillai's or Roy's test can be used.

2. Distributions of quadratic forms

Let $\mathrm{L}BA$ be a set of linear estimable parametric functions, (Roy, [5]) where L and A are $a \times m$ and $p \times b$ matrices respectively.

The test of the estimable hypothesis

$$\mathrm{L}BA = 0 \tag{2.1}$$

can be deduced from the known Wilk's, Hotelling's or Roy's tests.

In constructing the corresponding test functions the following quadratic forms can be used:

$$S_H = (\mathrm{L}\hat{B}A)'L^-(\mathrm{L}\hat{B}A) = (\mathrm{L}\hat{B}A)'(\mathrm{L}C_4\mathrm{L}')^-(\mathrm{L}\hat{B}A), \tag{2.2}$$

$$S_E = A'U'C_1UA, \tag{2.3}$$

where

$$\hat{B} = (X'T^-X)^-X'T^-U = C_2'U = C_3U \tag{2.4}$$

$$\begin{cases} C_1 = T^- - T^-X(X'T^-X)^-X'T^- = T^-(I - XC_3) \\ C_2' = C_3 = (X'T^-X)^-X'T^- \end{cases} \tag{2.5}$$

$$L = (\mathrm{L}C_3)V(\mathrm{L}C_3)' = \mathrm{L}C_4\mathrm{L}' \tag{2.6}$$

(Oktaba, [2] (2.1), p. 179).

The symbol L^- is reserved for any choice of the g-inverse, i.e. the following relation holds:

$$LL^-L = L. \tag{2.7}$$

Let the symbol $W_b(\nu, \Sigma)$ denote the central Wishart distribution with ν degrees of freedom and with the dispersion matrix Σ. We recall that

$$\begin{pmatrix} V & X \\ X' & 0 \end{pmatrix}^- = \begin{pmatrix} C_1 & C_2 \\ C_3 & -C_4 \end{pmatrix}.$$

Theorem 2.1. *If*

$$U \sim N_{np}(XB, \Sigma \otimes \sigma^2 V) \tag{2.8}$$

where $N_{np}(\ ,\ )$ denotes an np-variate normal distribution N_{np} with parameters defined in the "Introduction", LBA is a set of estimable linear combinations of parameters and the hypothesis (2.1) *is true, then*

$$S_H \sim W_b\left[r(L), \sigma^2 A'\Sigma A\right]$$

Proof. Subject to the assumptions that the hypothesis (2.1) is true and LBA is a set of estimable parametric functions, the quadratic form (2.2) can be presented in the form:

$$\begin{aligned} S_H &= (L\hat{B}A)'L^-(L\hat{B}A) = (L\hat{B}A - LBA)'L^-(L\hat{B}A - LBA) \\ &= (LC_3UA - LC_3XBA)'L^-(LC_3UA - LC_3XBA) \\ &= (UA - XBA)'(LC_3)'L^-LC_3(UA - XBA) \\ &= (UA - XBA)'D(UA - XBA) = \left[UA - \varepsilon(UA)\right]'D\left[UA - \varepsilon(UA)\right], \end{aligned} \tag{2.9}$$

where

$$D = \left[L(X'T^-X)^-X'T^-\right]'L^-\left[L(X'T^-X)^-XT^-\right] = (LC_3)'L^-(LC_3). \tag{2.10}$$

Using (2.8) we obtain that

$$UA \sim N_{nb}(XBA, A'\Sigma A \otimes \sigma^2 V). \tag{2.11}$$

By the definition of Wishart distribution (cf. Rao, [3], p. 534) we have $S_H \sim W_b\left[r(L), \sigma^2 A'\Sigma A\right]$ if and only if

$$VDVDV = VDV \quad \text{and} \quad r(L) = \mathrm{tr}(VD), \tag{2.12}$$

where $tr(A)$ denotes the trace of the matrix A.

We will prove now that the relation (2.12) holds. In fact,

$$
\begin{aligned}
(2.13) \qquad VDVDV &= V(\mathrm{L}C_3)'L^-\mathrm{L}C_3V(\mathrm{L}C_3)'L^-\mathrm{L}C_3V \\
&= V(\mathrm{L}C_3)'L^-LL^-\mathrm{L}C_3V = V(\mathrm{L}C_3)'L^-\mathrm{L}C_3V = VDV,
\end{aligned}
$$

and moreover,

$$
\begin{aligned}
\mathrm{tr}(VD) &= \mathrm{tr}\left[V(\mathrm{L}C_3)'L^-\mathrm{L}C_3\right] = \mathrm{tr}\left[L^-\mathrm{L}C_3V(\mathrm{L}C_3)'\right] \\
&= \mathrm{tr}(L^-L) = r(L).
\end{aligned}
$$

Hence the result (2.12) follows. □

Let $(V\vdots X)$ denote a partitioned matrix.

Theorem 2.2. *Subject to the assumption* (2.8) *we have*

$$
S_E \sim W_b\left[r(V\vdots X) - r(X), \sigma^2 A'\Sigma A\right]
$$

where S_E *is as defined in* (2.3).

Proof. By the relation

$$
(2.14) \qquad X'C_1 = X'\left[T^- - T^-X(X'T^-X)^-X'T^-\right] = 0
$$

the matrix (2.3) can be presented in the form

$$
\begin{aligned}
(2.15) \qquad S_E &= (UA)'C_1(UA) = (UA)'(T^- - T^-X(X'T^-X)^-X'T^-)UA \\
&= (UA - XBA)'\left[T^- - T^-X(X'T^-X)^-X'T^-\right](UA - XBA) \\
&= \left[UA - \varepsilon(UA)\right]'C_1\left[UA - \varepsilon(UA)\right].
\end{aligned}
$$

In virtue of (2.11) the matrix (2.15) has the Wishart distribution with the parameters as in Theorem 2.2 if and only if (cf. proof of Theorem 2.1)

$$
(2.16) \qquad VC_1VC_1V = VC_1V
$$

and

$$
(2.17) \qquad \mathrm{tr}\, VC_1 = r(V\vdots X) - r(X).
$$

Let us see that conditions (2.16) and (2.17) are fulfilled. In fact, by (2.14) and the definition of the matrix T we have $VC_1 = TC_1$ and the right hand side of (2.16) can be written as

(2.18)
$$\begin{aligned} VC_1VC_1V &= TC_1TC_1V \\ &= T[T^- - T^-X(X'T^-X)^-X'T^-]T[T^- - T^-X(X'T^-X)^-X'T^-]V \\ &= T[T^- - T^-X(X'T^-X)^-X'T^-]V = TC_1V = VC_1V. \end{aligned}$$

The relations (2.18) show that S_E has the central Wishart distribution.

Now we prove (2.17). By the definition of T and (2.14) we obtain

$$\begin{aligned} \operatorname{tr} VC_1 &= \operatorname{tr} TC_1 = \operatorname{tr}\left[TT^- - TT^-X(X'T^-X)^-X'T^-\right] \\ &= \operatorname{tr} TT^- - \operatorname{tr} X(X'T^-X)^-X'T^- = r(T) - \operatorname{tr}(X'T^-X)^-X'T^-X \\ &= r(V\vdots X) - r(X'T^-X) = r(V\vdots X) - r(X). \end{aligned} \tag{2.19}$$

□

Theorem 2.3. *If the assumptions of Theorems 2.1 and 2.2 concerning matrices S_H and S_E are fulfilled, then S_H and S_E are stochastically independent.*

Proof: A necessary and sufficient condition for S_H and S_E to be independently distributed is

$$VDVC_1V = 0, \tag{2.20}$$

where C_1 and D are defined as in (2.5) and (2.10) respectively. In virtue of (2.14), $X'T^-T = X'$, we have

$$\begin{aligned} VDVC_1V &= VDTC_1V = V[\mathrm{L}(X'T^-X)^-X'T^-]'L^-[\mathrm{L}(X'T^-X)^-X'T^-]TC_1V \\ &= V[\mathrm{L}(X'T^-X)^-XT^-]'L^-\mathrm{L}(X'T^-X)^-X'C_1V = 0. \end{aligned}$$

It means that the condition (2.20) holds for S_H and S_E, so the forms considered are independent. □

References

[1] *Mardia, K.V., Kent, J.T. and Bibby, J.M.*: Multivariate Analysis, Academic Press, London, 1979.

[2] *Oktaba, W.*: Estimation of the parametric functions in the multivariate general Gauss–Markoff model, XV Colloquium Metodol., Agrobiom., Pol. Acad. Sci., Warsaw, 1985, pp. 178–182. (In Polish.)

[3] *Rao, C.R.*: Linear Statistical Inference and Its Applications, sec. ed., J. Wiley, New York, 1973.

[4] *Rao, C.R. and Mitra, S.K.*: Generalized Inverse of Matrices and its Applications, J. Wiley, New York, 1971.

[5] *Roy, S.N.*: Some aspects of multivariate analysis, Indian Statistical Institute, New York, Calcutta, 1957.

[6] *Seber, G.A.F.*: Multivariate Observations, J. Wiley, New York, 1984.

[7] *Srivastava, M.S. and Khatri, C.G.*: An Introduction to Multivariate Statistics, Elsevier North Holland, Inc., New York, 1979.

Authors' address: *Wiktor Oktaba*, *Andrzej Kieloch*, Institute of Applied Mathematics, Academy of Agriculture, Akademicka 13, P.O. Box 325, 20-950 Lublin, Poland.