

K. Denecke; Jörg Koppitz
Pre-solid varieties of semigroups

Archivum Mathematicum, Vol. 31 (1995), No. 3, 171--181

Persistent URL: <http://dml.cz/dmlcz/107538>

Terms of use:

© Masaryk University, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

PRE-SOLID VARIETIES OF SEMIGROUPS

K. DENECKE AND J. KOPPITZ

ABSTRACT. Pre-hyperidentities generalize the concept of a hyperidentity. A variety V is said to be pre-solid if every identity in V is a pre-hyperidentity. Every solid variety is pre-solid. We consider pre-solid varieties of semigroups which are not solid, determine the smallest and the largest of them, and some elements in this interval.

1. INTRODUCTION

An identity $t \approx t'$ is called a *hyperidentity* in a variety V if whenever the operation symbols occurring in t and t' are replaced by any terms of the appropriate arity, the identity which results holds in V ([14]). Hyperidentities can be defined more precisely using the concept of a *hypersubstitution* ([2]).

We fix a type $\tau = (n_i)_{i \in I}$, $n_i > 0$ for all $i \in I$, and operation symbols $(f_i)_{i \in I}$, where f_i is n_i -ary. Let $W_\tau(X)$ be the set of all terms of type τ over some fixed alphabet X , and let $Alg(\tau)$ be the class of all algebras of type τ . Then a mapping

$$\sigma : \{f_i | i \in I\} \rightarrow W_\tau(X)$$

which assigns to every n_i -ary operation symbol f_i an n_i -ary term will be called a hypersubstitution of type τ (for short, a hypersubstitution). For a term $t \in W_\tau(X)$ by $\hat{\sigma}[t]$ we define the application of the hypersubstitution σ to the term t . The term $\hat{\sigma}[t]$ can be defined inductively by:

- (i) $\hat{\sigma}[x] := x$ for any variable x in the alphabet X , and
- (ii) $\hat{\sigma}[f_i(t_1, \dots, t_{n_i})] := \sigma(f_i)^{W_\tau(X)}(\hat{\sigma}[t_1], \dots, \hat{\sigma}[t_{n_i}])$.

It is clear that $\sigma(f_i)^{W_\tau(X)}$ on the right hand side of (ii) is the operation induced by $\sigma(f_i)$ on the term algebra $W_\tau(X)$.

An identity $t \approx t'$ where t, t' are terms of type τ is a hyperidentity of type τ (for short a hyperidentity) in an algebra $\mathcal{A} \in Alg(\tau)$ if $\hat{\sigma}[t] \approx \hat{\sigma}[t']$ is an identity in \mathcal{A} for every hypersubstitution σ .

An important example of a hyperidentity is the type $\tau = (2)$ medial hyperidentity

$$F(F(x, y), F(z, t)) \approx F(F(x, z), F(y, t))$$

1991 *Mathematics Subject Classification*. 20M07, 08B15.

Key words and phrases. hyperidentity, pre-hyperidentity, pre-solid variety.

Received April 6, 1994

which is a hyperidentity in the variety of all medial semigroups (defined by the identity $xyzt \approx xzyt$ ([15])).

On the basis of a hypersubstitution in [2] a closure operator Ξ defined on classes of algebras of type τ and of sets of equations was introduced. If $t \approx t'$ is an equation then by $\Xi[t \approx t']$ we denote the set $\{\bar{\sigma}[t] \approx \bar{\sigma}[t'] \mid \sigma \text{ is a hypersubstitution of type } \tau\}$ and by $\Xi[\Sigma]$ for a set Σ of equations the union of the sets $\Xi[t \approx t']$ for $t \approx t'$ in Σ . Let $\mathcal{A} = (A; (f_i^A)_{i \in I})$ be an algebra in $\text{Alg}(\tau)$ and let σ be a hypersubstitution. Then we make the following definitions:

$$\begin{aligned} \sigma[\mathcal{A}] &:= (A; (\sigma(f_i)^A)_{i \in I}), \\ \Xi[\mathcal{A}] &:= \{\sigma[\mathcal{A}] \mid \sigma \text{ is a hypersubstitution of type } \tau\}, \\ \Xi[K] &:= \bigcup_{\mathcal{A} \in K} \Xi[\mathcal{A}]. \end{aligned}$$

A variety V of type τ is *solid* if $\Xi[V] = V$. In [2] it was shown that a variety V is solid if and only if every identity in V is a hyperidentity. Equivalently, solid varieties can be characterized as classes of algebras satisfying a given set of equations as hyperidentities (*hyperequational classes*). Although the concept of a solid variety is very strong there are infinitely many solid varieties of semigroups ([3]). All solid varieties of a given type τ form a lattice which is a sublattice of the lattice of all varieties of type τ .

Note that the clone of a solid variety V (denoted by *clone* V see [14]) is a free heterogeneous algebra with respect to the variety of heterogeneous algebras generated by *clone* V ([6]). There is a lot of equations which cannot be hyperidentities in a nontrivial algebra (or variety). For instance, substituting one of the binary projections for F in the commutative law $F(x, y) \approx F(y, x)$ we get $x \approx y$ which is satisfied only in a trivial algebra. This observation motivates to weaken the concept of a hyperidentity. The simplest way for weakness could be to substitute only terms different from variables. The set of all term functions of an algebra which are different from projections forms a so-called pre-iterative algebra ([11]). Therefore these weaker hyperidentities are called pre-hyperidentities in [7]. At first we will set up the preliminary results and notation we will need and then we will consider the set of all pre-solid varieties of semigroups.

2. BASIC CONCEPTS

According to the ideas explained in the introduction we define a *pre-hypersubstitution* of type τ as a mapping

$$\sigma_p : \{f_i \mid i \in I\} \rightarrow W_\tau(X) \setminus X$$

which assigns to every operation symbol f_i an n_i -ary term which is different from a variable. (Note that we consider the first n_i variables x_0, \dots, x_{n_i-1} of the standard alphabet $X = \{x_0, \dots, x_{n_i-1}, \dots\}$ as n_i -ary terms. A composed term is called n_i -ary if it is built up from operation symbols of the corresponding arities and variables from this alphabet X .)

The extension $\hat{\sigma}_p[t]$ of a pre-hypersubstitution to a term t is defined inductively by

- (i) $\hat{\sigma}_p[x] := x$ for any variable x in the alphabet X , and
- (ii) $\hat{\sigma}_p[f_i(t_1, \dots, t_{n_i})] := \sigma_p(f_i)^{\mathcal{W}_\tau(X)}(\hat{\sigma}_p[t_1], \dots, \hat{\sigma}_p[t_{n_i}])$.

The expression $\sigma_p(f_i)^{\mathcal{W}_\tau(X)}$ on the right hand side of (ii) is the operation induced by $\sigma(f_i)$ on the term algebra $\mathcal{W}_\tau(X)$.

If $t \approx t'$ is an equation, then we denote by $\Xi_p[t \approx t']$ the set

$$\{\hat{\sigma}_p[t] \approx \hat{\sigma}_p[t'] \mid \sigma_p \text{ is a pre-hypersubstitution}\}$$

If Σ is a set of equations, we use $\Xi_p[\Sigma]$ for the union of the sets $\Xi_p[t \approx t']$, for $t \approx t'$ in Σ .

Let $\mathcal{A} = (A; (f_i^A)_{i \in I})$ be an algebra in $Alg(\tau)$, and let K be a class of algebras of type τ . Then we define:

$$\begin{aligned} \sigma_p[\mathcal{A}] &:= (A; (\sigma_p(f_i)^A)_{i \in I}), \\ \Xi_p[\mathcal{A}] &:= \{\sigma_p[\mathcal{A}] \mid \sigma_p \text{ is a pre-hypersubstitution of type } \tau\}, \\ \Xi_p[K] &:= \bigcup_{\mathcal{A} \in K} \Xi_p[\mathcal{A}]. \end{aligned}$$

In [7] we proved the following proposition:

Result 2.1. Ξ_p is a closure operator on sets of equations Σ and on classes of algebras K of type τ , i.e.

- (i) $\Sigma \subseteq \Xi_p[\Sigma]$,
- (ii) $\Sigma' \subseteq \Sigma \Rightarrow \Xi_p[\Sigma'] \subseteq \Xi_p[\Sigma]$,
- (iii) $\Xi_p[\Xi_p[\Sigma]] = \Xi_p[\Sigma]$,
- (i') $K \subseteq \Xi_p[K]$,
- (ii') $K' \subseteq K \Rightarrow \Xi_p[K'] \subseteq \Xi_p[K]$,
- (iii') $\Xi_p[\Xi_p[K]] = \Xi_p[K]$. □

Since every pre-hypersubstitution is a hypersubstitution we have

Result 2.2. Let K be a class of algebras of type τ and let Σ be a set of equations of type τ . Then

- (i) $\Xi_p[\Sigma] \subseteq \Xi[\Sigma]$ and
- (ii) $\Xi_p[K] \subseteq \Xi[K]$. □

Using the concept of a pre-hypersubstitution we define pre-hyperidentities in the following way:

Definition 2.3. Let $\mathcal{A} \in Alg(\tau)$ be an algebra of type τ . Then the identity $t \approx t'$, where t, t' are terms of type τ is a pre-hyperidentity of type τ in \mathcal{A} (\mathcal{A} pre-hypersatisfies $t \approx t'$) if $\hat{\sigma}_p[t] \approx \hat{\sigma}_p[t']$ is an identity of \mathcal{A} for every pre-hypersubstitution σ_p .

Clearly, every hyperidentity of type τ is a pre-hyperidentity of this type. In general, the converse is false.

For a class K of algebras of type τ and for a set Σ of equations of this type we fix the following notations:

IdK - the class of all identities of K ,

$HIdK$ - the class of all hyperidentities of K ,

H_pIdK - the class of all pre-hyperidentities of K ,

$Mod\Sigma = \{\mathcal{A} \in Alg(\tau) \mid \mathcal{A} \text{ satisfies } \Sigma\}$ - the variety defined by Σ

$HMod\Sigma = \{\mathcal{A} \in Alg(\tau) \mid \mathcal{A} \text{ hypersatisfies } \Sigma\}$ - the hyperequational class defined by Σ ,

$H_pMod\Sigma = \{\mathcal{A} \in Alg(\tau) \mid \mathcal{A} \text{ pre-hypersatisfies } \Sigma\}$ - the pre-hyperequational class defined by Σ ,

$VarK = ModIdK$ - the variety generated by K ,

$HVarK = HModHIdK = \{\mathcal{A} \in Alg(\tau) \mid \mathcal{A} \text{ hypersatisfies } HIdK\}$ - the hypervariety of type τ generated by K .

For these sets we get the following inclusions:

$$HIdK \subseteq H_pIdK, HMod\Sigma \subseteq H_pMod\Sigma.$$

By definition every hyperidentity or every pre-hyperidentity is an identity. Very natural there arises the problem to find algebras or varieties for which every identity is a hyperidentity or such that every identity is a pre-hyperidentity.

Definition 2.4. Let V be a variety of type τ . Then V is called pre-solid if $\Xi_p[V] = V$.

In [7] pre-solid varieties were characterized in the following manner:

Result 2.5. ([7]) *Let $K \subseteq Alg(\tau)$ be a variety. Then the following conditions are equivalent:*

- (i) K is a pre-hyperequational class,
- (ii) K is pre-solid,
- (iii) $IdK \subseteq H_pIdK$, i.e. every identity of K is a pre-hyperidentity,
- (iv) $\Xi_p[IdK] = IdK$, i.e. IdK is closed under pre-hypersubstitutions. □

For a given type τ by $\mathcal{L}(\tau)$ we denote the lattice of all varieties of this type and by $S(\tau)$ the set of all solid varieties of this type. $S_p(\tau)$ is the set of all pre-solid varieties of type τ . Then we have the following results:

Result 2.6. ([10], [13])

- (i) The set $S(\tau)$ forms a sublattice of $\mathcal{L}(\tau)$,
- (ii) The set $S_p(\tau)$ forms a sublattice of $\mathcal{L}(\tau)$ containing $S(\tau)$ as a sublattice.
- (iii) If τ is a finite type then the lattice $\mathcal{S}(\tau)$ is atomic. The unique atom is the variety RA_τ of all rectangular algebras of type τ . (RA_τ is the variety generated by all algebras of type τ whose fundamental operations are projections). □

3. SOLID AND PRE-SOLID VARIETIES OF SEMIGROUPS

By $\mathcal{L}(S)$ we denote the lattice of all semigroup varieties. To describe a bit more of the structure of all pre-solid varieties of semigroups we note that a variety of semigroups to be solid it must satisfy the associative law as a hyperidentity. That means, the greatest solid variety of semigroups is the hypermodel class of the associative law: $HMod\{F(F(x, y), z) \approx F(x, F(y, z))\}$. In [1] we obtained an equational basis for this variety:

$HMod\{F(F(x, y), z) \approx F(x, F(y, z))\} = V_{HS}$ with $V_{HS} = Mod\{I_1 \cup I_2 \cup \{x^2 \approx x^4\}\}$ where I_1 and I_2 are the following sets of identities:

$$I_1 := \{(x^{k_1}y^{k_2} \dots x^{k_{n-1}}y^{k_n})^{k_1}z^{k_2} \dots (x^{k_1}y^{k_2} \dots x^{k_{n-1}}y^{k_n})^{k_{n-1}}z^{k_n} \approx x^{k_1}(y^{k_1}z^{k_2} \dots y^{k_{n-1}}z^{k_n})^{k_2} \dots x^{k_{n-1}}(y^{k_1}z^{k_2} \dots y^{k_{n-1}}z^{k_n})^{k_n} \mid n \in \{2, 4, 6\} \text{ for } 1 \leq k_1, \dots, k_n \leq 3\}.$$

$$I_2 := \{(x^{k_1}(y^{k_1}z^{k_2}y^{k_3} \dots z^{k_{n-1}}y^{k_n})^{k_2} \dots (y^{k_1}z^{k_2}y^{k_3} \dots z^{k_{n-1}}y^{k_n})^{k_{n-1}}x^{k_n} \approx (x^{k_1}y^{k_2}x^{k_3} \dots y^{k_{n-1}}x^{k_n})^{k_1}z^{k_2} \dots (x^{k_1}y^{k_2}x^{k_3} \dots y^{k_{n-1}}x^{k_n})^{k_n} \mid n \in \{3, 5\} \text{ for } 1 \leq k_1, \dots, k_n \leq 3\}.$$

Note that L. Polák recently proved that $V_{HS} = Mod\{x(yz) \approx (xy)z, xyxzyx \approx xzyyx, x^2 \approx x^4, xy^2z^2 \approx xyzy^2z^2, x^2y^2z \approx x^2yx^2yz\}$.

By $S(V_{HS})$ we denote the lattice of all solid semigroup varieties (indeed, the set of all solid semigroup varieties forms a sublattice of $\mathcal{L}(S)$ since $S(V_{HS})$ is the intersection of the subvariety lattice of the variety V_{HS} and the lattice of all solid varieties of type $\tau = (2)$.)

According to Result 2.6 every non trivial solid semigroup variety contains the variety RB of all rectangular bands which is defined by the identities $x(yz) \approx (xy)z, x^2 \approx x, xyz \approx xz$. Then we have:

Result 3.1. ([5]) *The variety RB is the least nontrivial element of $S(V_{HS})$.* □

We call an equation $t \approx t'$ to be left-most (right-most) if the left-most (right-most) variables in t and t' are the same. An equation $t \approx t'$ will be called outer-most if it is left-most and right-most. These notions were used by E. Graczyńska in [9]. Let $Out(2)$ be the set of all outer-most equations of type $\tau = (2)$. Clearly, $IdRB = Out(2)$. As a consequence, a hyperidentity must be an outer-most equation. The following useful fact is obvious:

Proposition 3.2. *Let V be a nontrivial variety of semigroups and let $t \approx t'$ be an outer-most equation of type $\tau = (2)$. Then $t \approx t'$ is a hyperidentity in V if and only if $t \approx t'$ is a pre-hyperidentity in V .*

Proof. Clearly, if $t \approx t'$ is a hyperidentity satisfied in V then it is a pre-hyperidentity in V . Let $t \approx t'$ be a pre-hyperidentity satisfied in V . Then for every pre-hypersubstitution σ_p we have $\sigma_p[t] \approx \sigma_p[t'] \in IdV$. A hypersubstitution which is no pre-

hypersubstitution assigns to the binary fundamental operation symbol a binary projection. Since $t \approx t'$ is outer-most the equation $\sigma[t] \approx \sigma[t']$ is equal to $x \approx x$ or to $y \approx y$ which is an identity in V . Altogether, for every hypersubstitution σ we have $\sigma[t] \approx \sigma[t'] \in IdV$ and $t \approx t'$ is a hyperidentity in V . \square

Clearly, the associative law is an outermost equation and applying Proposition 3.2 we have:

Corollary 3.3. *The variety V_{HS} is pre-solid and for any pre-solid variety V of semigroups, $V \subseteq V_{HS}$.*

Proof. As a solid variety V_{HS} is pre-solid. Since V_{HS} is the hyperequational class generated by the associative law it is also the pre-hyperequational class generated by the associative law and thus the greatest pre-solid variety of semigroups. \square

Let $S_p(V_{HS})$ be the set of all pre-solid semigroup varieties. We want to discuss the following question:

Are there pre-solid semigroup varieties in the interval between RB and V_{HS} which are not solid?

The answer is given in [7], namely

Lemma 3.4. ([7]) *Let V be a variety of type $\tau = (2)$ such that $RB \subseteq V$. Then V is solid iff V is pre-solid.*

Proof. If V is solid then V is also pre-solid. Let V be pre-solid. The inclusion $RB \subseteq V$ means that every identity in V is outer-most. Therefore, by Proposition 3.2 every pre-hyperidentity is a hyperidentity and since every identity is a pre-hyperidentity V must be solid. \square

Lemma 3.4 shows that a pre-solid variety of semigroups which is not solid must be outside of the interval between RB and V_{HS} . Now we ask for the greatest pre-solid semigroup variety which is not solid.

Proposition 3.5. *The variety $V_{PS} := Mod\{(xy)z \approx x(yz), xyxzxxyx \approx xyzyx, x^2 \approx y^2, x^3 \approx y^3\}$ is pre-solid, but not solid.*

Proof. Since the identity $x^2 \approx y^2$ is no outer-most equation and since V_{PS} is nontrivial the variety V_{PS} is not solid. We are going to show that V_{PS} is the pre-hyperequational class defined by the associative law, by $F(x, x) \approx F(y, y)$, and by $F(x, F(x, x)) \approx F(y, F(y, y))$. Now, $V_{PS} \subseteq Mod\{I_1 \cup I_2 \cup \{x^2 \approx x^4\}\}$ since $I_1 \cup I_2 \cup \{x^2 \approx x^4\} \subseteq \{(xy)z \approx x(yz), xyxzxxyx \approx xyzyx\} \cup \{uv^2w \approx u'v'^2w'|u, v, w, u', v', w' \text{ are binary terms}\} \subseteq IdMod\{(xy)z \approx x(yz), xyxzxxyx \approx xyzyx, x^2 \approx y^2, x^3 \approx y^3\}$ because of $uv^2w \approx u^3w \approx w^4 \approx u'^3w' \approx u'v'^2w'$. Since $Mod\{I_1 \cup I_2 \cup \{x^2 \approx x^4\}\} = H_p Mod\{F(x, F(y, z)) \approx F(F(x, y), z)\}$ the associative law is a pre-hyperidentity in V_{PS} . We have to check that $F(x, x) \approx F(y, y)$ and $F(F(x, x), x) \approx F(F(y, y), y)$ are pre-hyperidentities in V_{PS} . For every binary term different from a variable there are natural numbers $r, s > 1$ with $t(x, x) = x^r$, $t(y, y) = y^r$, respectively with $t(t(x, x), x) = x^s$ and $t(t(y, y), y) = y^s$. Because of

$x^2 \approx y^2, x^3 \approx y^3 \in IdV_{PS}$ we get $t(x, x) \approx t(y, y), t(t(x, x), x) \approx t(t(y, y), y) \in IdV_{PS}$. (Note that $t(x, y)$ means, the term is constructed only from the variables x and y .) Altogether, $V_{PS} \subseteq H_p Mod\{F(F(x, y), z) \approx F(x, F(y, z)), F(x, x) \approx F(y, y), F(F(x, x), x) \approx F(F(y, y), y)\}$. On the other hand we see that the identities $x(yz) \approx (xy)z, xyxzxyx \approx xyzyx, x^2 \approx y^2, x^3 \approx y^3$ are satisfied in $H_p Mod\{F(F(x, y), z) \approx F(x, F(y, z)), F(x, x) \approx F(y, y), F(F(x, x), x) \approx F(F(y, y), y)\}$. This shows the equality $V_{PS} = H_p Mod\{F(F(x, y), z) \approx F(x, F(y, z)), F(x, x) \approx F(y, y), F(F(x, x), x) \approx F(F(y, y), y)\}$ and by Result 2.5 the variety V_{PS} is pre-solid. \square

Theorem 3.6. *For every nontrivial pre-solid variety V of semigroups the following propositions are equivalent:*

- (i) $V \subseteq V_{PS}$,
- (ii) V is not solid.

Proof. (i) \Rightarrow (ii): Since the equation $F(x, x) \approx F(y, y)$ is no hyperidentity in V the variety V is not solid.

(ii) \Rightarrow (i): Since V is pre-solid and not solid by Lemma 3.4 the variety of rectangular bands is not included in V . Then there is an identity in IdV which is not outer-most. We conclude that there are natural numbers m, n and variables $u_0, \dots, u_m, v_0, \dots, v_n \in X$ such that $u_0 \neq v_0$ or $u_m \neq v_n$ and $u_0 \dots u_m \approx v_0 \dots v_n \in IdV$. Without restriction of the generality assume that $u_0 \neq v_0$. Substituting $t(x, y) = x^2$ and $t(x, y) = x^3$ we obtain $u_0^2 \approx v_0^2 \in IdV$, respectively $u_0^3 \approx v_0^3 \in IdV$ and thus $V \subseteq Mod\{x^2 \approx y^2, x^3 \approx y^3\}$. Since V is pre-solid by Corollary 3.3, $V \subseteq V_{HS}$. Altogether we have $V \subseteq V_{PS}$. \square

Theorem 3.6 shows that V_{PS} is the greatest pre-solid variety of semigroups which is not solid. It is very natural to ask for the least pre-solid variety of semigroups which is not solid. Let $Z = Mod\{x(yz) \approx (xy)z, xy \approx zt\}$ be the variety of all zero-semigroups. It is easy to see that Z is pre-solid but not solid.

Theorem 3.7. *Let V be a pre-solid variety of semigroups. If V is not solid then $Z \subseteq V$.*

Proof. Since V is not solid V cannot be trivial. Any nontrivial variety V of semigroups must include at least one of the atoms of the lattice of all semigroup varieties (listed for example in [8]). Since V is pre-solid by Corollary 3.3 we have $V \subseteq V_{HS}$ and every atom included in V must be hyperassociative. In [1] we determined all hyperassociative atoms in the lattice of all semigroup varieties: zero-semigroups ($xy \approx zt$), right semigroups ($xy \approx x$), left semigroups ($xy \approx y$), semilattices ($xy \approx yx, x^2 \approx x$), and 2-groups ($xy \approx yx, x^2y \approx y$). Since by Theorem 3.6 $V \subseteq V_{PS}$ the identities $x^2 \approx y^2$ and $x^3 \approx y^3$ must be satisfied in an atom included in V . These identities are satisfied only in the variety Z of all zero-semigroups, thus $Z \subseteq V$. \square

Considering the set $S_p(V_{HS})$ we get:

Proposition 3.8. *The set $S_p(V_{HS}) \setminus S(V_{HS})$ forms a sublattice of $\mathcal{S}_p(V_{HS})$.*

Proof. Consider two varieties $V_1, V_2 \in S_p(V_{HS}) \setminus S(V_{HS})$. Then $V_1 \vee V_2$ and $V_1 \wedge V_2$ are pre-solid by Result 2.6. Because of $V_1 \vee V_2, V_1 \wedge V_2 \subseteq V_{PS}$ the equation $x^2 \approx y^2$ is satisfied in $V_1 \vee V_2$ and in $V_1 \wedge V_2$. The equation $x^2 \approx y^2$ cannot be a hyperidentity in these varieties. This shows that both varieties are not solid. \square

A solid variety V must contain the variety RB . Therefore the join of a solid variety V and the pre-solid variety Z can be written as $V \vee Z = V \vee (RB \vee Z)$. It is well-known that $RB \vee Z = Mod\{xyz \approx xz\}$ is solid ([9], [15], [3]) and that the join of two solid varieties is solid (Result 2.6). This example motivates the following question: Is the join of a solid and an arbitrary pre-solid variety solid? To attack this question we use Theorem 2.7 and obtain:

Proposition 3.9. *Let V_1 be a nontrivial solid and let V_2 be a pre-solid variety of semigroups. Then the variety $V_1 \vee V_2$ is solid.*

Proof. By Result 2.6 the variety $V_1 \vee V_2$ is pre-solid and because of $RB \subseteq V_1 \subseteq V_1 \vee V_2$ and Lemma 3.4 solid. \square

Remark that the varieties RB and V_{PS} have certain “splitting” properties for the lattice $S_p(V_{HS})$. Indeed, if the variety V is nontrivial, pre-solid and contains RB then V is solid. If the variety RB is not contained in V then V belongs to $S_p(V_{HS}) \setminus S(V_{HS})$. If V is nontrivial, pre-solid and contained in V_{PS} then V belongs to $S_p(V_{HS}) \setminus S(V_{HS})$. If V is not contained in V_{PS} then V is solid.

4. APPLICATIONS

Proposition 3.8 is useful to derive new solid varieties from given ones. We will consider a special case of this theorem.

An identity $t \approx t'$ is called normal (see e.g. [9]) if t and t' are the same variable or neither t nor t' are variables. By $N(\tau)$ we denote the set of all normal identities of type τ . For a variety V of type τ we set $N(V) := N(\tau) \cap IdV$. The variety $V_N := ModN(V)$ is called the normalization of V . It is easy to see that IdZ is the set of all normal identities of type $\tau = (2)$. Therefore, for a variety V of semigroups $V \vee Z$ is the normalization of V and we have:

Corollary 4.1. *The normalization of a pre-solid variety of semigroups is pre-solid.*

Proof. Let V be a pre-solid but not solid variety of semigroups, then by Theorem 3.7, $V_N = V = V \vee Z$. If V is solid then by Theorem 3.8 V_N is solid and thus pre-solid. \square

(For solid varieties Corollary 4.1 was shown in [4]).

One could ask whether the solidity of $V \vee Z$ implies the solidity of V .

To formulate the following Theorem we need all solid varieties of bands. There are exactly the following solid varieties of bands ([15]):

1. $RB = Mod\{x(yz) \approx (xy)z \approx xz, x^2 \approx x\}$ – rectangular bands,
2. $RegB = Mod\{x(yz) \approx (xy)z, x^2 \approx x, xyxzx \approx xyzx\}$ – regular bands,
3. $NB = Mod\{x(yz) \approx (xy)z, x^2 \approx x, xyzt \approx xzyt\}$ – normal bands.

Theorem 4.2. *Let V be a nontrivial variety of semigroups. Then the following is equivalent:*

- (i) V is solid,
- (ii) $V \vee Z$ is solid and for V there holds $V \supseteq Z$ or $V \in \{RB, NB, RegB\}$

Proof. Let V be solid, then $V \vee Z$ is solid by Corollary 4.1. Assume that $x^2 \approx x \notin IdV$. We will show that IdV consists only of normal identities. At first we check identities of the form $x^m \approx x$ with $m > 2$. Let $k > 2$ be the least natural number with $x^k \approx x \in IdV$. From $F(x^{k-1}, x) \approx x$ we obtain by the hypersubstitution $\sigma : F \mapsto x$ the identity $x^{k-1} \approx x \in IdV$ in contradiction of the minimality of k . Assume there is an identity $u \approx v \in IdV$ which is not normal and so that the sets of variables in u and in v are different. Each of the terms u and v contains at most two variables since otherwise $x^m \approx x \in IdV$ with $m \geq 3$. Therefore $u \approx v$ is one of the following identities: $yz \approx x$, or $xz \approx x$, or $zx \approx x$. These identities cannot be hyperidentities in V in contradiction to the solidity of V . This shows $Z \subseteq V$. Assume now that $x^2 \approx x \in IdV$, that is, V is a variety of bands. Then $V \in \{RB, RegB, NB\}$.

If $Z \subseteq V$ then from the solidity of $V \vee Z$ follows the solidity of $V = V \vee Z$. The solidity of $RB, RegB$, and NB is clear. □

5. EXAMPLES FOR PRE-SOLID VARIETIES OF SEMIGROUPS

In the introduction we remarked that there is no nontrivial solid variety of commutative semigroups. In [5] we determined all pre-solid varieties of commutative semigroups. Clearly, the pre-hyperequational class defined by the associative and the commutative law must be the greatest pre-solid variety of commutative semigroups (Result 2.5). In [5] we determined an equational basis for this variety. There holds:

$$H_p Mod\{F(F(x, y), z) \approx F(x, F(y, z)), F(x, y) \approx F(y, x)\} = Mod\{(xy)z \approx x(yz), xy \approx yx, xy^2 \approx x^2y, x^2 \approx y^2\} =: V_{PC}.$$

We fix the following denotations:

$$p_n : x_0x_1 \dots x_n \approx y_0y_1 \dots y_n,$$

$$I_n = \{(xy)z \approx x(yz), xy \approx yx, xy^2 \approx x^2y, x^2 \approx y^2, p_n\}$$

$$P_n = ModI_n \text{ for every natural number } n.$$

Then we have:

Result 5.1. ([5]) *The varieties $P_n, n \in N$ and V_{PC} are all pre-solid varieties of commutative semigroups. They form a lattice, namely the subvariety lattice of V_{PC} (which is a chain).* □

It is a well-known fact ([15]) that in every variety of medial semigroups the medial law is a hyperidentity. The greatest solid variety of medial semigroups is the hyperequational class defined by the medial and the associative law, $HMod\{F(F(x, y), F(z, t)) \approx F(F(x, z), F(y, t)), F(F(x, y), z) \approx F(x, F(y, z))\} = Mod\{x(yz) \approx (xy)z, xyz t \approx xzyt, x^2 \approx x^4, x^3y^2zx \approx xy^2zx\} = HM$. This variety is pre-solid. We ask for pre-solid varieties of medial semigroups which are not

solid.

Lemma 5.2. *The variety $V_{MPS} := Mod\{(xy)z \approx x(yz), xyzt \approx xzyt, x^2 \approx y^2, x^3 \approx y^3\}$ is pre-solid.*

Proof. We show that V_{MPS} is a pre-hyperequational class. The inclusion $V_{MPS} \subseteq HM$ implies that V_{MPS} satisfies the associative and the medial law as hyperidentities. These equations are outer-most and therefore pre-hyperidentities in V_{MPS} . Further, $V_{MPS} \subseteq V_{PS}$ since V_{PS} is the greatest pre-solid variety of semigroups which is not solid. Since the equations $F(x, x) \approx F(y, y)$ and $F(F(x, x), x) \approx F(F(y, y), y)$ are pre-hyperidentities in V_{PS} they are pre-hyperidentities in $V_{MPS} \subseteq V_{PS}$. This shows: $V_{MPS} \subseteq H_p Mod\{F(F(x, y), z) \approx F(x, F(y, z)), F(F(x, y), F(z, t)) \approx F(F(x, z), F(y, t)), F(x, x) \approx F(y, y), F(F(x, x), x) \approx F(F(y, y), y)\}$. On the other hand, it is easy to show that this pre-hyperequational class is included in V_{MPS} . \square

Theorem 5.3. *For every pre-solid variety V of semigroups the following is equivalent:*

- (i) V is medial but not solid,
- (ii) $V \subseteq V_{MPS}$.

Proof. (i) \Rightarrow (ii): Since V is pre-solid the associative law is a pre-hyperidentity in V and hence it is a hyperidentity in V . The variety V is medial and therefore, $V \subseteq HM = Mod\{x(yz) \approx (xy)z, xyzt \approx xzyt, x^2 \approx x^4, x^3y^2zx \approx xy^2zx\}$. Further, we have $V \subseteq V_{PS}$ since V is pre-solid but not solid (Theorem 3.6). Consequently, $V \subseteq HM \cap V_{PS} \subseteq Mod\{x(yz) \approx (xy)z, xyzt \approx xzyt, x^2 \approx y^2, x^3 \approx y^3\}$.

(ii) \Rightarrow (i): Obviously V is medial and $V \subseteq V_{MPS} \subseteq V_{PS}$. Then by Theorem 3.6 V is not solid. \square

Theorem 5.3 means that V_{MPS} is the greatest pre-solid variety of medial semigroups which is not solid.

REFERENCES

1. Denecke, K. and Koppitz, J., *Hyperassociative semigroups*, Semigroup Forum, Vol. **49**,(1994) 41-48.
2. Denecke, K., Lau, D., Pöschel, R. and Schweigert, D., *Hyperidentities, hyperequational classes and clone congruences*, Contributions to General Algebra 7, Verlag Hölder-Pichler-Tempsky, Wien 1991 - Verlag B.G. Teubner, Stuttgart (1991) 97-118.
3. Denecke, K. and Wismath, S. L., *Solid varieties of semigroups*, Semigroup Forum, Vol. **48**, (1994) 219-234.
4. Denecke, K. and Płonka, J., *On regularizations and normalizations of solid varieties*, in: General Algebra and Discrete Mathematics, Berlin 1995, 83-92.
5. Denecke, K. and Koppitz, J., *Pre-solid varieties of commutative semigroups*, preprint 1993.
6. Denecke, K., Lau, D., Pöschel, R. and Schweigert, D., *Free Clones and Solid Varieties*, preprint 1993.
7. Denecke, K., *Pre-solid varieties*, Demonstratio Mathematica, Vol. **27**, 3-4 (1994), 741-750.

8. Evans, T., *The lattice of semigroup varieties*, Semigroup Forum Vol.2, No 1 (1971) 1-43.
9. Graczyńska, E., *On Normal and Regular Identities and Hyperidentities*, Universal and Applied Algebra, Proceedings of the V Universal Algebra Symposium, Turawa, Poland, 3-7 May, 1988, World Scientific, Singapore, New Jersey, London, Hong Kong (1989) 107-135.
10. Graczyńska, E. and Schweigert, D., *Hyperidentities of a given type*, Algebra Universalis **27**(1990), 305-318.
11. Mal'cev, I. A., *Iterative Post's Algebras (russian)* Novosibirsk 1976.
12. Plonka, J., *On hyperidentities of some varieties*, preprint 1993.
13. Pöschel, R. and Reichel, M., *Projection algebras and rectangular algebras*, General Algebra and Applications, Heldermann-Verlag, Berlin, (1993) 180-194.
14. Taylor, W., *Hyperidentities and hypervarieties*, Aequationes Mathematicae **23**(1981), 111-127.
15. Wismath, S., *Hyperidentity Bases for rectangular bands and other semigroup varieties*, to appear in Journal of the Australian Math. Society.

K. DENECKE, J. KOPFITZ
UNIVERSITY OF POTSDAM
INSTITUTE OF MATHEMATICS
14415 POTSDAM
AM NEUEN PALAIS
E-mail address: kdenecke@hp.rz.uni-potsdam.de