

Jong Taek Cho

A contact metric manifold satisfying a certain curvature condition

*Archivum Mathematicum*, Vol. 31 (1995), No. 4, 319--333

Persistent URL: <http://dml.cz/dmlcz/107554>

## Terms of use:

© Masaryk University, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**A CONTACT METRIC MANIFOLD SATISFYING  
A CERTAIN CURVATURE CONDITION**

JONG TAEK CHO

**ABSTRACT.** In the present paper we investigate a contact metric manifold satisfying (C)  $(\bar{\nabla}_{\dot{\gamma}}R)(\cdot, \dot{\gamma})\dot{\gamma} = 0$  for any  $\bar{\nabla}$ -geodesic  $\gamma$ , where  $\bar{\nabla}$  is the Tanaka connection. We classify the 3-dimensional contact metric manifolds satisfying (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ . Also, we prove a structure theorem for a contact metric manifold with  $\xi$  belonging to the  $k$ -nullity distribution and satisfying (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ .

1. INTRODUCTION

A Riemannian manifold  $M = (M, g)$  with Riemannian metric tensor  $g$  is called (E.Cartan [6]) a locally symmetric space if  $M$  satisfies  $\nabla R = 0$ , where  $\nabla$  is the Levi-Civita connection. In [1] a locally symmetric space  $M$  is characterized by the remarkable property that the Jacobi operator field  $R_{\dot{\gamma}} = R(\cdot, \dot{\gamma})\dot{\gamma}$  is diagonalizable by a  $\nabla$ -parallel orthonormal frame field along  $\gamma$  and their eigenvalues are constant along  $\gamma$  for any geodesic  $\gamma$  on  $M$ .

On the other hand, T.Takahashi ([11]) introduced the notion of Sasakian locally  $\phi$ -symmetric spaces which may be considered as the analogues of locally Hermitian symmetric spaces. A contact metric locally  $\phi$ -symmetric space is defined as a generalization of the notion of the Sasakian locally  $\phi$ -symmetric spaces and investigated by D.E.Blair ([3]).

In [9], we have introduced a class of contact metric manifolds satisfying

$$(C) \quad (\bar{\nabla}_{\dot{\gamma}}R)(\cdot, \dot{\gamma})\dot{\gamma} = 0$$

for any unit  $\bar{\nabla}$ -geodesic  $\gamma(\bar{\nabla}_{\dot{\gamma}}\dot{\gamma} = 0)$ , where  $\bar{\nabla}$  is a linear connection such that the structure tensors are parallel. We note that the connection coincides with the Tanaka connection ([13]) on a strongly pseudo-convex integrable CR-manifold whose structure is determined by a given contact metric structure, particularly for 3-dimensional contact metric manifolds and contact metric manifolds with

---

1991 *Mathematics Subject Classification*: 53C15, 53C35.

*Key words and phrases*: contact metric manifolds, Tanaka connection, Jacobi operator.

This work was partially supported by TGRC-KOSEF.

Received August 21, 1995.

the structure vector field  $\xi$  belonging to the  $k$ -nullity distribution (see section 1), and also note that the geodesics of the Levi-Civita connection and the Tanaka connection do not coincide in general. We easily observe that a contact metric manifold satisfies the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$  if and only if the Jacobi operator field  $R_{\dot{\gamma}}$  is diagonalizable by a  $\bar{\nabla}$ -parallel orthonormal frame field along  $\gamma$  and their eigenvalues are constant along  $\gamma$  for any  $\bar{\nabla}$ -geodesic  $\gamma$  in the manifold.

The present paper is a continuation of the preceding papers [8], [9] in which we proved that a 3-dimensional contact metric manifold satisfying the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$  is locally  $\phi$ -symmetric (in the sense of D.E.Blair). In the present paper, we determine all 3-dimensional contact metric manifolds satisfying the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ . Namely, we prove

**Theorem A.** *Let  $M$  be a 3-dimensional contact metric manifold. If  $M$  satisfies the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ , then  $M$  is a Sasakian locally  $\phi$ -symmetric or a contact metric manifold of constant sectional curvature.*

It was proved ([5]) that a 3-dimensional Sasakian  $\phi$ -symmetric space (simply connected and complete Sasakian locally  $\phi$ -symmetric space) is isometric to the unit sphere  $S^3$  in  $\mathbb{E}^4$ ,  $SU(2)$ , the universal covering space  $\widetilde{SL(2, \mathbb{R})}$  of  $SL(2, \mathbb{R})$  or the Heisenberg group  $H$ , each with a special left invariant metric (see [15]). Also, it was proved ([4]) recently that a 3-dimensional contact metric locally symmetric space is of constant sectional curvature 0 or 1. Thus from Theorem A we have

**Corollary B.** *Let  $M$  be a simply connected and complete 3-dimensional contact metric manifold. If  $M$  satisfies the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ , then  $M$  is isometric to the unit sphere  $S^3$  in  $\mathbb{E}^4$ ,  $SU(2)$ , the universal covering space  $\widetilde{SL(2, \mathbb{R})}$  of  $SL(2, \mathbb{R})$  or the Heisenberg group  $H$ , each with a special left invariant metric, or the Euclidean space  $\mathbb{E}^3$ .*

A contact metric on  $\mathbb{E}^3$ , for example, is explicitly expressed as  $\mathbb{R}^3(x^1, x^2, x^3)$  with  $\eta = \frac{1}{2}(\cos x^3 dx^1 + \sin x^3 dx^2)$  and  $g_{ij} = \frac{1}{4}\delta_{ij}$ . Also, in section 3 we prove that

**Theorem C.** *Let  $M^{2n+1}(n \geq 2)$  be a contact metric manifold with  $\xi$  belonging to the  $k$ -nullity distribution. If  $M$  satisfies the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ , then  $M$  is a Sasakian locally  $\phi$ -symmetric space or  $M$  is locally the product of a flat  $(n + 1)$ -dimensional manifold and an  $n$ -dimensional manifold of positive constant sectional curvature equal to 4.*

We remark that a contact manifold  $M^{2n+1}(n \geq 2)$  can not admit a contact metric structure of vanishing curvature (cf. pp. 115 in [2]). All manifolds in the present paper are assumed to be connected and of class  $C^\infty$ .

The author thanks to Professor K.Sekigawa and L.Vanhecke for their advices and constant encouragements.

## 2. PRELIMINARIES

A  $(2n + 1)$ -dimensional manifold  $M^{2n+1}$  is said to be a contact manifold if it admits a global 1-form  $\eta$  such that  $\eta \wedge (d\eta)^n \neq 0$  everywhere. Given a contact

form  $\eta$ , we have a unique vector field  $\xi$ , which is called the characteristic vector field, satisfying  $\eta(\xi) = 1$  and  $d\eta(\xi, X) = 0$  for any vector field  $X$ . It is well-known that there exists a Riemannian metric  $g$  and a  $(1, 1)$ -tensor field  $\phi$  such that

$$(2.1) \quad \eta(X) = g(X, \xi), \quad d\eta(X, Y) = g(X, \phi Y), \quad \phi^2 X = -X + \eta(X)\xi,$$

where  $X$  and  $Y$  are vector fields on  $M$ . From (2.1) it follows that

$$(2.2) \quad \phi\xi = 0, \quad \eta \circ \phi = 0, \quad g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y).$$

A Riemannian manifold  $M$  equipped with structure tensors  $(\phi, \xi, \eta, g)$  satisfying (2.1) is said to be a contact metric manifold and is denoted by  $M = (M, \phi, \xi, \eta, g)$ . Given a contact metric manifold  $M$ , following D.E.Blair([2]), we define a  $(1, 1)$ -tensor field  $h$  by  $h = -\frac{1}{2}L_\xi\phi$ , where  $L$  denotes Lie differentiation. Then we may observe that  $h$  is symmetric and satisfies

$$(2.3) \quad h\xi = 0 \quad \text{and} \quad h\phi = -\phi h,$$

$$(2.4) \quad \nabla_X \xi = -\phi X - \phi hX,$$

where  $\nabla$  is Levi-Civita connection. From (2.3) and (2.4) we see that each trajectory of  $\xi$  is a geodesic. We denote by  $R$  Riemannian curvature tensor defined by

$$R(X, Y)Z = \nabla_X(\nabla_Y Z) - \nabla_Y(\nabla_X Z) - \nabla_{[X, Y]}Z$$

for all vector fields  $X, Y, Z$ . Along a trajectory of  $\xi$ , the Jacobi operator  $R_\xi = R(\cdot, \xi)\xi$  is a symmetric  $(1, 1)$ -tensor field. We have

$$(2.5) \quad (\text{trace } R_\xi) = g(Q\xi, \xi) = 2n - (\text{trace } h^2),$$

$$(2.6) \quad \nabla_\xi h = \phi - \phi R_\xi - \phi h^2,$$

(cf.[2] or [3]) where  $Q$  is Ricci  $(1, 1)$ -tensor on  $M$ .

A contact metric manifold for which  $\xi$  is Killing is called a  $K$ -contact metric manifold. It is easy to see that a contact metric manifold is  $K$ -contact if and only if  $h = 0$ . For a contact metric manifold  $M$  one may define naturally an almost complex structure on  $M \times \mathbb{R}$ . If this almost complex structure is integrable,  $M$  is said to be Sasakian. A Sasakian manifold is characterized by a condition

$$(2.7) \quad (\nabla_X \phi)Y = g(X, Y)\xi - \eta(Y)X$$

for all vector fields  $X$  and  $Y$  on the manifold.

Let  $M$  be a contact metric manifold. It is well-known that  $M$  is Sasakian if and only if

$$(2.8) \quad R(X, Y)\xi = \eta(Y)X - \eta(X)Y$$

for all vector fields  $X$  and  $Y$  ([2]).

Let  $T$  be a  $(1, 2)$ -tensor field on  $M$  defined by

$$(2.9) \quad T_X Y = -\frac{1}{2}\phi(\nabla_X \phi)Y + \frac{1}{2}\eta(Y)(\phi X + \phi hX) - \eta(X)\phi Y - g(\phi X + \phi hX, Y)\xi.$$

Particularly, for a Sasakian manifold, from (2.7) and (2.9) we see that

$$(2.10) \quad T_X Y = g(X, \phi Y)\xi + \eta(Y)\phi X - \eta(X)\phi Y,$$

where  $X$  and  $Y$  are vector fields on  $M$ . Using the tensor field  $T$ , we define a linear connection  $\bar{\nabla}$  on  $M$  by

$$(2.11) \quad \bar{\nabla}_X Y = \nabla_X Y + T_X Y$$

(cf. [7] or [8]). Then the linear connection  $\bar{\nabla}$  has the torsion given by  $T_X Y - T_Y X$ . Using (2.1), (2.2) and (2.3), we have

$$(2.12) \quad \bar{\nabla}\phi = 0, \quad \bar{\nabla}\xi = 0, \quad \bar{\nabla}\eta = 0, \quad \bar{\nabla}g = 0.$$

We remark that the above connection  $\bar{\nabla}$  coincides with the Tanaka connection (defined in [12]) on a strongly pseudo-convex integrable CR-manifold whose structure is determined by a contact metric manifold which satisfies  $(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$  for any vector fields  $X$  and  $Y$  (see Proposition 2.1 in [15]). The tangent space  $T_p M$  of  $M$  at each point  $p \in M$  is decomposed as  $T_p M = \mathfrak{D}_p \oplus \{\xi\}_p$  (direct sum), where we denote  $\mathfrak{D}_p = \{v \in T_p M \mid \eta(v) = 0\}$ . Then  $\mathfrak{D} : p \rightarrow \mathfrak{D}_p$  defines a distribution orthogonal to  $\xi$ . Let  $\gamma$  be a  $\bar{\nabla}$ -geodesic parametrized with the arc-length parameter  $s$ , where a  $\bar{\nabla}$ -geodesic means a geodesic with respect to  $\bar{\nabla}$ . From (2.9) and (2.11) we see that a  $\bar{\nabla}$ -geodesic does not coincide with a  $\nabla$ -geodesic in general. We denote  $\dot{\gamma} = \gamma_*\left(\frac{d}{ds}\right)$  and by  $\gamma_*$  the differential of  $\gamma : I \rightarrow M$ . Define the Jacobi operator  $R_{\dot{\gamma}}$  by  $R_{\dot{\gamma}} = R(\cdot, \dot{\gamma})\dot{\gamma}$  along  $\gamma$ .  $R_{\dot{\gamma}}$  is a symmetric  $(1, 1)$ -tensor field along  $\gamma$ . Moreover, from (2.12) we observe that  $\eta(\dot{\gamma})$  is constant along  $\gamma$ , and thus a  $\bar{\nabla}$ -geodesic whose tangent initially belongs to  $\mathfrak{D}$  remains in  $\mathfrak{D}$ . We call such a  $\bar{\nabla}$ -geodesic which is tangent to  $\mathfrak{D}$  a *horizontal  $\bar{\nabla}$ -geodesic*.

Now, recall the definition of a Sasakian locally  $\phi$ -symmetric space ([11]).

**Definition 2.1.** A Sasakian manifold  $M = (M, \phi, \xi, \eta, g)$  is said to be locally  $\phi$ -symmetric if  $\phi^2(\nabla_V R)(X, Y)Z = 0$  for all vector fields  $V, X, Y, Z \in \mathfrak{D}$ .

As a generalization of the above Sasakian one, a contact metric locally  $\phi$ -symmetric space is defined by D.E.Blair([3]) by the same condition. In [7] we characterized a Sasakian locally  $\phi$ -symmetric space by following

**Theorem 2.2.** A Sasakian manifold  $M$  is locally  $\phi$ -symmetric if and only if  $M$  satisfies the condition (C) for any horizontal  $\bar{\nabla}$ -geodesic.

Concerning Theorem 2.2 we prove

**Theorem 2.3.** *A Sasakian manifold  $M$  is locally  $\phi$ -symmetric if and only if  $M$  satisfies the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ .*

**Proof.** From (2.8) and (2.12) we see that

$$(\bar{\nabla}_\xi R)(Y, X)\xi = 0$$

for all vector fields  $X$  and  $Y$  on  $M$ . Then, taking account of Theorem 2.2, it suffices to prove  $g((\bar{\nabla}_\xi R)(Y, V)V, X) = 0$  for all vector fields  $V, X, Y \in \mathfrak{D}$ . It follows from (2.10) and (2.11) that

$$(2.13) \quad \begin{aligned} g((\bar{\nabla}_\xi R)(Y, V)V, X) = & (\nabla_\xi R)(Y, V)V, X) - g(\phi R(Y, V)V, X) + g(R(\phi Y, V)V), X) \\ & + g(R(X, \phi V)V, Y) + g(R(X, V)\phi V, Y) \end{aligned}$$

for all vector fields  $V, X, Y \in \mathfrak{D}$ . From (2.8) and the second Bianchi identity, we have

$$(2.14) \quad \begin{aligned} ((\bar{\nabla}_\xi R)(Y, V)V, X) = & g(\phi Y, V)g(V, X) - g(\phi Y, X)g(V, V) + g(R(V, X)\phi Y, V) \\ & + g(\phi V, X)g(V, Y) - g(R(V, X)\phi V, Y). \end{aligned}$$

Thus, from (2.13) and (2.14), we have

$$(2.15) \quad \begin{aligned} ((\bar{\nabla}_\xi R)(Y, V)V, X) = & g(\phi Y, V)g(V, X) - g(\phi Y, X)g(V, V) + 2g(R(V, X)\phi Y, V) \\ & + g(\phi V, X)g(V, Y) - 2g(R(V, X)\phi V, Y) \\ & + g(R(Y, V)\phi V, X) - g(\phi R(Y, V)V, X) \end{aligned}$$

for all vector fields  $V, X, Y \in \mathfrak{D}$ . From the definition of the curvature tensor, taking account of (2.4) and (2.7), we obtain

$$(2.16) \quad R(Y, X)\phi Z - \phi R(Y, X)Z = g(\phi Y, Z)X - g(X, Z)\phi Y - g(\phi X, Z)Y + g(Y, Z)\phi X,$$

where  $X, Y$  and  $Z$  are vector fields on  $M$ . By using (2.16), from (2.15) we see that  $g((\bar{\nabla}_\xi R)(Y, V)V, X) = 0$  for all vector fields  $V, X, Y \in \mathfrak{D}$ . □

S. Tanno ([13]) defined the  $k$ -nullity distribution of Riemannian manifold  $(M, g)$ , for a real number  $k$ , by

$$N(k) : p \rightarrow N_p(k) = \{z \in T_p M \mid R(x, y)z = k(g(y, z)x - g(x, z)y) \text{ for any } x, y \in T_p M\},$$

and he proved

**Proposition 2.4.** *Let  $M = (M, \phi, \xi, \eta, g)$  be a contact metric manifold. If  $\xi$  belong to the  $k$ -nullity distribution, then  $k \leq 1$ . If  $k < 1$ , then  $M$  admits three mutually orthogonal and integral distributions  $D(0)$ ,  $D(\lambda)$  and  $D(-\lambda)$ , defined by the eigenspaces of  $h$ , where  $\lambda = \sqrt{1-k}$ .*

In [8], we proved

**Theorem 2.5.** *Let  $M$  be a contact metric manifold with  $\xi$  belonging to the  $k$ -nullity distribution. Then  $M$  is locally  $\phi$ -symmetric (in the sense of D.E.Blair) if and only if  $M$  satisfies the condition (C) for any horizontal  $\bar{\nabla}$ -geodesic.*

Since a contact metric manifold  $M$  with  $\xi$  belonging to the 1-nullity distribution is a Sasakian manifold, the above Theorem 2.5 is an extension of Theorem 2.2. For a contact metric manifold with  $\xi$  belonging to the 0-nullity distribution, D.E. Blair ([2]) proved

**Theorem 2.6.** *Let  $M$  be a contact metric manifold with  $\xi$  belonging to the 0-nullity distribution. Then  $M$  is locally the product of a flat  $(n + 1)$ -dimensional manifold and an  $n$ -dimensional manifold of positive constant sectional curvature equal to 4.*

### 3. 3-DIMENSIONAL CONTACT METRIC MANIFOLDS

In this section we prove Theorem A. Recently, it was proved in [14] that a 3-dimensional contact metric manifold always satisfies

$$(3.1) \quad (\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX)$$

for all vector fields  $X, Y$ .

**Lemma 3.1.** *A 3-dimensional contact metric manifold is Sasakian if and only if  $h = 0$ .*

**Proof.** Assume that  $M^3$  is a contact metric manifold. Then from (2.7) and (3.1) we get  $g(hX, Y)\xi - \eta(Y)hX = 0$ . Taking account of (2.3), we have  $g(hX, Y) = 0$  for all vector fields  $X, Y$  on  $M$  and hence, we have  $h = 0$ . The converse is obvious.  $\square$

Now we prove Theorem A.

**Proof of Theorem A.** Let  $M^3 = (M^3, \phi, \xi, \eta, g)$  be a 3-dimensional contact metric manifold satisfying the condition (C) for any  $\bar{\nabla}$ -geodesic  $\gamma$ . It is well-known that the curvature tensor  $R$  of a 3-dimensional Riemannian manifold is expressed by

$$(3.2) \quad R(Y, X)Z = \rho(X, Z)Y - \rho(Y, Z)X + g(X, Z)QY - g(Y, Z)QX - \frac{1}{2}\tau\{g(X, Z)Y - g(Y, Z)X\}$$

for all vector fields  $X, Y, Z$ , where  $\rho(Y, X) = g(QY, X)$  and  $\tau$  is the scalar curvature of the manifold.

From (3.2) and the assumption we have

$$\begin{aligned}
 (3.3) \quad 0 &= (\bar{\nabla}_x R)(y, x)x \\
 &= (\bar{\nabla}_x \rho)(x, x)y - (\bar{\nabla}_x \rho)(y, x)x + g(x, x)(\bar{\nabla}_x Q)y - g(y, x)(\bar{\nabla}_x Q)x \\
 &\quad - \frac{1}{2}(x\tau)\{g(x, x)y - g(y, x)x\},
 \end{aligned}$$

for any  $x, y \in T_p M$  and any  $p \in M$ . For any unit  $v$  orthogonal to  $\xi$ , let  $\{v, \phi v, \xi\}$  be an adapted orthonormal basis of  $T_p M$  ( $p \in M$ ). Then from (3.3) we get  $g((\bar{\nabla}_x R)(v, x)x, v) = 0$ ,  $g((\bar{\nabla}_x R)(\phi v, x)x, \phi v) = 0$  and  $g((\bar{\nabla}_x R)(\xi, x)x, \xi) = 0$ , and summing up these three equalities, we have

$$(3.4) \quad (\bar{\nabla}_x \rho)(x, x) = 0.$$

Also, from (3.3) we get  $(\bar{\nabla}_v R)(\phi v, v)v = 0$ ,  $(\bar{\nabla}_v R)(\xi, v)v = 0$  and thus we have

$$(3.5) \quad (\bar{\nabla}_v \rho)(\phi v, \phi v) = (\bar{\nabla}_v \rho)(\xi, \xi)$$

and

$$(3.6) \quad (\bar{\nabla}_v \rho)(\phi v, \xi) = 0.$$

Taking account of (3.1), we see that

$$(3.7) \quad T_x y = \eta(y)(\phi x + \phi h x) - \eta(x)\phi y - g(\phi x + \phi h x, y)\xi$$

for  $x, y \in T_p M$  and  $p \in M$ . From (2.11) and (3.7) we have the formulas (3.8),(3.9) and (3.10) which are equivalent to (3.4),(3.5) and (3.6), respectively:

$$(3.8) \quad (\nabla_x \rho)(x, x) = 2\{\eta(x)\rho(\phi h x, x) - g(\phi h x, x)\rho(\xi, x)\},$$

$$(3.9) \quad (\nabla_v \rho)(\xi, \xi) - (\nabla_v \rho)(\phi v, \phi v) = 2\{(2 + g(hv, v))\rho(\xi, \phi v) + \rho(\phi hv, \xi)\},$$

$$(3.10) \quad (\nabla_v \rho)(\phi v, \xi) = \rho(\phi v, \phi v) + \rho(\phi v, \phi h v) - \{1 + g(hv, v)\}\rho(\xi, \xi)$$

for any unit  $x \in T_p M$  and unit vector  $v$  orthogonal to  $\xi$ .

Let  $W$  be the subset of  $M$  on which the number of distinct eigenvalues of  $h$  is constant. Then  $W$  is an open and dense subset of  $M$ . We fix any point  $q$  in  $W$ . Then from (2.3) there exists a  $C^\infty$  function  $\lambda$  such that  $he_1 = \lambda e_1$ ,  $he_2 = -\lambda e_2$ ,  $h\xi = 0$  where  $\{e_1, e_2, \xi\}$  is a local orthonormal frame field on a neighborhood  $N_q \subset W$  containing  $q$ . We denote  $\Gamma_{ijk} = g(\nabla_{e_i} e_j, e_k)$ ,

$\rho_{ij} = \rho(e_i, e_j)$ ,  $\nabla_i \rho_{jk} = (\nabla_{e_i} \rho)(e_j, e_k)$  and  $\nabla_h R_{ijkl} = g((\nabla_h R)(e_i, e_j)e_k, e_l)$  for  $h, i, j, k, l = 1, 2, 3$ . Then from (2.4) we get

$$(3.11) \quad \Gamma_{132} = -(1 + \lambda), \quad \Gamma_{231} = 1 - \lambda$$

and

$$(3.12) \quad \Gamma_{131} = \Gamma_{232} = 0.$$

Also, from (2.6) and taking account of (2.5) and (3.2), we have

$$(3.13) \quad \xi \lambda = \rho_{12}$$

and

$$(3.14) \quad 4\lambda \Gamma_{312} = \rho_{22} - \rho_{11}.$$

Moreover, from (3.8) we get

$$(3.15) \quad \nabla_1 \rho_{11} = 0, \quad \nabla_2 \rho_{22} = 0$$

and

$$(3.16) \quad \nabla_3 \rho_{33} = 0.$$

Substituting  $x = \frac{1}{\sqrt{2}}(e_1 + e_2)$  and  $x = \frac{1}{\sqrt{2}}(e_1 - e_2)$ , respectively in (3.8) and taking account of (3.15), we have

$$2\nabla_1 \rho_{12} + 2\nabla_2 \rho_{12} + \nabla_1 \rho_{22} + \nabla_2 \rho_{11} = -4\lambda(\rho_{31} + \rho_{32})$$

and

$$-2\nabla_1 \rho_{12} + 2\nabla_2 \rho_{12} + \nabla_1 \rho_{22} - \nabla_2 \rho_{11} = 4\lambda(\rho_{31} - \rho_{32}).$$

By summing these two equalities, we have

$$(3.17) \quad \nabla_1 \rho_{22} + 2\nabla_2 \rho_{12} = -4\lambda \rho_{23}$$

and subtracting (3.17) from the preceding one, we have

$$(3.18) \quad \nabla_2 \rho_{11} + 2\nabla_1 \rho_{12} = -4\lambda \rho_{13}.$$

Also, substituting  $x = \frac{1}{\sqrt{2}}(e_1 + e_3)$  and  $x = \frac{1}{\sqrt{2}}(e_1 - e_3)$ , respectively in (3.8) and taking account of (3.16), we have

$$2\nabla_1 \rho_{13} + 2\nabla_3 \rho_{31} + \nabla_1 \rho_{33} + \nabla_3 \rho_{11} = 2\lambda \rho_{23}$$

and

$$-2\nabla_1\rho_{13} + 2\nabla_3\rho_{31} + \nabla_1\rho_{33} - \nabla_3\rho_{11} = 2\lambda\rho_{23}.$$

Summing these two equalities we have

$$(3.19) \quad \nabla_1\rho_{33} + 2\nabla_3\rho_{13} = 2\lambda\rho_{23}$$

and subtracting (3.19) from the preceding one, we have

$$(3.20) \quad \nabla_3\rho_{11} + 2\nabla_1\rho_{31} = 0.$$

A similar calculation for  $x = \frac{1}{\sqrt{2}}(e_2 + e_3)$  and  $x = \frac{1}{\sqrt{2}}(e_2 - e_3)$  gives

$$(3.21) \quad \nabla_2\rho_{33} + 2\nabla_3\rho_{23} = 2\lambda\rho_{13}$$

and

$$(3.22) \quad \nabla_3\rho_{22} + 2\nabla_2\rho_{32} = 0.$$

On the one hand, from the second Bianchi identity, we have

$$(3.23) \quad 2\nabla_2\rho_{12} + 2\nabla_3\rho_{13} - \nabla_1\rho_{22} - \nabla_1\rho_{33} = 0.$$

$$(3.24) \quad 2\nabla_1\rho_{21} + 2\nabla_3\rho_{23} - \nabla_2\rho_{11} - \nabla_2\rho_{33} = 0.$$

From (3.17), (3.19) and (3.23) (resp.(3.18), (3.21) and (3.24)), we have (3.25) (resp.(3.26)):

$$(3.25) \quad \nabla_1\rho_{22} + \nabla_1\rho_{33} = -\lambda\rho_{23},$$

$$(3.26) \quad \nabla_2\rho_{11} + \nabla_2\rho_{33} = -\lambda\rho_{13}.$$

On the other hand, from (3.5) we have

$$(3.27) \quad \nabla_1\rho_{33} - \nabla_1\rho_{22} = 4(\lambda + 1)\rho_{23}$$

and

$$(3.28) \quad \nabla_2\rho_{33} - \nabla_2\rho_{11} = 4(\lambda - 1)\rho_{13}.$$

Thus, from (3.25)-(3.28) we have

$$(3.29) \quad \nabla_1\rho_{33} = \frac{1}{2}(3\lambda + 4)\rho_{23}, \quad \nabla_2\rho_{33} = \frac{1}{2}(3\lambda - 4)\rho_{13}$$

and

$$(3.30) \quad \nabla_1\rho_{22} = -\frac{1}{2}(5\lambda + 4)\rho_{23}, \quad \nabla_2\rho_{11} = -\frac{1}{2}(5\lambda - 4)\rho_{13}.$$

Also, from (3.17),(3.18) and (3.30), we have

$$(3.31) \quad \nabla_1\rho_{12} = -\frac{1}{4}(3\lambda + 4)\rho_{13} \quad \text{and} \quad \nabla_2\rho_{21} = -\frac{1}{4}(3\lambda - 4)\rho_{23}.$$

**Lemma 3.2.**  $\rho_{ij} = 0$  on  $N_q(\subset W)$ , where  $i \neq j, i, j = 1, 2, 3$ .

**Proof.** . Differentiating (2.5) in the direction  $\xi$  and taking account of (3.16) we have  $\xi\lambda = 0$ . Thus from (3.13) we have  $\rho_{12} = 0$  on  $N_q$ .

Now we prove  $\rho_{13} = 0$  and  $\rho_{23} = 0$  on  $N_q$ . Differentiating (2.5) in the directions  $e_1$  and  $e_2$  and taking account of (3.11), (3.12) and (3.29) we have

$$(3.32) \quad \rho_{23} = 8(e_1\lambda)$$

and

$$(3.33) \quad \rho_{13} = 8(e_2\lambda),$$

respectively.

Also, differentiating  $\rho_{12} = 0$  in the direction  $\xi$ , we have

$$(3.34) \quad \nabla_3\rho_{12} = \Gamma_{312}(\rho_{11} - \rho_{22}).$$

Substituting  $x = \xi$  in (3.3), we get  $\bar{\nabla}_3\rho_{12} = 0$ , and from (3.7) we get  $\bar{\nabla}_3\rho_{12} = \nabla_3\rho_{12} + \rho_{22} - \rho_{11}$ . Thus we see that

$$(3.35) \quad \nabla_3\rho_{12} = \rho_{11} - \rho_{22}.$$

At first, if there exists a point in  $N_q(\subset W)$  such that  $\rho_{11} = \rho_{22}$ , then  $\rho_{13} = \rho_{23} = 0$  at that point. In fact, differentiating  $\rho_{12} = 0$  in the direction  $e_1$  and  $e_2$ , then from the assumption and (3.11) we have  $\nabla_1\rho_{12} = -(1 + \lambda)\rho_{13}$  and  $\nabla_2\rho_{21} = (1 - \lambda)\rho_{23}$ , respectively. Thus taking account of (3.31) we have  $\rho_{13} = \rho_{23} = 0$  at the point in  $N_q$ . Next, suppose there exists a point  $m$  such that  $\rho_{11}(m) \neq \rho_{22}(m)$ . Then we see that  $\rho_{11} \neq \rho_{22}$  on a sufficiently small neighborhood  $U(m)$  of  $m$ . From (3.34) and (3.35) we get  $\Gamma_{312} = 1$  on  $U(m)$ . Thus (3.14) becomes  $4\lambda = \rho_{22} - \rho_{11}$  on  $U(m)$ . Differentiating this equation in the directions  $e_1$  and  $e_2$  and taking account of (3.11), (3.12), (3.32) and (3.33) we have  $\nabla_1\rho_{22} = -\frac{1}{2}(4\lambda + 3)\rho_{23}$  and  $\nabla_2\rho_{11} = -\frac{1}{2}(4\lambda - 3)\rho_{13}$ . Thus taking account of (3.30) we have

$$(3.36) \quad (\lambda + 1)\rho_{23} = 0$$

and

$$(3.37) \quad (\lambda - 1)\rho_{13} = 0$$

on  $U(m)$ . Differentiating (3.36)(resp.(3.37)) in the direction  $e_1$ (resp. $e_2$ ) and taking account of (3.32) and (3.33), we have

$$(3.38) \quad \frac{1}{8}\rho_{23}^2 + (\lambda + 1)(e_1\rho_{23}) = 0$$

and

$$(3.39) \quad \frac{1}{8}\rho_{13}^2 + (\lambda - 1)(\epsilon_2\rho_{13}) = 0$$

on  $U(m)$ . If there exists a point  $n$  in  $U(m)$  such that  $\rho_{13}(n) \neq 0$ , then from (3.37) we get  $\lambda(n) = 1$ , and from (3.39) we get  $\rho_{13}(n) = 0$ , a contradiction. Also, if there exist a point  $n$  in  $U(m)$  such that  $\rho_{23}(n) \neq 0$ , then from (3.36) we get  $\lambda(n) = -1$ , and from (3.38) we get  $\rho_{23}(n) = 0$ , a contradiction. Thus we have  $\rho_{13} = \rho_{23} = 0$  on  $U(m)$ . At last, we conclude that  $\rho_{13} = \rho_{23} = 0$  also on  $N_q$ .  $\square$

From Lemma 3.2, we see that  $\lambda$  is locally constant on  $N_q(\subset W)$ . Since  $\rho_{13} = \rho_{23} = 0$ , from (3.29)-(3.31), we get

$$(3.40) \quad \begin{aligned} \nabla_1\rho_{12} &= 0, \quad \nabla_1\rho_{22} = 0, \quad \nabla_1\rho_{33} = 0, \\ \nabla_2\rho_{12} &= 0, \quad \nabla_2\rho_{11} = 0, \quad \nabla_2\rho_{33} = 0. \end{aligned}$$

Also, taking account of (3.12), we have

$$(3.41) \quad \nabla_1\rho_{13} = 0 \quad \text{and} \quad \nabla_2\rho_{23} = 0.$$

The equations (3.19)-(3.22), together with (3.40) and (3.41), yield

$$(3.42) \quad \nabla_3\rho_{11} = 0, \quad \nabla_3\rho_{13} = 0, \quad \nabla_3\rho_{22} = 0, \quad \nabla_3\rho_{23} = 0.$$

From (3.15), (3.16), (3.40) and (3.42), we see that the scalar curvature  $\tau$  is constant. Returning to the condition (C), from (3.3), by using polarization, we have

$$(3.43) \quad \begin{aligned} 0 = & S_{x,z,w} [(\nabla_x\rho)(z,w)y + \eta(Qz)g(\phi x + \phi hx, w)y - \eta(x)g(\phi Qz, w)y \\ & - g(\phi x + \phi hx, Qz)\eta(w)y - \eta(z)g(Q\phi x + Q\phi hx, w)y + \eta(x)g(Q\phi z, w)y \\ & + g(\phi x + \phi hx, z)\eta(Qw)y - (\nabla_x\rho)(y,z)w - \eta(Qw)g(\phi x + \phi hx, y)z \\ & + \eta(x)g(\phi Qw, y)z + g(\phi x + \phi hx, Qw)\eta(y)z + \eta(w)g(Q\phi x + Q\phi hx, y)z \\ & - \eta(x)g(Q\phi w, y)z - g(\phi x + \phi hx, w)\eta(Qy)z + g(x,z)\{(\nabla_w Q)y \\ & + \eta(Qy)(\phi w + \phi hw) - \eta(w)\phi Qy - g(\phi w + \phi hw, Qy)\xi \\ & - \eta(y)(Q\phi w + Q\phi hw) + \eta(w)Q\phi y + g(\phi w + \phi hw, y)Q\xi\} - g(y,x)\{(\nabla_z Q)w \\ & + \eta(Qw)(\phi z + \phi hz) - \eta(z)\phi Qw - g(\phi z + \phi hz, Qw)\xi \\ & - \eta(w)(Q\phi z + Q\phi hz) + \eta(z)Q\phi w + g(\phi z + \phi hz, w)Q\xi\}] \end{aligned}$$

for any  $x, y, z, w \in T_qM$ , where  $S_{x,z,w}$  denotes the cyclic sum for tangent vectors  $x, z, w$ . First, substitute  $y = e_1, x = e_1, z = e_2, w = e_3$  into (3.43). Then taking account of (3.40) and (3.41) we have

$$(3.44) \quad \begin{aligned} \nabla_1\rho_{23} + \nabla_3\rho_{12} - \nabla_2\rho_{31} \\ - \lambda\rho_{22} + (3\lambda - 1)\rho_{33} - (2\lambda - 1)\rho_{11} = 0. \end{aligned}$$

Next, substitute  $y = e_2$ ,  $x = e_1$ ,  $z = e_2$ ,  $w = e_3$  into (3.43). Then taking account of (3.41) and (3.42) we have

$$(3.45) \quad \begin{aligned} \nabla_1 \rho_{23} + \nabla_2 \rho_{31} - \nabla_3 \rho_{12} \\ - (\lambda - 1)\rho_{11} - (\lambda + 1)\rho_{22} - 2\lambda\rho_{33} = 0. \end{aligned}$$

Finally, substitute  $y = e_3$ ,  $x = e_1$ ,  $z = e_2$ ,  $w = e_3$  into (3.43). Then taking account of (3.40) we have

$$(3.46) \quad \begin{aligned} \nabla_2 \rho_{31} + \nabla_3 \rho_{12} - \nabla_1 \rho_{23} \\ + (\lambda - 1)\rho_{33} + \rho_{22} - \lambda\rho_{11} = 0. \end{aligned}$$

From (3.44), (3.45) and (3.46), we have

$$(3.47) \quad \begin{aligned} 2\nabla_2 \rho_{31} &= (2\lambda - 1)\rho_{11} + \lambda\rho_{22} - (3\lambda - 1)\rho_{33}, \\ 2\nabla_3 \rho_{12} &= (3\lambda - 1)\rho_{11} + (\lambda - 1)\rho_{22} - (4\lambda - 2)\rho_{33}, \\ 2\nabla_1 \rho_{23} &= (3\lambda - 2)\rho_{11} + (2\lambda + 1)\rho_{22} - (5\lambda - 1)\rho_{33}. \end{aligned}$$

Now suppose there exists a point  $m \in N_q$  such that  $\rho_{11}(m) \neq \rho_{22}(m)$ . Then we see that  $\Gamma_{312}(m) = 1$  in the proof of Lemma 3.2, and from (3.10) and (3.14) we obtain

$$(3.48) \quad \begin{aligned} \nabla_2 \rho_{31} &= (\lambda - 1)(\rho_{11} - \rho_{33}), \\ \nabla_3 \rho_{12} &= \rho_{11} - \rho_{22}, \\ \nabla_1 \rho_{23} &= (\lambda + 1)(\rho_{22} - \rho_{33}) \end{aligned}$$

at  $m$ . Thus from (3.47) and (3.48) we have

$$(3.49) \quad \begin{aligned} \rho_{11} + \lambda\rho_{22} - (\lambda + 1)\rho_{33} &= 0, \\ 3(\lambda - 1)\rho_{11} + (\lambda + 1)\rho_{22} - 2(2\lambda - 1)\rho_{33} &= 0, \\ (3\lambda - 2)\rho_{11} - \rho_{22} - 3(\lambda - 1)\rho_{33} &= 0 \end{aligned}$$

at  $m$ . Since  $\rho_{22}(m) - \rho_{11}(m) = 4\lambda(m)$  from (3.14), the above (3.49) gives

$$\begin{aligned} (\lambda + 1)(\rho_{11} - \rho_{33}) &= -4\lambda^2, \\ 2(2\lambda - 1)(\rho_{11} - \rho_{33}) &= -4\lambda(\lambda + 1), \\ 3(\lambda - 1)(\rho_{11} - \rho_{33}) &= 4\lambda \end{aligned}$$

which yields  $\lambda(m) = 0$ . Since  $\lambda$  is locally constant on  $N_q$ , we see that  $\lambda = 0$ . Now, we consider  $\|h\|^2$ . Then  $\|h\|^2 = 2\lambda^2$  is a function on  $M$ , and by the continuity argument we observe that  $h = 0$  on  $M$ . Thus by Lemma 3.1 we see that  $M$  is Sasakian, and by Theorem 3.1 we see that  $M$  is locally  $\phi$ -symmetric.

Or suppose  $\rho_{11} = \rho_{22}$  on  $N_q$ . Then from (3.10) and (3.14) we have

$$(3.50) \quad \begin{aligned} \nabla_2 \rho_{31} &= (\lambda - 1)(\rho_{11} - \rho_{33}), \\ \nabla_3 \rho_{12} &= 0, \\ \nabla_1 \rho_{23} &= (\lambda + 1)(\rho_{22} - \rho_{33}). \end{aligned}$$

From (3.47) and (3.50), we see that

$$\begin{aligned} (\lambda + 1)(\rho_{11} - \rho_{33}) &= 0, \\ 2(2\lambda - 1)(\rho_{11} - \rho_{33}) &= 0, \\ 3(\lambda - 1)(\rho_{11} - \rho_{33}) &= 0 \end{aligned}$$

which yields  $\rho_{11} = \rho_{33}$  on  $N_q$ . In this case, taking account of Lemma 3.2, we see that  $M$  is a Einstein manifold and hence, of constant sectional curvature. At last, we have our conclusion. □

#### 4. A CONTACT METRIC MANIFOLD WITH $\xi$ BELONGING TO THE $k$ -NULLITY DISTRIBUTION

In the present section we prove Theorem C. The following Lemma is known (cf. p. 446-447 in [13] or p. 251 in [10]).

**Lemma 4.1.** *Let  $M = (M^{2n+1}, \phi, \xi, \eta, g)$  be a contact metric manifold with  $\xi$  belonging to the  $k$ -nullity distribution. Then*

$$(4.1) \quad (\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)(X + hX).$$

**Proof of Theorem C.** Let  $M^{2n+1}(n \geq 2)$  be a contact metric manifold with  $\xi$  belonging to the  $k$ -nullity distribution, i.e.,

$$(4.2) \quad R(X, Y)\xi = k(\eta(Y)X - \eta(X)Y),$$

where  $k$  is a real number. From (4.2) we see that

$$(\bar{\nabla}_\xi R)(Y, X)\xi = 0$$

for all vector fields  $X$  and  $Y$  on  $M$ . Thus, by virtue of Theorem 2.5, it only remains to examine  $g((\bar{\nabla}_\xi R)(Y, V)V, X) = 0$  for all vector fields  $V, X, Y \in \mathfrak{D}$ . From (2.9) and (4.1) we get

$$(4.3) \quad T_X Y = \eta(Y)(\phi X + \phi hX) - \eta(X)\phi Y - g(\phi X + \phi hX, Y)\xi.$$

Then it follows from (2.11) and (4.3), together with (2.1) and (2.2), that

$$(4.4) \quad \begin{aligned} g((\bar{\nabla}_\xi R)(Y, V)V, X) &= (\nabla_\xi R)(Y, V)V, X - g(\phi R(Y, V)V, X) + g(R(\phi Y, V)V, X) \\ &\quad + g(R(X, \phi V)V, Y) + g(R(X, V)\phi V, Y) \end{aligned}$$

for all vector fields  $V, X, Y \in \mathfrak{D}$ . On the other hand, from (4.2) and the second Bianchi identity we obtain

$$(4.5) \quad \begin{aligned} g((\nabla_{\xi} R)(Y, V)V, X) = & k\{g(\phi Y, V)g(V, X) + g(\phi hY, V)g(V, X) \\ & - g(\phi Y, X)g(V, V) - g(\phi hY, X)g(V, V)\} \\ & - g(\phi hV, V)g(Y, X) + g(\phi V, X)g(V, Y) \\ & + g(\phi hV, X)g(V, Y)\} \\ & + g(R(V, X)\phi Y, V) + g(R(V, X)\phi hY, V) \\ & - g(R(V, X)\phi V, Y) - g(R(V, X)\phi hV, Y), \end{aligned}$$

where  $X, Y \in \mathfrak{D}$ . From the definition of the curvature tensor, taking account of (2.4) and (4.1), we obtain

$$(4.6) \quad \begin{aligned} & g(R(Y, X)\phi Z, W) - g(\phi R(Y, X)Z, W) \\ & = g(\phi Y + \phi hY, Z)g(X + hX, W) - g(X + hX, Z)g(\phi Y + \phi hY, W) \\ & \quad - g(\phi X + \phi hX, Z)g(Y + hY, W) + g(Y + hY, Z)g(\phi X + \phi hX, W), \end{aligned}$$

where  $X, Y, Z, W \in \mathfrak{D}$ . Since  $g((\bar{\nabla}_{\xi} R)(Y, V)V, X) = 0$ , from (4.4), (4.5) and (4.6), we have

$$(4.7) \quad \begin{aligned} & (k-1)\{g(\phi Y, V)g(X, V) - g(\phi Y, X)g(V, V) + g(\phi V, X)g(V, Y)\} \\ & \quad + (k+3)\{g(\phi hY, V)g(X, V) - g(\phi hY, X)g(V, V) - g(\phi hV, V)g(X, Y) \\ & \quad + g(\phi hV, X)g(V, Y)\} \\ & = g(\phi Y, V)g(hX, V) - g(\phi Y, X)g(hV, V) + g(\phi V, X)g(hV, Y) \\ & \quad - 3\{g(\phi hY, V)g(hX, V) - g(\phi hY, X)g(hV, V) - g(\phi hV, V)g(hX, Y) \\ & \quad + g(\phi hV, X)g(hV, Y)\} + g(R(V, X)\phi hV, Y) - g(R(V, X)\phi hY, V), \end{aligned}$$

for all vector fields  $V, X, Y \in \mathfrak{D}$ . Since  $h$  is symmetric operator and  $2n+1 \geq 5$ , we assume that  $hY = \lambda Y$  and  $hV = \lambda V$ , where  $Y$  and  $V$  are unit and mutually orthogonal. Then from (4.7) we obtain

$$(4.8) \quad \begin{aligned} & (k-1)g(Y, \phi X) + (k+3)\lambda g(Y, \phi X) \\ & = \lambda g(Y, \phi X) - 3\lambda^2 g(Y, \phi X) \\ & \quad + \lambda g(\phi R(V, X)Y, V) - \lambda g(R(V, X)\phi Y, V). \end{aligned}$$

Also, from (4.6) we have

$$(4.9) \quad g(\phi R(V, X)Y, V) - g(\phi R(V, X)\phi Y, V) = (1 - \lambda^2)g(Y, \phi X).$$

The equations (4.8) and (4.9), together with  $\lambda = \sqrt{1-k}$  (by Proposition 2.4), yield

$$\sqrt{1-k} - (1-k) = 0,$$

which yields  $k = 0$  or  $k = 1$ . Thus we see that  $M$  is Sasakian (when  $k = 1$ ) or  $M$  is a contact metric manifold whose structure vector  $\xi$  belongs to the 0-nullity distribution. Therefore by virtue of Theorems 2.3 and 2.6, we have our conclusion.  $\square$

#### REFERENCES

- [1] Berndt, J. and Vanhecke, L., *Two natural generalizations of locally symmetric spaces*, Diff. Geom. Appl. **2** (1992), 57-80.
- [2] Blair, D. E., *Contact manifolds in Riemannian geometry*, Lecture Notes in Math. Springer-Verlag, Berlin-Heidelberg-New-York. **509** (1976).
- [3] Blair, D. E., Koufogiorgos, T., and Sharma, R., *A classification of 3-dimensional contact metric manifolds with  $Q\phi = \phi Q$* , Kodai Math.J. **13** (1990), 391-401.
- [4] Blair, D. E. and Sharma, R., *Three-dimensional locally symmetric contact metric manifolds*, to appear in Boll.Un.Mat.Ital..
- [5] Blair, D. E. and Vanhecke, L., *Symmetries and  $\phi$ -symmetric spaces*, Tôhoku Math.J. **39** (1987), 373-383.
- [6] Cartan, E., *Lecons sur la géométrie des espaces de Riemann*, 2nd éd., Gauthier-Villars, Paris (1946).
- [7] Cho, J. T., *On some classes of almost contact metric manifolds*, Tsukuba J. Math. **19** (1995), 201-217.
- [8] Cho, J. T., *On some classes of contact metric manifolds*, Rend.Circ.Mat. Palermo **XLIII** (1994), 141-160.
- [9] Cho, J. T., *Generalizations of locally symmetric spaces and locally  $\phi$ -symmetric spaces*, Niigata Univ. Doctorial Thesis (1994).
- [10] Olszak, Z., *On contact metric manifolds*, Tôhoku Math. J. **31** (1979).
- [11] Takahashi, T., *Sasakian  $\phi$ -symmetric spaces*, Tôhoku Math. J. **29** (1977), 91-113.
- [12] Tanaka, N., *On non-degenerate real hypersurfaces, graded Lie algebras and Cartan connections*, Japan J. Math. **2** (1976), 131-190.
- [13] Tanno, S., *Ricci curvature of contact Riemannian manifolds*, Tôhoku Math. J. **40** (1988), 441-448.
- [14] ———, *Variational problems on contact Riemannian manifolds*, Trans. Amer. Math. Soc. **314** (1989), 349-379.
- [15] Tricerri, F. and Vanhecke, L., *Homogeneous structures on Riemannian manifolds*, London Math. Soc. Lecture Note Ser. 83, Cambridge University Press, London (1983).

JONG TAEK CHO  
 TOPOLOGY AND GEOMETRY RESEARCH CENTER  
 KYUNGPOOK NATIONAL UNIVERSITY  
 TAEGU 702-701, KOREA