

Peter Vassilev Danchev

Notes on countable extensions of $p^{\omega+n}$ -projectives

Archivum Mathematicum, Vol. 44 (2008), No. 1, 37--40

Persistent URL: <http://dml.cz/dmlcz/108094>

Terms of use:

© Masaryk University, 2008

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

NOTES ON COUNTABLE EXTENSIONS
OF $p^{\omega+n}$ -PROJECTIVES

PETER DANCHEV

ABSTRACT. We prove that if G is an Abelian p -group of length not exceeding ω and H is its $p^{\omega+n}$ -projective subgroup for $n \in \mathbb{N} \cup \{0\}$ such that G/H is countable, then G is also $p^{\omega+n}$ -projective. This enlarges results of ours in (Arch. Math. (Brno), 2005, 2006 and 2007) as well as a classical result due to Wallace (J. Algebra, 1971).

Unless we do not specify some else, by the term “group” we mean “an Abelian p -group”, written additively as is the custom when dealing with such groups, for some arbitrary but a fixed prime p . All unexplained exclusively, but however used, notions and notations are standard and follow essentially those from [7]. For instance, a group is called *separable* if it does not contain elements of infinite height. As usual, for any group A , A_r denotes the reduced part of A .

A recurring theme is the relationship between the properties of a given group and its countable extension (see, e.g., [1]). The study in that aspect starts incidentally by Wallace [12] in order to establish a complete set of invariants for a concrete class of mixed Abelian groups. Specifically, his remarkable achievement states as follows.

Theorem (Wallace, 1971). *Let G be a reduced group with a totally projective subgroup H so that G/H is countable. Then G is totally projective.*

Since any reduced group is summable precisely when its socle is a free valuated vector space, as application of ([8], Lemma 7) one can derive the following.

Theorem (Fuchs, 1977). *Let G be a reduced group with a summable isotype subgroup H so that G/H is countable. Then G is summable.*

Without knowing then the cited attainment of Fuchs, we have proved in [1] an analogous assertion for summable groups of countable length via the usage of a more direct group-theoretical approach. In [5] was also showed via the construction of a concrete example that when the summable subgroup H is not isotype in G , G may not be summable.

Likewise, in [5] (see [1] too) it was obtained the following affirmation.

2000 *Mathematics Subject Classification*: Primary: 20K10; Secondary: 20K25.

Key words and phrases: abelian groups, countable factor-groups, $p^{\omega+n}$ -projective groups.

Received May 28, 2007. Editor J. Trlifaj.

Theorem (Danchev and Keef, 2005 and 2008). *Let G be a group with a subgroup H so that G/H is countable. If*

- (a) *H is σ -summable, then G is σ -summable provided that it is of limit length and H is isotype in G (when H is not isotype in G , G may not be σ -summable);*
- (b) *H is a Σ -group, then G is a Σ -group;*
- (c) *H is a Q -group, then G is a Q -group provided that it is separable;*
- (d) *H is weakly ω_1 -separable, then G is weakly ω_1 -separable provided that it is separable.*

In [1] and [3] we have shown the following statement as well.

Theorem (Danchev, 2005 and 2006). *Let G be a group with a $p^{\omega+n}$ -projective subgroup H so that G/H is countable and $n \in \mathbb{N} \cup \{0\}$. If*

- (e) *H is pure and nice in G , then G is $p^{\omega+n}$ -projective;*
- (f) *H is pure in the separable G , then G is $p^{\omega+n}$ -projective.*

Note that in [4] we have established such type results for ω -elongations of totally projective groups by $p^{\omega+n}$ -projective groups or summable groups by $p^{\omega+n}$ -projective groups, respectively.

The purpose of the present brief work is to discuss some questions as those alluded to above concerning when a given separable group is $p^{\omega+n}$ -projective provided that it has a modulo countable proper $p^{\omega+n}$ -projective subgroup, but by removing the pureness of the subgroup in the whole group.

We are now in a position to proceed by proving the next extension of point (f) (see [5] as well).

Theorem. *Suppose that G is a group of length at most ω which contains a subgroup H such that G/H is countable. Then G is $p^{\omega+n}$ -projective if and only if H is $p^{\omega+n}$ -projective, whenever $n \in \mathbb{N} \cup \{0\}$.*

Proof. The necessity is immediate because $p^{\omega+n}$ -projectives are closed with respect to subgroups (see, for example, [10]). As for the sufficiency, according to the classical Nunke's criterion for $p^{\omega+n}$ -projectivity (see [10]), there exists $P \leq H[p^n]$ with H/P a direct sum of cyclic groups. But observing that $(G/P)_r / ((G/P)_r \cap H/P) \cong ((G/P)_r + H/P) / H/P \subseteq G/P/H/P \cong G/H$ is countable with $(G/P)_r \cap H/P \subseteq H/P$ a direct sum of cyclic groups, we appeal to Wallace's theorem, quoted above, to infer that $(G/P)_r$ is totally projective. Hence G/P is simply presented. Referring now to [11] (see [7], v. II, too), we deduce that $G/P / (G/P)^1 = G/P / P_G^- / P \cong G / P_G^-$ is a direct sum of cycles, where $P_G^- = \bigcap_{i < \omega} (P + p^i G)$ is the closure of P in G . It is a straightforward argument that $p^n P_G^- \subseteq p^\omega G$. Since G is separable, that is $p^\omega G = 0$, we derive that $p^n P_G^- = 0$, so employing once again the Nunke's criterion we are finished. \square

The condition on separability may be avoided if the following strategy is realizable: Since P is bounded, one can write $P = \bigcup_{m < \omega} P_m$, where $P_m \subseteq P_{m+1} \leq P$ with $p^k P_m = 0$ for each $m < \omega$ and some $k \in \mathbb{N}$. It is readily seen that

$P_G^- = \cup_{m < \omega} K_m$, where $K_m = \cap_{i < \omega} (P_m + p^i G)$. The crucial moment is whether we may choose a nice subgroup N of G such that $N \subseteq P_m$ and such that $P_m \cap p^m G \subseteq N$ for each integer $m \geq 1$; thus P/N is strongly bounded in G/N in terms of [2]. Consequently, complying with the modular law from [7], we calculate that $K_m \cap p^m G = \cap_{m \leq i < \omega} (P_m + p^i G) \cap p^m G = \cap_{m \leq i < \omega} (P_m \cap p^m G + p^i G) \subseteq \cap_{m \leq i < \omega} (N + p^i G) = N + \cap_{m \leq i < \omega} p^i G = N + p^\omega G \leq P_G^- [p^n]$. Furthermore, we elementarily observe that $P_G^- / (N + p^\omega G) = \cup_{m < \omega} [K_m / (N + p^\omega G)]$ where, for each $m < \omega$, we compute with the aid of the modular law in [7] and the foregoing calculations that $(K_m / (N + p^\omega G)) \cap p^m (G / (N + p^\omega G)) = [K_m \cap (p^m G + N)] / (N + p^\omega G) = (N + K_m \cap p^m G) / (N + p^\omega G) \subseteq (N + p^\omega G) / (N + p^\omega G) = \{0\}$. Besides, by what we have already shown above, $G / (N + p^\omega G) / P_G^- / (N + p^\omega G) \cong G / P_G^-$ is a direct sum of cyclic groups. Knowing this, we apply the Dieudonné's criterion from [6] (see also [2]) to deduce that $G / (N + p^\omega G)$ is, in fact, a direct sum of cycles. Hence and from Nunke's criterion in [10], we conclude that G is $p^{\omega+n}$ -projective, as asserted. This completes our conclusions in all generality.

Remark. Actually, $G/P = (G/P)_r$ since $p^{\omega+n}(G/P) = 0$ by seeing that $(G/P)^1 = \cap_{i < \omega} (p^i G + P) / P \subseteq G[p^n] / P$ with $P \leq G[p^n]$ and $G^1 = 0$. However, our approach in the proof gives a more general strategy even for inseparable groups. Nevertheless, this general case is still in question.

A group A is said to be C -decomposable if $A = B \oplus K$, where B is a direct sum of cycles with $\text{fin } r(B) = \text{fin } r(A)$.

We also pose the following conjecture.

Conjecture. Suppose G is a group whose subgroup H is C -decomposable and G/H is countable. Then G is C -decomposable.

In closing, we notice that Hill jointly with Megibben have found in ([9], Proposition 1) that if G is a reduced group which possesses a torsion-complete subgroup H such that $G/H \cong \mathbf{Z}(p^\infty)$, then G is torsion-complete.

So, we are ready to state the following.

Problem. Suppose G is a group with a subgroup H which belongs to the class \mathcal{K} of Abelian p -groups. If $(G/H)[p]$ is finite, then whether or not G also belongs to \mathcal{K} ?

Investigate with a priority when \mathcal{K} coincides with the class of thick groups, torsion-complete groups, semi-complete groups, quasi-complete groups or pure-complete groups, respectively.

It is worthwhile noticing that according to the main result, stated above, the results from [4] can be improved by dropping off some unnecessary additional limitations.

REFERENCES

- [1] Danchev, P., *Countable extensions of torsion abelian groups*, Arch. Math. (Brno) **41** (3) (2005), 265–272.
- [2] Danchev, P., *Generalized Dieudonné criterion*, Acta Math. Univ. Comenian. **74** (1) (2005), 15–26.

- [3] Danchev, P., *A note on the countable extensions of separable $p^{\omega+n}$ -projective abelian p -groups*, Arch. Math. (Brno) **42** (3) (2006), 251–254.
- [4] Danchev, P., *On countable extensions of primary abelian groups*, Arch. Math. (Brno) **43** (1) (2007), 61–66.
- [5] Danchev, P., Keef, P., *Generalized Wallace theorems*, Math. Scand. (to appear).
- [6] Dieudonné, J., *Sur les p -groupes abéliens infinis*, Portugal. Math. **11** (1) (1952), 1–5.
- [7] Fuchs, L., *Infinite Abelian Groups, I, II*, Mir, Moskva, 1974 and 1977, (in Russian).
- [8] Fuchs, L., *Subfree valued vector spaces*, Lecture Notes in Math. **616** (1977), 158–167.
- [9] Hill, P., Megibben, C., *Extensions of torsion-complete groups*, Proc. Amer. Math. Soc. **44** (2) (1974), 259–262.
- [10] Nunke, R., *Topics in Abelian Groups*, ch. Purity and subfunctors of the identity, pp. 121–171, Scott, Foresman and Co., Chicago, 1963.
- [11] Nunke, R., *Homology and direct sums of countable abelian groups*, Math. Z. **101** (3) (1967), 182–212.
- [12] Wallace, K., *On mixed groups of torsion-free rank one with totally projective primary components*, J. Algebra **17** (4) (1971), 482–488.

13, GENERAL KUTUZOV STREET, BL. 7, FLOOR 2, FLAT 4
4003 PLOVDIV, BULGARIA
E-mail: pvdanchev@yahoo.com