

Jerzy Kąkol

Simple construction of spaces without the Hahn-Banach extension property

Commentationes Mathematicae Universitatis Carolinae, Vol. 33 (1992), No. 4, 623--624

Persistent URL: <http://dml.cz/dmlcz/118533>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1992

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Simple construction of spaces without the Hahn-Banach extension property

JERZY KAKOL*

Abstract. An elementary construction for an abundance of vector topologies ξ on a fixed infinite dimensional vector space E such that (E, ξ) has not the Hahn-Banach extension property but the topological dual $(E, \xi)'$ separates points of E from zero is given.

Keywords: Hahn-Banach extension property, topological vector space

Classification: 46A22

A topological vector space (tvs) $E = (E, \tau)$ over $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ is said to have the Hahn-Banach extension property (HBEP) if for every subspace F of E and every continuous linear functional f on F there is a continuous linear extension of f to the whole space E . Then the topological dual E' of E is separating, i.e. E' separates points of E from zero. Clearly every locally convex space has HBEP. In [1] Duren, Romberg, Shields gave an example of an F -space E , i.e. a metrizable and complete tvs, such that E' is separating and E has a subspace M such that E/M has a closed weakly dense subspace; hence E has not HBEP. In [5] Shapiro showed that every sequence space ℓ^p , $0 < p < 1$, also contains such subspaces. In [2] Kalton developing basic sequence techniques to F -spaces proved that all F -spaces with HBEP are locally convex. (It is still unknown if the completeness in the above result is necessary.) Hence no non-locally convex F -space has HBEP.

This note gives an elementary construction for an abundance of vector topologies ξ on a fixed infinite dimensional vector space E such that (E, ξ) has not the HBEP, but $(E, \xi)'$ separates points of E from zero. We prove the following

Theorem. *Let E be an infinite dimensional vector space and $\Delta = \{(x, x) : x \in E\}$ the diagonal of $E \times E$. For every non-zero linear functional f on Δ there are two vector topologies ϑ_1, ϑ_2 on E such that f is $\vartheta_1 \times \vartheta_2$ $|\Delta$ -continuous but f cannot be extended to a continuous linear functional on $(E, \vartheta_1) \times (E, \vartheta_2)$.*

We shall need the following simple

Lemma ([6, Example 2.5] or [4, Theorem 1.6]). *Let (E, F) be a dual pair and $\sigma(E, F)$ the weak topology on E . Let ξ be a Hausdorff nearly exotic vector topology on E , i.e. $(E, \xi)' = 0$. Then the topology $\inf\{\sigma(E, F), \xi\}$, i.e. the strongest vector topology on E weaker than ξ and $\sigma(E, F)$, is indiscrete.*

Note that every infinite dimensional vector space E admits a nearly exotic Hausdorff vector topology, cf. e.g. [4].

*This work is supported by AvH.

PROOF OF THEOREM: By Q we denote the map $x \rightarrow (x, x)$ of E onto Δ . Choose on E a normed topology ξ such that $h := f \circ Q$ is ξ -discontinuous. Set $E' = (E, \xi)'$. Let τ_1 be a Hausdorff nearly exotic topology on E . Fix $y \in E$ such that $h(y) = 2$. Let $\tau_2 := T(\tau_1)$ be the image topology on E under the map $T(x) = x - h(x)y$. Since T is injective, τ_2 is Hausdorff and nearly exotic. Set $\vartheta_i = \sup\{\sigma(E, E'), \tau_i\}$, $i = 1, 2$. By Lemma, $\inf\{\sigma(E, E'), \tau_i\}$, $i = 1, 2$, is indiscrete, so Δ is dense in $(E, \sigma(E, E') \times (E, \tau_i))$, $i = 1, 2$; clearly (E, ϑ_i) is isomorphic to $(\Delta, \sigma(E, E') \times \tau_i \upharpoonright \Delta)$, $i = 1, 2$. This and the fact that τ_i is nearly exotic, $i = 1, 2$, imply that ϑ_i and $\sigma(E, E')$ have the same continuous linear functionals. On the other hand h is continuous in $\sup\{\vartheta_1, \vartheta_2\}$ (since it is $\sup\{\tau_1, \tau_2\}$ -continuous). Hence $h \circ Q^{-1}(= f)$ is $\vartheta_1 \times \vartheta_2 \upharpoonright \Delta$ -continuous. If $h \circ Q^{-1}$ had a continuous linear extension to the whole space, then h would be ξ -continuous. This completes the proof. \square

Remark. It is known that every infinite dimensional separable normed space (E, ξ) admits a weaker metrizable nearly exotic vector topology, cf. e.g. [3, Lemma 2.1]. Let h be a ξ -discontinuous linear functional on E . Then (E, ϑ_1) (where ϑ_1 is defined as above) has HBEP. We do not know however if (E, ϑ_2) has also HBEP.

Problem. Are there tvs E, F with HBEP whose product has not HBEP?

REFERENCES

- [1] Duren P.L., Romberg R.C., Shields A.L., *Linear functionals in H^p -spaces with $0 < p < 1$* , J. Reine Angew. Math. **238** (1969), 32–60.
- [2] Kalton N.J., *Basic sequences in F -spaces and their applications*, Proc. Edinburgh Math. Soc. **19** (1974), 151–167.
- [3] Kakol J., *Nonlocally convex spaces and the Hahn-Banach extension property*, Bull. Acad. Polon. Sci. **33** (1985), 381–393.
- [4] Klee V., *Exotic topologies for linear spaces*, Proc. Symposium on General Topology and its Relations to Modern Algebra, Prague, 1961.
- [5] Shapiro J.H., *Examples of proper closed weakly dense subspaces in non-locally convex F -spaces*, Israel J. Math. **7** (1969), 369–380.
- [6] Wilansky A., *Topics in Functional Analysis*, Springer Verlag 45 (1967).

INSTITUTE OF MATHEMATICS, A. MICKIEWICZ UNIVERSITY, 60-769 POZNAŃ, POLAND

(Received April 28, 1992)