

Jan Čerych

On essential sets of function algebras in terms of their orthogonal measures

Commentationes Mathematicae Universitatis Carolinae, Vol. 36 (1995), No. 3, 471--474

Persistent URL: <http://dml.cz/dmlcz/118775>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1995

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On essential sets of function algebras in terms of their orthogonal measures

JAN ČERYCH

Abstract. In the present note, we characterize the essential set of a function algebra defined on a compact Hausdorff space X in terms of its orthogonal measures on X .

Keywords: compact Hausdorff space X , the sup-norm algebra $C(X)$ of all complex-valued continuous functions on X , its closed subalgebras (called function algebras), measure orthogonal to a function algebra

Classification: 46J10

Let X be a compact Hausdorff topological space. Denote by $C(X)$ the commutative Banach algebra, consisting of all continuous complex-valued functions on X (with respect to usual point-wise algebraic operations) endowed with the sup-norm.

By a *function algebra* on X we mean any closed subalgebra of $C(X)$ which contains constant functions on X and which separates points of X .

Definition. A function algebra A on X is said to be a *maximal* one if it is a proper subset (i.e., a proper subalgebra) of $C(X)$ and has the following property: whenever B is a function algebra on X , $B \supset A$, then either $B = A$ or $B = C(X)$.

A being a function algebra on X , the closed subset E is said to be an *essential set* of A if the following conditions are fulfilled:

- (*) A consists of all continuous prolongations of functions in the algebra of restrictions A/E (i.e., the algebra of all restrictions of functions in A from the set X to its subset E).
- (**) Whenever a closed subset F of X has the same property as E in (*), then $E \subset F$ (or, E is a unique minimal closed subset of X satisfying the condition (*)).

The notion “essential set” is due to Bear, who proved in [1] that any maximal algebra on X has an essential set.

Hoffman and Singer in [2] found an essential set of any, not necessarily maximal, function algebra on X .

Denote by $M(X)$ the space of all complex Borel regular measures on X , i.e., by the Riesz Representation Theorem, the dual space of $C(X)$.

The *annihilator* A^\perp of a function algebra A is defined to be the set of all measures $m \in M(X)$ such that $\int f dm = 0$ for any $f \in A$, or the set of all measures *orthogonal* to A . The dual space A' of A is then canonically isomorphic to the quotient space $M(X)/A^\perp$.

Now endow $M(X)$ with the weak-star topology: it is well known that $M(X)$ becomes a locally convex topological linear space with the dual space $C(X)$.

Our aim here is to characterize the essential set of a function algebra A by means of the properties of the measures in A^\perp . Remark that our construction is rather simpler than the classical one.

Theorem. *Let A be a function algebra on X . Denote by E the closure of the union of all closed supports of measures in A^\perp . Then E is the essential set of A .*

PROOF: Let $f \in C(X)$, $g \in A$ and let $f/E = g/E$, where f/E denotes the restriction of the function f from X to E . If for $m \in A^\perp$ we denote $M = \text{spt}(m)$, then

$$\int f dm = \int_M f dm = \int_M g dm = \int g dm = 0,$$

hence f is orthogonal to A^\perp and, by Banach theorem, $f \in A$. It means that E has the property (*) from Definition.

Now let a closed subset K have the property (*); we shall prove that $K \supset E$. Suppose that $K \not\supset E$. Then there is a measure $m \in A^\perp$, whose closed support is not a subset of K . Take $x \in \text{spt}(m) \setminus K$. Let V be an open neighbourhood of x in X such that its closure \bar{V} is disjoint with K . We shall find a function $f \in C(\bar{V})$ which fulfills the following two conditions:

$$\text{spt}(f) \subset V, \quad \int_V f dm \neq 0,$$

where $\text{spt}(f)$ means the closed support of f . Denote by g such a function in $C(X)$, which is equal to f on \bar{V} and equal to 0 off \bar{V} . Then $g/K = 0 \in A/K$, but

$$\int g dm = \int_V g dm = \int_V f dm \neq 0$$

and then $g \notin A^\perp$, so $g \notin A$. It follows that K has not the property (*). □

Now the following question arises: whether the word “closure” in Theorem may be omitted, or whether the essential set E of a function algebra A on X is composed of the union of closed supports of all measures in A^\perp , without closure. We shall show that it is true if X is a metric space (Proposition), but in general it is not the case (Example).

Proposition. *Let X be a compact metric space, A a function algebra on X . Then the essential set E of A is equal to the union of closed supports of all measures in A^\perp . (Especially, the union of closed supports of all orthogonal measures is a closed set.)*

PROOF: Let $x \in E$. We shall find the measure $m \in A^\perp$ such that $\text{spt}(m) \ni x$. Denote by $U_n, n = 1, 2, \dots$, the open balls in X with centres at x and radii $\frac{1}{n}$. We shall construct a finite or infinite sequence of measures $m_n \in A^\perp$ such that

- (1) $|m_n|(X) \leq 1,$
- (2) $(\text{spt}(m_n) \setminus \bigcup_{k=1}^{n-1} \text{spt}(m_k)) \cap U_n \stackrel{\text{def}}{=} M_n \neq \emptyset$ and then $|m_n|(M_n) > 0,$
- (3) $|m_n|(X) < \min_{1 \leq k \leq n-1} |m_k|(M_k),$

where $|m|$ means a total variation of a measure m .

By the Theorem, we can find a measure $m_1 \in A^\perp$ such that $|m_1|(X) = 1$ for which $\text{spt}(m_1) \cap U_1 \neq \emptyset$. If $x \in \text{spt}(m_1)$, the proof is finished. If it is not the case, then, by the Theorem, there exists the measure $m_2 \in A^\perp$ such that $(\text{spt}(m_2) \setminus \text{spt}(m_1)) \cap U_2 \neq \emptyset$; (2) follows. Multiplying m_2 by a small enough nonzero constant, we can reach fulfilling (1) and (3). If $x \in \text{spt}(m_2)$, we are done. In the opposite case, we shall continue the construction ...

In the case the sequence $\{m_n\}$ is finite, the proof is finished. If it is not the case, put

$$m = \sum_{n=1}^{\infty} \frac{1}{2^n} m_n.$$

By (1), it is $m \in M(X)$. Also $m \in A^\perp$ because $m_n \perp A$.

Take an arbitrary n . By (2), it is $|m_n|(M_n) > 0$, while $|m_k|(M_n) = 0$ for $1 \leq k \leq n - 1$. By (3), it is

$$\begin{aligned} |m|(M_n) &= \left| \sum_{k=n}^{\infty} \frac{1}{2^k} m_k(M_n) \right| \geq \frac{1}{2^n} |m_n|(M_n) - \sum_{k=n+1}^{\infty} \frac{1}{2^k} |m_k|(X) \geq \\ &\geq \frac{1}{2^n} |m_n|(M_n) - \sum_{k=n+1}^{\infty} \frac{1}{2^k} |m_k|(X) > \frac{1}{2^n} |m_n|(M_n) - \frac{1}{2^n} |m_n|(M_n) = 0 \end{aligned}$$

and then $\text{spt}(m) \cap U_n \neq \emptyset$. Since n was arbitrary, Proposition follows. □

Now, we shall construct a function algebra A on X such that there exists a point $x \in E$ which is not contained in the closed support of any measure in A^\perp .

Example. Let us denote by ω_1 the first uncountable ordinal number, put

$$\Omega = \{\omega \text{ ordinal}; \omega \leq \omega_1\},$$

let C be the closed unit disk in the complex plane. Denote by Y the cartesian product $C \times \Omega$ and let X arise from Y by “collapsing” the “last disk” $C \times \{\omega_1\}$ into one point, say x_1 , i.e., $X = Y/C \times \{\omega_1\}$. Let the algebra A consist of all functions f continuous on X such that, for a fixed ordinal ω , $\omega < \omega_1$, the function $z \mapsto f(z, \omega)$ is holomorphic in $|z| < 1$. Then the singleton $\{x_1\}$ does not meet the closed support of any measure from A^\perp , while the essential set E of A is whole X .

PROOF: (1) Any function $f \in C(\Omega)$ is constant on a neighbourhood of ω_1 .

Let us suppose that $f(\omega_1) = 0$. Put, for natural n ,

$$U_n = \{\omega \text{ ordinal}; \omega \leq \omega_1, |f(\omega)| < \frac{1}{n}\},$$

$$\omega^n = \sup \{\Omega \setminus U_n\}, \quad \omega^0 = \sup_n \omega^n.$$

It follows from the properties of ordinal numbers that $\omega^n < \omega_1$, so $\omega^0 < \omega_1$, and $f = 0$ identically on the “ordinal interval” $[\omega^0, \omega_1]$.

(2) Any function in $C(X)$ is constant on some neighbourhood of x_1 : this follows from (1).

(3) If $m \in A^\perp$ then $\text{spt}(m) \cap \{x_1\} = \emptyset$.

Let $m \in A^\perp$. Then the ordinal

$$\omega_2 = \sup\{\omega \text{ ordinal}; \omega < \omega_1, (z, \omega) \in \text{spt}(m) \text{ for some } z, z \in C\}$$

is less than ω_1 . Now let $f \in A$ be a function which is equal to 0 on the set $S = (C \times [1, \omega_2])$ and equal to 1 on $X \setminus S$. If the measure m contains a nonzero multiple of the one-point mass at $\{x_1\}$, it does not annihilate f , a contradiction. It follows that $\text{spt}(m) \subset S$.

(4) Any “non-collapsed” disk supports the measure $m \in M(X)$ for which

$$\int f dm = \int_0^1 \int_{C_r(0)} f(z) dz dr$$

where $C_r(0) = re^{it}$ for $t \in [0, 2\pi]$, $0 < r \leq 1$. But $\int f dm = 0$, by the classical Cauchy Integral Theorem, and $m \in A^\perp$. The union of such disks is $X \setminus \{x_1\}$. \square

REFERENCES

- [1] Bear H.S., *Complex function algebras*, Trans. Amer. Math. Soc. **90** (1959), 383–393.
- [2] Hoffman K., Singer I.M., *Maximal algebras of continuous functions*, Acta Math. **103** (1960), 217–241.

DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS,
CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 00 PRAHA 8, CZECH REPUBLIC

(Received December 21, 1994)