

Hans-Eberhard Porst  
Hu's Primal Algebra Theorem revisited

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 41 (2000), No. 4, 855--859

Persistent URL: <http://dml.cz/dmlcz/119217>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Hu’s Primal Algebra Theorem revisited

HANS-E. PORST

*Abstract.* It is shown how Lawvere’s one-to-one translation between Birkhoff’s description of varieties and the categorical one (see [6]) turns Hu’s theorem on varieties generated by a primal algebra (see [4], [5]) into a simple reformulation of the classical representation theorem of finite Boolean algebras as powerset algebras.

*Keywords:* Lawvere theory, equivalence between varieties, Hu’s theorem, primal algebra, Post algebras

*Classification:* Primary 06D25, 06B20; Secondary 18C05

Hu’s theorem ([4], [5]) characterizes the varieties (in Birkhoff’s sense) equivalent to the variety **Bool** of Boolean algebras as those varieties which are generated by some primal algebra. The original proof made use of Stone duality; later proofs ([3], [7]) were of a purely algebraic nature, but certainly not straightforward. Since the variety  $\mathbf{Post}_n$  of Post algebras of order  $n$  ( $n \in \mathbb{N}$ ,  $n \geq 2$ ) is generated by a primal  $n$ -element algebra (the  $n$ -chain)  $\mathbf{Post}_n$  is equivalent to **Bool** — a fact which also can be proved directly (see e.g. [1]).

In this note we are going to show how this result — plus an explicit description of all the varieties equivalent to **Bool** — can be obtained in the most simple manner using instead of Birkhoff’s description of varieties the categorical one due to Lawvere ([6]). Our analysis moreover shows that what is needed substantially to obtain these results (besides the appropriate language) is only the elementary fact that finite Boolean algebras are powersets or, more precisely, that the contravariant powerset functor is a dual equivalence between the categories of finite sets and finite Boolean algebras (we refer to this as “restricted Stone duality”).

To make this a selfcontained paper we start recalling briefly the fundamentals of Lawvere’s approach to varieties as far as they are needed here. As a first application we obtain descriptions of primality of an algebra and of the variety generated by an algebra in this language, which are obvious but seem not to have appeared in print yet.

**1 Definitions.** A (*Lawvere*) *theory* is a category  $\mathbb{T}$  with countably many objects  $T_0, T_1, T_2, \dots$  and, for each  $n \in \mathbb{N}$ , a distinguished family  $(\pi_i^n: T_n \rightarrow T_1)_{1 \leq i \leq n}$  of morphisms making  $T_n$  an  $n$ -fold power of  $T_1$ .

A functor  $\Phi: \mathbb{S} \rightarrow \mathbb{T}$  between theories is a *theory morphism* provided  $\Phi$  preserves the distinguished product families.

A  $\mathbb{T}$ -model is a product-preserving functor from  $\mathbb{T}$  into the category **Set** of sets and mappings.  $\text{Mod}\mathbb{T}$  denotes the category of all  $\mathbb{T}$ -models considered as a full subcategory of the category of all functors from  $\mathbb{T}$  to **Set**.

**2 Definitions.** Given concrete categories<sup>1</sup>  $(\mathcal{K}, \mathbf{U})$  and  $(\mathcal{L}, \mathbf{V})$ , a functor  $F: \mathcal{K} \rightarrow \mathcal{L}$  is called *concrete functor* provided  $\forall F = \mathbf{U}^2$ .

Concrete categories  $(\mathcal{K}, \mathbf{U})$  and  $(\mathcal{L}, \mathbf{V})$  are called *concretely equivalent* (or *concretely isomorphic*) provided there exists a concrete functor  $F: (\mathcal{K}, \mathbf{U}) \rightarrow (\mathcal{L}, \mathbf{V})$  which is an equivalence (an isomorphism, respectively) as a functor.

**3 Facts** ([6], [2, Chapter 3]). The category  $\text{Mod}\mathbb{T}$  is concrete by means of the underlying functor “evaluation at  $T_1$ ”, i.e., the functor  $\text{ev}_{T_1}: \text{Mod}\mathbb{T} \rightarrow \text{Set}$  mapping a model  $G$  to  $G(T_1)$  and a natural transformation  $\lambda$  to its  $T_1$ -component  $\lambda_{T_1}$ .

The fundamental observation leading to the concepts introduced above is the following. For any variety  $\mathcal{V}$  one can construct a theory  $\text{Th}\mathcal{V}$ , called the *theory of  $\mathcal{V}$* , as follows: objects of  $\text{Th}\mathcal{V}$  are the finite powers  $(F\omega)^n$ ,  $n \geq 0$ , of the free  $\mathcal{V}$ -algebra  $F\omega$  on a countable set of generators<sup>3</sup>; morphisms  $(F\omega)^m \rightarrow (F\omega)^n$  are all  $m$ -ary term operations of  $F\omega$  and morphisms  $(F\omega)^m \rightarrow (F\omega)^n$  are all maps  $t: \langle t_1, \dots, t_n \rangle$  where the  $i$ -th component  $t_i$  of  $t$  is an  $m$ -ary term operation. Note that the theory just described is nothing but the clone of  $F\omega$  extended by the maps  $\langle t_1, \dots, t_n \rangle$  and that, what often is called *clone composition*, becomes ordinary composition in this category.

Now any  $\mathcal{V}$ -algebra  $A$  determines a  $\text{Th}\mathcal{V}$ -model, denoted by  $A$ , mapping  $(F\omega)^n$  to  $A^n$  and an  $m$ -ary term (operation of  $F\omega$ ) to its interpretation on  $A$ . Then a homomorphism  $f: A \rightarrow B$  determines a natural transformation  $f: A \rightarrow B$  with  $f_{(F\omega)^n} = f^n = f \times \dots \times f$ . This construction yields a concrete equivalence between  $\mathcal{V}$  and the category of models of  $\text{Th}\mathcal{V}$ .

Conversely, for every Lawvere theory  $\mathbb{T}$  there exists a variety  $\mathcal{V}_{\mathbb{T}}$  — called the *variety determined by  $\mathbb{T}$*  — such that  $\text{Mod}\mathbb{T}$  and  $\mathcal{V}_{\mathbb{T}}$  are concretely equivalent. Moreover, the correspondence between varieties and Lawvere theories established above is essentially bijective (and functorial), i.e.,

1. every variety  $\mathcal{V}$  is concretely isomorphic to the variety  $\mathcal{V}_{\text{Th}\mathcal{V}}$  determined by  $\text{Th}\mathcal{V}$ ;
2. every Lawvere theory  $\mathbb{T}$  is isomorphic to the theory  $\text{Th}\mathcal{V}_{\mathbb{T}}$  of the variety  $\mathcal{V}_{\mathbb{T}}$  determined by  $\mathbb{T}$ ;
3. every theory morphism  $\Phi: \mathbb{S} \rightarrow \mathbb{T}$  determines a functor  $\Phi^*: \text{Mod}\mathbb{T} \rightarrow \text{Mod}\mathbb{S}$  given by  $\Phi^*(H \xrightarrow{\lambda} K) = H\Phi \xrightarrow{\lambda\Phi} K\Phi$  which is concrete;
4. every concrete functor  $F: \text{Mod}\mathbb{T} \rightarrow \text{Mod}\mathbb{S}$  is  $F = \Phi^*$  for a unique theory morphism  $\Phi: \mathbb{S} \rightarrow \mathbb{T}$ .

---

<sup>1</sup>By a *concrete category*  $(\mathcal{K}, \mathbf{U})$  is meant a category  $\mathcal{K}$  together with a faithful functor  $\mathbf{U}: \mathcal{K} \rightarrow \text{Set}$ ; a variety can always be considered as a concrete category by means of its canonical forgetful functor.

<sup>2</sup>Concrete functors between varieties are often called *interpretations*.

<sup>3</sup>We do not distinguish notationally between an algebra and its underlying set.

**4 Examples.** Each algebra  $A$  in a variety  $\mathcal{V}$  gives rise to two different theories as follows:

- a) The *theory of  $A$* , denoted by  $\mathbb{A}$ , is simply the image of  $\mathbf{A}: \mathbf{Th}\mathcal{V} \rightarrow \mathbf{Set}$ . Thus  $\mathbb{A}$  is the clone of the algebra  $A$  together with the additional maps  $A^n \xrightarrow{\langle t_i^A \rangle} A^m$  induced by families of  $(t_1^A, \dots, t_m^A)$  of  $n$ -ary derived operations. Note that for  $A = F\omega$ ,  $\mathbb{A} = \mathbf{Th}\mathcal{V}$ .
- b) The *theory generated by  $A$* , denoted by  $\mathbf{Th}_{\mathcal{V}}(A)$ , is the dual of the full subcategory of  $\mathcal{V}$  spanned by all finite copowers  $m \cdot A$ ,  $m \in \mathbb{N}$ . Explicitly:  $\mathbf{Th}_{\mathcal{V}}(A)$  has objects  $T_i = i \cdot A$  (the  $i$ -fold copower of  $A$  in  $\mathcal{V}$ ) and morphism sets  $\text{hom}_{\mathbf{Th}_{\mathcal{V}}(A)}(T_i, T_j) = \text{hom}_{\mathcal{V}}(j \cdot A, i \cdot A)$  and chosen coproduct injections  $\mu_i^n: A \rightarrow n \cdot A$  ( $i = 1, \dots, n$ ). Note that  $\mathbf{Th}_{\mathcal{V}}(F1)$  is isomorphic to  $\mathbf{Th}\mathcal{V}$ .

For each algebra  $A$  the functor  $\mathbf{A}$  factors, in the notation above, as

$$\mathbf{Th}\mathcal{V} \xrightarrow{\mathbf{A}} \mathbf{Set} = \mathbf{Th}\mathcal{V} \xrightarrow{\Phi_A} \mathbb{A} \xrightarrow{E_A} \mathbf{Set}$$

where  $E_A$  is the embedding of the subcategory  $\mathbb{A}$  into  $\mathbf{Set}$ .  $\Phi_A$  clearly is a theory-morphism which is surjective (on objects and morphisms). Note that in general  $\mathbb{A}$  is not a full subcategory of  $\mathbf{Set}$ . This being the case means that each map  $A^n \rightarrow A$  is a derived operation of the algebra  $A$ .  $A$  is then called a *primal algebra* (provided  $A$  is finite with more than one element). Thus we have shown the first of the following propositions.

**5 Proposition.** *Let  $A$  be a finite algebra in  $\mathcal{V}$  having more than one element. Then  $A$  is primal if and only if  $\mathbf{A}$  is full.*

**6 Proposition.** *For any algebra  $A$  in a variety  $\mathcal{V}$  the following hold:*

1.  $\mathbb{A}$  is the theory of  $\mathbf{HSP}(A)$ , the variety generated by  $A$ .
2.  $\mathcal{V} = \mathbf{HSP}(A)$  if and only if  $\mathbf{A}$  is faithful.

**PROOF:** Statement 1 is equivalent to the following lemma which — in the language of universal algebra — states that the variety determined by  $\mathbb{A}$  is the smallest subvariety of  $\mathcal{V}$  containing  $A^4$ . The second statement is an immediate consequence of the first one:  $\Phi_A: \mathbf{Th}\mathcal{V} \rightarrow \mathbb{A}$  is an isomorphism iff  $\mathbf{A}$  is a faithful model; thus, in this case  $\mathcal{V}$  and  $\mathbf{HSP}(A)$  coincide by 1. and bijectivity of the correspondence between theories and varieties. □

**7 Lemma.** *Let  $\mathbb{S}$  be a theory and  $\mathbf{E}: \mathbf{Mod}\mathbb{S} \rightarrow \mathbf{Mod}\mathbb{T}$  be a full concrete embedding. Assume that the  $\mathbb{T}$ -model  $\mathbf{A}$  is  $\mathbf{E}(A')$  for some  $\mathbb{S}$ -model  $A'$ . Then there exists a full concrete embedding  $\mathbf{Mod}\mathbb{A} \hookrightarrow \mathbf{Mod}\mathbb{S}$ .*

**PROOF:** Let  $\mathbf{E}$  be  $\Phi^*$  for  $\Phi: \mathbb{T} \rightarrow \mathbb{S}$ . Then  $\mathbf{A} = A' \circ \Phi$  and therefore  $\Phi_A = \Phi_{A'} \cdot \Phi: \mathbb{T} \rightarrow \mathbb{A}$ . Then  $\Phi_{A'}$  is surjective since  $\Phi_A$  is. Now apply the following lemma, the proof of which is an easy exercise. □

---

<sup>4</sup>A somewhat informal way of proving this is the following: an algebra  $B$  is in  $\mathbf{HSP}(A)$  iff  $B$  satisfies all equations which are valid in  $A$ . The  $\mathbb{T}$ -model  $\mathbf{B}$  “is” an  $\mathbb{A}$ -model iff  $\mathbf{B}$  factors over  $\Phi_A$  iff the algebra  $B$  satisfies all equations of  $A$ .

**8 Lemma.** *The functor  $\Phi^*: \text{Mod}\mathbb{T} \rightarrow \text{Mod}\mathbb{S}$  is a full embedding for every surjective theory morphism  $\Phi: \mathbb{S} \rightarrow \mathbb{T}$ .*

It is well known that varieties might be equivalent as categories without being concretely equivalent (in the latter case they would in fact be concretely isomorphic) as, e.g., the varieties of left modules over Morita-equivalent rings or the varieties of Post algebras of the various fixed (finite) orders. In order to describe the varieties  $\mathcal{W}$  equivalent to a given variety  $\mathcal{V}$  the following notion is crucial:

**9 Definition.** An algebra  $G$  in a variety  $\mathcal{V}$  is called a *variatal generator*<sup>5</sup>, provided  $G$  is a retract of some finitely generated free  $\mathcal{V}$ -algebra  $F_n$  and the free algebra  $F_1$  on one generator is a retract of some finite copower of  $G$ .

**10 Fact** ([6], [8]). Let  $\mathcal{V}$  be a variety with theory  $\mathbb{T}$ . A variety  $\mathcal{W}$  with theory  $\mathbb{S}$  is equivalent to  $\mathcal{V}$  iff  $\mathbb{S}$  is (isomorphic to) some theory  $\text{Th}_{\mathcal{V}}(G)$  where  $G$  is a variatal generator in  $\mathcal{V}$ .

**11 Example.** Let  $\mathcal{V}$  be the variety **Bool** of Boolean algebras. The variatal generators of **Bool** are (up to isomorphism) precisely the powerset algebras  $\mathbf{2}^n$ ,  $n \in \mathbb{N}$ ,  $n \geq 2$ . This is clear since each retract of a finitely generated free Boolean algebra (which is finite) is finite, thus  $\mathbf{2}^n$  for some  $n \in \mathbb{N}$  due to restricted Stone duality. Moreover  $\mathbf{2}^2$  is a retract of  $\mathbf{2}^n$  iff  $n \geq 2$  (again by restricted Stone duality).

It is easy to describe the theory generated by  $\mathbf{2}^n$ : the contravariant powerset functor, restricted to the full subcategory  $\mathbb{T}_n$  of **Set** spanned by the finite powers of the  $n$ -element set  $\{0, 1, \dots, n-1\}$  provides a theory-isomorphism  $\mathbb{T}_n \simeq \text{Th}_{\text{Bool}}(\mathbf{2}^n)$ .

As a consequence of the above a proof of the following sharpening of Hu's theorem ([4] [5]) becomes nearly trivial, making thereby clear in addition that this theorem is essentially — up to the well known categorical result stated as Fact 10 above — a reformulation of *restricted* Stone duality.

**12 Theorem.** *The following are equivalent for a variety  $\mathcal{V}$ :*

- (i)  $\mathcal{V}$  is equivalent to **Bool**, the variety of Boolean algebras;
- (ii)  $\mathcal{V}$  is generated by a primal algebra;
- (iii)  $\text{Th}\mathcal{V} \simeq \mathbb{T}_n$  for some  $n \in \mathbb{N}$ ,  $n \geq 2$ .

**PROOF:** By Fact 10 and Example 11  $\mathcal{V}$  is equivalent to **Bool** iff  $\text{Th}\mathcal{V} \simeq \mathbb{T}_n$  for some  $n \geq 2$ .

By Propositions 5 and 6 one gets:  $\mathcal{V} = \text{HSP}(A)$  with an  $n$ -element primal algebra  $A$  iff  $A$  is a full and faithful  $\text{Th}\mathcal{V}$ -model, iff  $\text{Th}\mathcal{V} \simeq \mathbb{T}_n$ . (Note that  $\mathbb{T}_n$  has a full and faithful  $n$ -element model trivially: its embedding into **Set**).  $\square$

---

<sup>5</sup>In categorical terms, this is equivalent to saying that  $G$  is an extremally projective, finitely presentable extremal generator (see e.g. [8]).

**13 Remark.** Since the correspondence between varieties and Lawvere theories is essentially bijective the above theorem shows in particular that, for each  $n \in \mathbb{N}$ ,  $n \geq 2$ , there is (up to concrete isomorphism) precisely one variety which is generated by an  $n$ -element primal algebra, namely the variety  $\mathcal{V}_{\mathbb{T}_n}$  determined by the theory  $\mathbb{T}_n$ , and that any variety equivalent to **Bool** has to be one of those (again up to concrete isomorphism). Now the variety  $\mathcal{V}_{\mathbb{T}_n}$  is easily identified as the variety **Post** $_n$  of Post algebras of order  $n$  since it is well known (see e.g. [1]) that the latter is equivalent to **Bool** and generated by an  $n$ -element primal algebra. For a more categorical argument see [8].

## REFERENCES

- [1] Balbes R., Dwinger Ph., *Distributive Lattices*, University of Missouri Press, Missouri, 1974.
- [2] Borceux F., *Handbook of Categorical Algebra Vol. 2*, Cambridge University Press, Cambridge, 1994.
- [3] Davey B.A., Werner H., *Dualities and equivalences for varieties of algebras*, in A.P. Huhn and E.T. Schmidt, editors, 'Contributions to Lattice Theory' (Proc. Conf. Szeged 1980), vol. 33 of Coll. Math. Soc. János Bolyai, North-Holland, 1983, pp. 101–275.
- [4] Hu T.K., *Stone duality for Primal Algebra Theory*, Math. Z. **110** (1969), 180–198.
- [5] Hu T.K., *On the topological duality for Primal Algebra Theory*, Algebra Universalis **1** (1971), 152–154.
- [6] Lawvere F.W., *Functorial semantics of algebraic theories*, PhD thesis, Columbia University, 1963.
- [7] McKenzie R., *An algebraic version of categorical equivalence for varieties and more general algebraic theories*, in A. Ursini and P. Agliano, editors, 'Logic and Algebra', vol. 180 of Lecture Notes in Pure and Appl. Mathematics, Marcel Dekker, 1996, pp. 211–243.
- [8] Porst H.-E., *Equivalence for varieties in general and for Bool in particular*, to appear in Algebra Universalis.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BREMEN, 28359 BREMEN, GERMANY

E-mail: porst@math.uni-bremen.de

(Received February 14, 2000)