

Roger Yue Chi Ming

On p -injectivity, YJ-injectivity and quasi-Frobeniusean rings

Commentationes Mathematicae Universitatis Carolinae, Vol. 43 (2002), No. 1, 33--42

Persistent URL: <http://dml.cz/dmlcz/119298>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On p -injectivity, YJ-injectivity and quasi-Frobeniusean rings

ROGER YUE CHI MING

Dedicated to Professor Carl Faith on his 75-th birthday

Abstract. A new characteristic property of von Neumann regular rings is proposed in terms of annihilators of elements. An ELT fully idempotent ring is a regular ring whose simple left (or right) modules are either injective or projective. Artinian rings are characterized in terms of Noetherian rings. Strongly regular rings and rings whose two-sided ideals are generated by central idempotents are characterized in terms of special annihilators. Quasi-Frobeniusean rings are characterized in terms of p -injectivity. Also, a commutative YJ-injective ring with maximum condition on annihilators and finitely generated socle is quasi-Frobeniusean.

Keywords: von Neumann regular, V -ring, Artinian ring, p -injectivity, YJ-injectivity, quasi-Frobeniusean

Classification: 16D30, 16D36, 16D50

Introduction

Von Neumann regular rings, V -rings, self-injective rings and generalizations are extensively studied since several years (cf. for example, [1], [3]–[16], [28]–[30]). This sequel of [21] and [24] contains the following results for a ring A : (1) A is von Neumann regular iff A is a semi-prime ring such that every finitely generated one-sided ideal is the annihilator of an element of A (Theorem 1); (2) A is strongly regular iff for every $b \in A$, $Ab + I(AbA)$ is a complement right ideal of A (Proposition 3); (3) Left Noetherian rings whose essential right ideals are idempotent two-sided ideals are left Artinian (Proposition 4); (4) Special two-sided ideals are used to characterize rings whose two-sided ideals are generated by central idempotents (Proposition 8), (5) A is quasi-Frobeniusean iff A has a p -injective left generator and projective p -injective left A -modules are injective (Theorem 9); (6) If every simple right A -module is flat and every maximal left ideal of A is either injective or a two-sided ideal of A , then A is either a left self-injective regular left V -ring or strongly regular; (7) If A is commutative, then A is quasi-Frobeniusean iff A is a YJ-injective ring with maximum condition on annihilators and a finitely generated socle (Theorem 11).

Throughout, A denotes an associative ring with identity and A -modules are unital. Recall that a left A -module M is p -injective (resp f -injective) if, for any

principal (resp. finitely generated) left ideal I of A , every left A -homomorphism of I into M extends to A (cf. [4, p. 122], [14, p. 340], [17]). A is called left p -injective (resp. f -injective) if ${}_A A$ is p -injective (resp. f -injective). Similarly, p -injectivity and f -injectivity are defined on the right side. (P -injectivity is also called principal injectivity in the literature). Following C. Faith, A is called a left V -ring if every simple left A -module is injective. A well-known theorem of I. Kaplansky asserts that a commutative ring is von Neumann regular iff it is a V -ring. In general, von Neumann regular rings need not be V -rings and the converse is not true either. A theorem of M. Ikeda-T. Nakayama asserts that A is a left p -injective ring iff every principal right ideal of A is a right annihilator. It is well-known that A is von Neumann regular iff every left (right) A -module is flat ([4, p. 91]). This remains true if “flat” is replaced by “ p -injective” ([17]).

Our first result is motivated by [19, Question 1] and [27, Question 1].

Theorem 1. *The following conditions are equivalent:*

- (1) A is von Neumann regular;
- (2) A is a semi-prime ring whose finitely generated one-sided ideals are annihilators of an element of A ;
- (3) A is a semi-prime ring such that every finitely generated left ideal is the left annihilator of an element of A and every principal right ideal of A is the right annihilator of an element of A .

PROOF: It is clear that (1) implies (2) while (2) implies (3). Assume (3). Let F be a finitely generated left ideal of A . Then $F = I(u)$, $u \in A$. Since $uA = r(w)$, $w \in A$, $F = I(uA) = I(r(w)) = I(r(Aw))$. But Aw is a left annihilator which implies that $Aw = I(r(Aw))$. Therefore $F = Aw$ which shows that every finitely generated left ideal of A is principal. Since A is semi-prime, A is left semi-hereditary by [4, Theorem 7.5B] and it follows that every principal left ideal of A is projective. Since every principal right ideal is a right annihilator, by Ikeda-Nakayama’s theorem, A is left p -injective. Now A being a left $p.p.$ ring is equivalent to “every quotient of a p -injective left A -module is p -injective” ([18, Remark 2]). Since A is left p -injective, every cyclic left A -module is p -injective which yields that A is von Neumann regular ([17, Lemma 2]). Thus (3) implies (1). \square

In Theorem 1, the term “semi-prime” cannot be omitted (otherwise, any principal ideal quasi-Frobeniusean ring would be von Neumann regular!).

Remark 1. If every finitely generated one-sided ideal of A is the annihilator of an element, then A is a left and right IF-ring whose finitely generated one-sided ideals are principal (cf. [4, Theorem 6.9]).

Remark 2. The fact that a strongly regular ring is unit-regular follows from a cancellation theorem of G. Ehrlich (cf. [4, Corollary 6.3C] and [5, Corollary 4.2]).

The proof of [17, Proposition 1] shows in an elementary way that this result holds. Note that a left and right V -ring whose essential left ideals are two-sided is a unit-regular ring (whose prime factor rings are Artinian).

We may note the following characterization of principal ideal rings in terms of p -injectivity.

Remark 3. A is a principal left ideal ring iff every finitely generated left ideal of A is principal and every p -injective left A -module is injective.

Another result on annihilators.

Proposition 2. *If every finitely generated left ideal of A is the left annihilator of a finite subset of A and every finitely generated right ideal of A is a right annihilator, then A is left f -injective and right p -injective.*

PROOF: Since every principal one-sided ideal of A is an annihilator, A is a left and right p -injective ring by Ikeda-Nakayama's theorem. Let F, K be finitely generated left ideals of A . By hypothesis, we have $F = I(U)$, $K = I(V)$, where U, V are finitely generated right ideals of A . Then $U = r(I(U))$, $V = r(I(V))$ which imply $r(F) + r(K) = r(I(U)) + r(I(V)) = U + V = r(I(U + V)) = r(I(U) \cap I(V)) = r(F \cap K)$. By Ikeda-Nakayama's theorem, A is left f -injective. \square

Question 1. If A is left p -injective such that every finitely generated left ideal of A is the left annihilator of an element of A , is A left f -injective?

Remark 4. In Proposition 2, $\text{Soc}({}_A A) = \text{Soc}(A_A)$.

Proposition 3. *The following conditions are equivalent:*

- (1) A is strongly regular;
- (2) for any $b \in A$, $Ab + I(AbA)$ is a complement right ideal of A .

PROOF: Assume (1). For any $b \in A$, $Ab = Ae$, where e is a central idempotent. Then $I(AbA) = I(b) = A(1-e)$ and $Ab + I(AbA) = Ae \oplus A(1-e) = A$. Therefore (1) implies (2) evidently.

Assume (2). Suppose that $c \in A$ such that $(Ac)^2 = 0$. Then $I(AcA)$ is an essential right ideal of A . By hypothesis, $I(AcA) = Ac + I(AcA)$ is a complement right ideal of A which proves that $A = I(AcA)$, whence $c = 0$. This proves that A must be semi-prime. It follows that for any $b \in A$, $I = Ab \oplus I(AbA)$ (because $Ab \cap I(Ab) = 0$) is a complement right ideal of A . Now there exists a complement right ideal C of A such that $I \oplus C$ is an essential right ideal of A . Then $CAb \subseteq C \cap Ab \subseteq C \cap I = 0$ implies that $C \subseteq I(AbA) \subseteq I$ and hence $C \subseteq C \cap I = 0$. Therefore I is an essential right ideal of A which yields $I = A$. Therefore $A = Ab \oplus I(AbA)$ which proves that A is von Neumann regular. Since $AbA \cap I(AbA) = 0$ (in as much as A is semi-prime), then $A = Ab \oplus I(AbA) \subseteq AbA \oplus I(AbA)$ which yields $A = AbA \oplus I(AbA)$. For any $u \in AbA$, $u = v + w$,

$v \in Ab$, $w \in I(AbA)$. Then $u - v \in AbA \cap I(AbA) = 0$ which implies $u = v \in Ab$, proving that $Ab = AbA$ is generated by a central idempotent (because A is semi-prime). Thus (2) implies (1). \square

Question 2. If $Ab + r(AbA)$ is a complement left ideal of A for every $b \in A$, is A regular, biregular?

We now give a sufficient condition for left Noetherian rings to be left Artinian.

Proposition 4. *If A is a left Noetherian ring such that every essential right ideal of A is an idempotent two-sided ideal, then A is left Artinian.*

PROOF: Let B be a prime factor ring of A . Then every essential right ideal of B is an idempotent two-sided ideal of B . For any $0 \neq b \in B$, set $T = BbB$. Let K be a complement right subideal of T such that $E = bB \oplus K$ is an essential right subideal of T . Since T is an essential right ideal of B (B being prime), E is an essential right ideal of B which implies that E is an idempotent two-sided ideal of B . Now $b \in E = E^2$ implies that $b = \sum_{i=1}^n (bb_i + k_i)(bc_i + s_i)$, $b_i, c_i \in B$ and $k_i, s_i \in K$, whence $b - \sum_{i=1}^n bb_i(bc_i + s_i) = \sum_{i=1}^n k_i(bc_i + s_i) \in bB \cap K = 0$. Then $b \in BbB$ which proves that every right ideal of B is idempotent. Since every essential right ideal of B is two-sided, then B is von Neumann regular by [1, Theorem 3.1]. Since B is left Noetherian, it is well-known that B must be simple Artinian. If A is prime, then A is Artinian as just seen. If A is not prime, then by [3, Lemma 18.34B], A is left Artinian. This establishes the proposition. \square

Note that the ring in Proposition 4 needs not be right duo. The proof of Proposition 4 yields a characterization of Artinian rings.

Theorem 5. *The following conditions are equivalent:*

- (1) A is left Artinian;
- (2) A is a left Noetherian ring such that for any prime factor ring B of A , every essential right ideal of B is an idempotent two-sided ideal of B .

An ideal of A will always mean a two-sided ideal of A . Recall that A is ELT (resp. ERT) if every essential left (resp. right) ideal of A is an ideal of A . As usual, A is called fully (resp. (1) fully left; (2) fully right) idempotent if every ideal (resp. (1) left ideal; (2) right ideal) of A is idempotent.

Note that if A is fully idempotent and every maximal left (with even every maximal right) ideal of A is an ideal, then A needs not be von Neumann regular ([28, Theorem 1]).

Theorem 6. *If A is an ELT fully idempotent ring, then A is a von Neumann regular ring whose simple right (or left) modules are either projective or injective.*

PROOF: Let B be a prime factor ring of A . Then B is an ELT fully idempotent ring. The proof of Proposition 4 shows that B is fully left idempotent. By

[1, Theorem 3.1], B is von Neumann regular. Looking carefully at the proof of [1, Theorem 3.1], we see that A is also ERT. Let M be a maximal right ideal of A . If A/M_A is not projective, then M_A is essential in A_A which implies that M is an ideal of A and is therefore a maximal left ideal of A . For any $y \in M$, $y \in yAyu \subseteq yM$ which implies that the simple left A -module A/M is flat ([2, p. 458]). By [21, Lemma 1], A/M_A is injective. This proves that every simple right A -module is either injective or projective. Similarly, every simple left A -module is either injective or projective. \square

Corollary 7. *An ELT fully idempotent ring is either regular with non-zero socle or strongly regular.*

The next remark is connected with [6, Corollary 6], [18, Lemma 1] and [26, Question 1].

Remark 5. If A is an ERT (or ELT) ring whose simple left modules are p -injective, then A is regular and every simple one-sided A -module is either injective or projective.

Remark 6. If A is a semi-prime ELT ring containing an injective maximal left ideal, then A is a left and right self-injective, left and right V -ring of bounded index. Consequently, A is left and right FPF by a theorem of S. Page [4, Theorem 5.49].

We now consider a particular class of biregular rings which generalizes simple non-Artinian rings and semi-simple Artinian rings.

We introduce two definitions.

Definitions. Let E, T be ideals of A , $E \subseteq T$. We say that

- (1) T is an essential extension of E (or E is essential in T) if $E \cap N \neq 0$ for any non-zero ideal N of A contained in T ;
- (2) E is a complement ideal of A if E has no proper essential extension in A .

Proposition 8. *The following conditions are equivalent for a ring A :*

- (1) *every ideal of A is generated by a central idempotent;*
- (2) *for every ideal T of A , $T + (I(T) \cap r(T))$ is a complement ideal of A .*

PROOF: Assume (1). Let T be an ideal of A . If $I = T + (I(T) \cap r(T))$, since $T = Ae$, where e is a central idempotent, then $I(T) = I(eA) = A(1 - e) = (1 - e)A = r(T)$ and $A = Ae \oplus A(1 - e) = T \oplus (I(T) \cap r(T))$. Therefore (1) implies (2).

Assume (2). Let T be an ideal of A such that $T^2 = 0$. Then $r(T)$ is an essential left ideal of A which implies $r(T)$ is an essential ideal of A . Similarly, $I(T)$ is an essential ideal of A . Therefore $r(T) \cap I(T)$ is an essential ideal of A which implies that $T + I(T) \cap r(T)$ is an essential ideal of A . By hypothesis, $T + (I(T) \cap r(T))$ is a complement ideal of A which yields $T + (I(T) \cap r(T)) = A$.

But $T^2 = 0$ implies that $T \subseteq I(T) \cap r(T)$, whence $A = I(T) \cap r(T)$, yielding $A = I(T) = r(T)$. This implies $T = 0$ and proves that A must be semi-prime. Now for any ideal U of A , $U \cap I(U) = 0$ and $I(U) = r(U)$. If $I = U + (I(U) \cap r(U))$, then $I = U + I(U) = U \oplus I(U)$. Suppose that I is not an essential ideal of A : there exists a non-zero ideal N of A such that $I \cap N = 0$. Now $NU \subseteq N \cap U \subseteq N \cap I = 0$ which implies that $N \subseteq I(U) = r(U)$, whence $N = N \cap I(U) \subseteq N \cap I = 0$. This contradiction proves that I must be essential in A . By hypothesis, $I = A$ which proves that U is generated by a central idempotent (in as much as A is semi-prime). Thus (2) implies (1). \square

The following property of p -injectivity seems interesting.

Remark 7. If T is an ideal of A such that ${}_A A/T$ is p -injective, then the factor ring A/T is left p -injective.

Quasi-Frobeniusean rings are now characterized in terms of p -injectivity.

Theorem 9. *The following conditions are equivalent:*

- (1) A is quasi-Frobeniusean;
- (2) A is left pseudo-Frobeniusean and projective p -injective right A -modules are injective;
- (3) there exists a p -injective left generator of $A\text{-Mod}$ and projective p -injective left A -modules are injective.

PROOF: (1) implies (2) and (3) by [3, Theorem 24.20].

Assume (2). Since A is left pseudo-Frobeniusean, then every left ideal of A is a left annihilator which implies that A is right p -injective. For any projective right A -module P , there exist B , a direct sum of copies of ${}_A A$, and an epimorphism p of B onto P_A . Then $B/\ker p \approx P_A$ implies that $B \approx \ker p \oplus B/\ker p$. Since B is a direct sum of p -injective right A -modules, then B_A is p -injective. Therefore $B/\ker p$ is a p -injective right A -module and hence P_A is p -injective. By hypothesis, P_A is injective. Then (2) implies (1) by [3, Theorem 24.20].

Assume (3). Let G be a p -injective left generator of $A\text{-Mod}$. For any projective left A -module F , there exists C , a direct sum of copies of ${}_A G$, and an epimorphism $q : {}_A C \rightarrow {}_A F$. As before, we obtain a p -injective left A -module F which is injective by hypothesis. Thus (3) implies (1) by [3, Theorem 24.20]. \square

Recall that a left A -module M is YJ-injective if, for any $o \neq a \in A$, there exists a positive integer n with $a^n \neq 0$ such that every left A -homomorphism of Aa^n into M extends to A (cf. [15], [24], [30]). A is called left YJ-injective if ${}_A A$ is YJ-injective. Similarly, YJ-injectivity is defined on the right side. In [15], it is shown that YJ-injectivity generalizes p -injectivity even for rings (quasi-injectivity generalizes injectivity but the two concepts coincide for rings). Also left YJ-injective rings generalize right IF-rings. If A is left YJ-injective, then every

non-zero-divisor is invertible in A . Consequently, A coincides with $Q_{cl}^l(A)$ and $Q_{cl}^r(A)$, the classical left and right quotient rings of A . It is well-known that A is left non-singular iff A has a maximal left quotient ring, noted $Q_{\max}^l(A)$, which is a left self-injective regular ring. If A is a reduced ring, $Q_{\max}^l(A)$, is not necessarily strongly regular. However, if A is reduced and admits a classical left quotient ring $Q_{cl}^l(A)$, then by [22, Proposition 1.5], $Q_{cl}^l(A)$ is a reduced ring (this is the case when A is left non-singular, left duo). In that case, if $Q_{cl}^l(A)$ is left or right YJ-injective, then it is strongly regular. Consequently, a reduced left YJ-injective ring is strongly regular.

A well-known theorem of Y. Utumi asserts that if A is left and right non-singular, then $Q_{\max}^l(A)$ and $Q_{\max}^r(A)$ coincide iff every complement one-sided ideal of A is an annihilator. (The terms “annihilator” and “complement” should be permuted in [4, p. 181]).

Left (or right) Johns rings are studied in [4].

Question 3. Is a left Johns, left YJ-injective ring quasi-Frobeniusean?

(We know that a left p -injective left Johns ring is quasi-Frobeniusean).

Now let J , Y , Z denote respectively the Jacobson radical, the right singular ideal and the left singular ideal of the ring A .

Proposition 10. *Let A be a ring whose simple right modules are flat. If every maximal left ideal of A is either injective or an ideal of A , then either A is a left self-injective regular left V -ring or A is strongly regular. Consequently, A must be a regular left V -ring.*

PROOF: First suppose that every maximal left ideal of A is an ideal of A . For any maximal left ideal N of A , N is a maximal right ideal of A and by hypothesis, A/N_A is flat. Then ${}_A A/N$ is injective by [21, Lemma 1] which implies that A is a left V -ring, whence A is strongly regular (cf [17, Proposition 3]). Now suppose that there exists a maximal left ideal M of A which is not an ideal of A . Then ${}_A M$ is injective which implies $A = M \oplus U$, where U is a minimal projective left ideal of A . Let V be an arbitrary minimal projective left ideal of A . Write $V = Av$, $0 \neq v \in A$. If $MV = 0$, then $MAv = 0$ which implies that $MA = M$ (because $MA \neq A$). This contradicts the hypothesis that M is not an ideal of A . Therefore $MV \neq 0$ and $Mw \neq 0$ for some $0 \neq w \in V$. Now $Mw = V$ and we have an epimorphism $p : M \rightarrow V$ defined by $p(m) = mw$ for all $m \in M$. Then $M/\ker p \approx V$ which yields $M \approx \ker p \oplus M/\ker p$ (in as much as ${}_AV$ is projective). Since ${}_A M$ is injective, then so is $M/\ker p$, proving that ${}_AV$ is injective. In particular, ${}_AU$ is injective which implies that A is left self-injective. Now for any maximal left ideal L of A , if ${}_A L$ is injective, then ${}_A A/L$ is injective as just seen. If L is an ideal of A , then A/L_A is flat which implies that ${}_A A/L$ is injective ([21, Lemma 1]). In any case, ${}_A A/L$ is injective, proving that A is a left V -ring, whence $J = 0$ (cf. [18, Lemma 1]). Since A is left self-injective, $Z = J = 0$

and hence A is von Neumann regular. We conclude that A must be a regular, left V -ring. \square

We now turn to a characterization of commutative quasi-Frobeniusean rings.

Theorem 11. *The following conditions are equivalent for a commutative ring A :*

- (1) A is quasi-Frobeniusean;
- (2) A is a YJ-injective ring with maximum condition on annihilators and $\text{Soc}(A)$, the socle of A , is finitely generated.

PROOF: It is clear that (1) implies (2).

Assume (2). Since A is a commutative YJ-injective, then A coincides with its classical quotient ring. Since A satisfies the maximum condition on annihilators, then A/J is Artinian ([4, Theorem 16.31]) and also Y is nilpotent. Now A being YJ-injective implies that $J = Y$ ([24, p. 103]), whence A is semi-primary. Therefore A has an essential socle. Given $\text{Soc}(A)$ finitely generated, we then have a finitely embedded ring A satisfying the maximum condition on annihilators which yields A Artinian ([4, p. 164]). Since A is YJ-injective, every minimal ideal of A must be an annihilator. In that case, A is quasi-Frobeniusean by a theorem of H.H. Storrer. Thus (2) implies (1). \square

Remark 8. A right YJ-injective ring whose simple right modules are either p -injective or projective is fully right idempotent (this is because $Y = J = Y \cap J = 0$).

Theorem 11 motivates the next question on YJ-injectivity.

Question 4. Is a commutative YJ-injective ring with maximum condition on annihilators quasi-Frobeniusean?

We add a remark on flatness and p -injectivity.

Remark 9. We know that A is von Neumann regular if every cyclic singular left A -module is flat (Math. J. Okayama Univ. 20 (1978), 123–129 (Theorem 5)). If every singular left A -module is injective, A needs not be von Neumann regular ([4, p. 92]). Consequently, this is also the case when every cyclic singular left A -module is p -injective. However, A is von Neumann regular if A is also left p -injective. We may also recall the following: If I is a p -injective left ideal of A , then $_A A/I$ is flat.

In 1974, we introduced the concept of p -injective modules to study von Neumann regular rings, V -rings and associated rings ([17]). In 1985, this is generalized to YJ-injective modules ([24]). In 1998, Xue Weimin showed that even for rings, YJ-injectivity effectively generalizes p -injectivity [15]. Finally, Zhang-Wu [30] proved that if every left A -module is YJ-injective, then A is von Neumann regular (which answers a question raised in [24]).

K.R. Goodearl's volume on von Neumann regular rings [5] has motivated numerous papers in that area during the last twenty years. It is now a classic for people interested in VNR rings (cf. [4]).

In view of [17, Proposition 1] and our Theorem 1 here, we raise the last question.

Question 5. Is A von Neumann regular if A is a semi-prime ring such that every principal one-sided ideal is the annihilator of an element of A ?

Note that semi-prime rings whose principal one-sided ideals are annihilators need not be von Neumann regular (cf. K. Beidar–R. Wisbauer, Comm. Algebra 23 (1995), 841–861 (Example 4.8), which answered in the negative a question raised in 1981).

In recent years, Xue Weimin and Zhang Jule solved several problems raised in my papers. Among the still unanswered questions, we recall the following:

U.Q.1. Is A von Neumann regular if A satisfies any one of the following conditions: (a) A is left semi-hereditary and every maximal left ideal of A is p -injective; (b) A is a left p -injective left V -ring; (c) every principal left ideal of A is a projective left annihilator; (d) A is left semi-hereditary and every simple left A -module is flat; (e) A is a semi-prime left self-injective ring whose maximal essential left ideals are two-sided.

U.Q.2. Is A strongly regular if A is a reduced ring whose principal left ideals are complement left ideals?

U.Q.3. Is A fully left idempotent if every simple left A -module is YJ-injective? (The answer is “yes” if “YJ-injective” is replaced by “ p -injective”).

U.Q.4. Is A Artinian if A is a prime left self-injective ring whose maximal essential left ideals are two-sided?

U.Q.5. Is A left pseudo-Frobeniusean if A is a left Kasch ring containing an injective maximal left ideal?

REFERENCES

- [1] Baccella G., *Generalized V-rings and von Neumann regular rings*, Rend. Sem. Mat. Univ. Padova **72** (1984), 117–133.
- [2] Chase S.U., *Direct product of modules*, Trans. Amer. Math. Soc. **97** (1960), 457–473.
- [3] Faith C., *Algebra II: Ring Theory*, Grundlehren Math. Wiss. **191** (1976).
- [4] Faith C., *Rings and things and a fine array of twentieth century associative algebra*, AMS Math. Survey and Monographs **65** (1999).
- [5] Goodearl K.R., *Von Neumann Regular Rings*, Pitman, 1979.
- [6] Hirano Y., Tominaga H., *Regular rings, V-rings and their generalizations*, Hiroshima Math. J. **9** (1979), 137–149.
- [7] Hirano Y., *On non-singular p -injective rings*, Publ. Math. **38** (1994), 455–461.
- [8] Huynh D.V., Dung N.V., *A characterization of Artinian rings*, Glasgow Math. J. **30** (1988), 67–73.

- [9] Huynh D.V., Dung N.V., *Rings with restricted injective condition*, Arch. Math. **54** (1990), 539–548.
- [10] Huynh D.V., Dung N.V., Smith P.F., Wisbauer R., *Extending Modules*, Pitman, London, 1994.
- [11] Mohamed S.H., Mueller B.J., *Continuous and discrete modules*, LMS Lecture Note 47 (C.U.P.), 1990.
- [12] Puninski G., Wisbauer R., Yousif M.F., *On p -injective rings*, Glasgow Math. J. **37** (1995), 373–378.
- [13] Wang Ding Guo, *Rings characterized by injectivity classes*, Comm. Algebra **24** (1996), 717–726.
- [14] Wisbauer R., *Foundations of Module and Ring Theory*, Gordon and Breach, New-York, 1991.
- [15] Xue Wei Min, *A note on YJ-injectivity*, Riv. Mat. Univ. Parma (6) **1** (1998), 31–37.
- [16] Xue Wei Min, *Rings related to quasi-Frobenius rings*, Algebra Colloq. **5** (1998), 471–480.
- [17] Yue Chi Ming R., *On von Neumann regular rings*, Proc. Edinburgh Math. Soc. **19** (1974), 89–91.
- [18] Yue Chi Ming R., *On simple p -injective modules*, Math. Japonica **19** (1974), 173–176.
- [19] Yue Chi Ming R., *On regular rings and continuous rings*, Math. Japonica **24** (1979), 563–571.
- [20] Yue Chi Ming R., *On V-rings and prime rings*, J. Algebra **62** (1980), 13–20.
- [21] Yue Chi Ming R., *On regular rings and Artinian rings*, Riv. Mat. Univ. Parma (4) **8** (1982), 443–452.
- [22] Yue Chi Ming R., *On von Neumann regular rings, X*, Collectanea Math. **34** (1983), 81–94.
- [23] Yue Chi Ming R., *On regular rings and continuous rings, III*, Annali di Matematica **138** (1984), 245–253.
- [24] Yue Chi Ming R., *On regular rings and Artinian rings, II*, Riv. Mat. Univ. Parma (4) **11** (1985), 101–109.
- [25] Yue Chi Ming R., *On von Neumann regular rings, XV*, Acta Math. Vietnamica **13** (1988), no. 2, 71–79.
- [26] Yue Chi Ming R., *On V-rings, p -V-rings and injectivity*, Kyungpook Math. J. **32** (1992), 219–228.
- [27] Yue Chi Ming R., *On self-injectivity and regularity*, Rend. Sem. Fac. Sci. Univ. Cagliari **64** (1994), 9–24.
- [28] Zhang Jule, *Fully idempotent rings whose every maximal left ideal is an ideal*, Chinese Sci. Bull. **37** (1992), 1065–1068.
- [29] Zhang Jule, *A note on von Neumann regular rings*, Southeast Asian Bull. Math. **22** (1998), 231–235.
- [30] Zhang Jule, Wu Jun, *Generalizations of principal injectivity*, Algebra Colloq. **6** (1999), 277–282.

UNIVERSITÉ PARIS VII-DENIS DIDEROT, UFR MATHS-UMR 9994 CNRS, 2, PLACE JUSSIEU,
75251 PARIS CEDEX 05, FRANCE

(Received July 19, 2001)