

Kaori Yamazaki

A proof for the Blair-Hager-Johnson theorem on absolute  $z$ -embedding

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 43 (2002), No. 1, 175--179

Persistent URL: <http://dml.cz/dmlcz/119310>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## A proof for the Blair-Hager-Johnson theorem on absolute $z$ -embedding

KAORI YAMAZAKI

*Abstract.* In this paper, a simple proof is given for the following theorem due to Blair [7], Blair-Hager [8] and Hager-Johnson [12]: A Tychonoff space  $X$  is  $z$ -embedded in every larger Tychonoff space if and only if  $X$  is almost compact or Lindelöf. We also give a simple proof of a recent theorem of Bella-Yaschenko [6] on absolute embeddings.

*Keywords:* absolute  $z$ -embedding, absolute  $C$ -embedding, absolute  $C^*$ -embedding, absolute embeddings, almost compact, Lindelöf, compact, pseudocompact

*Classification:* 54C25, 54D20

All spaces are assumed to be Tychonoff spaces. For a subspace  $A$  of a space  $X$ ,  $A$  is said to be  $C$  (resp.  $C^*$ )-*embedded* in  $X$  if every real-valued (resp. every bounded real-valued) continuous function on  $A$  can be continuously extended over  $X$ . A subspace  $A$  of a space  $X$  is said to be  $z$ -*embedded* in  $X$  if for every zero-set  $Z$  of  $A$  there exists a zero-set  $Z'$  of  $X$  with  $Z' \cap A = Z$ . Clearly,  $C$ -embedding implies  $C^*$ -embedding, and the latter implies  $z$ -embedding. As it is known, there have been several results on  $z$ -embeddings, which are closely related to those on  $C$  or  $C^*$ -embeddings and other extension properties (see [1]). Let us recall the following theorem which describes the so-called absolute  $C$ -embedding or absolute  $C^*$ -embedding; a Tychonoff space  $X$  is said to be *almost compact* if  $|\beta X - X| \leq 1$ , where  $\beta X$  denotes the Stone-Čech compactification of  $X$ .

**Theorem 1** (Doss [9], Hewitt [14], Smirnov [16]; see also [1], [11]). *A Tychonoff space  $X$  is  $C$  (or equivalently,  $C^*$ )-embedded in every larger Tychonoff space if and only if  $X$  is almost compact.*

As concerns  $z$ -embeddings, recall the following result due to Jerison (see [13, Lemma 5.3] or [1, Theorem 7.8]).

**Theorem 2** (Jerison). *If  $X$  is a Lindelöf subspace of a Tychonoff space  $Y$ , then  $X$  is  $z$ -embedded in  $Y$ .*

Corresponding to Theorem 1, the result characterizing the so-called absolute  $z$ -embedding has been established by Blair [7], Blair-Hager [8] and Hager-Johnson [12] as the following; the “if” part directly follows from Theorems 1 and 2.

---

This research was partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Encouragement of Young Scientists, No. 13740034, 2001.

**Theorem 3** (Blair [7], Blair-Hager [8], Hager-Johnson [12]). *A Tychonoff space  $X$  is  $z$ -embedded in every larger Tychonoff space if and only if  $X$  is almost compact or Lindelöf.*

The proofs of the “only if” part of Theorem 3 given in [7], [8] and [12] are obtained through several consequences under their own interests on realcompactness or rings of continuous functions. In the present paper, we give an alternative and simple proof to this theorem, with only a use of the following well-known fact: a Tychonoff space  $X$  is Lindelöf if and only if for every compact subspace  $F$  of  $\beta X$  with  $F \subset \beta X - X$  there exists a zero-set  $Z$  of  $\beta X$  such that  $F \subset Z \subset \beta X - X$  (see [10, 3.12.25(b)]). Other terminology may be found in [1], [10] and [11].

**PROOF OF THE “ONLY IF” PART OF THEOREM 3:** Assume that  $X$  is  $z$ -embedded in every larger Tychonoff space. Suppose that  $X$  is not almost compact. We show that  $X$  is Lindelöf. Let  $F$  be a compact subspace of  $\beta X$  with  $F \subset \beta X - X$ .

**Claim.** *For every  $x \in F$ , there exist an open neighborhood  $U_x$  of  $x$  in the subspace  $F$  and a zero-set  $Z_x$  of  $\beta X$  such that  $U_x \subset Z_x \subset \beta X - X$ .*

**PROOF OF CLAIM:** Let  $x \in F$ . Since  $|\beta X - X| \geq 2$ , pick up a point  $y \in \beta X - X$  with  $y \neq x$ . Let  $f : \beta X \rightarrow [0, 1]$  be a continuous function satisfying that  $f(x) = 0$  and  $f(y) = 1$ . Let  $Y = \beta X / (F \cup \{y\})$  be the quotient space obtained from  $\beta X$  by identifying  $F \cup \{y\}$  with a single point and  $q : \beta X \rightarrow Y$  be the natural quotient map. Since  $f^{-1}([0, 1/2]) \cap X$  is a cozero-set of  $X$  and  $X$  is  $z$ -embedded in  $Y$ , there exists a cozero-set  $V$  of  $Y$  such that  $V \cap X = f^{-1}([0, 1/2]) \cap X$ . Then,  $q(x) \notin V$ . Indeed, if  $q(x) \in V$ , then  $y \in q^{-1}(V) \cap f^{-1}((1/2, 1]) \subset \beta X - X$ , a contradiction. Put  $U_x = f^{-1}([0, 1/3]) \cap F$  and  $Z_x = f^{-1}([0, 1/3]) - q^{-1}(V)$ . These are the required sets. This completes the proof of Claim.  $\square$

Finally, for some finite points  $x_1, \dots, x_n \in F$  with  $F = \bigcup_{i=1}^n U_{x_i}$ , put  $Z = \bigcup_{i=1}^n Z_{x_i}$ . Then,  $Z$  is a zero-set of  $\beta X$  and  $F \subset Z \subset \beta X - X$ . Hence  $X$  is Lindelöf. This completes the proof.  $\square$

We apply our technique in the above proof to give a simple proof to the following theorem which was recently obtained by Bella and Yaschenko [6], their proof is long and complicated.

**Theorem 4** (Bella-Yaschenko [6]). *For a Tychonoff space  $X$ , the following statements are equivalent:*

- (1) *if a Tychonoff space  $Y$  contains two disjoint closed copies  $X_1$  and  $X_2$  of  $X$ , then these copies can be separated in  $Y$  by open sets;*
- (2)  *$X$  is Lindelöf.*

See [3] for the motivation of Theorem 4. For the proof, let us consider another condition (\*) below; recall that *separated subsets* mean those subsets  $A$  and  $B$  of  $X$  with  $\overline{A} \cap B = A \cap \overline{B} = \emptyset$ .

(\*) *If a Tychonoff space  $Y$  contains two copies  $X_1$  and  $X_2$  of  $X$  which are separated subsets, then these copies can be separated in  $Y$  by cozero-sets.*

PROOF OF THEOREM 4: (1)  $\Rightarrow$  (2): Assume (1). We shall first prove that (\*) holds. Let  $X_1$  and  $X_2$  be copies of  $X$  and assume that they are separated subsets of a Tychonoff space  $Y$ . Embed  $Y$  into the Tychonoff cube  $T$ . Moreover, embed  $T$  into the product space  $T \times [0, 1]$  as a subspace  $T \times \{0\}$  and denote the subspace  $(T \times (0, 1]) \cup ((X_1 \cup X_2) \times \{0\})$  by  $Z$ . Then,  $X_1$  and  $X_2$  are disjoint closed subsets in  $Z$ . From the assumption,  $X_i \times \{0\}$ ,  $i = 1, 2$ , can be separated by open sets  $U_1$  and  $U_2$  in  $Z$ . Since  $Z$  is a dense subset of  $T \times [0, 1]$ , these open sets can be extended to disjoint open sets  $U_1^*$  and  $U_2^*$  in  $T \times I$ . Since  $T \times I$  is a Tychonoff cube, it is known that  $U_1^*$  and  $U_2^*$  can be separated by cozero-sets  $G_1$  and  $G_2$  in  $T \times I$ , use [15, Theorem 2] and [17, Theorem 2]. Then, it follows that  $X_i \times \{0\} \subset G_i \cap (Y \times \{0\})$ ,  $i = 1, 2$ . Hence,  $X_1$  and  $X_2$  can be separated by cozero-sets in  $Y$ . So, (\*) holds.

We now prove  $X$  is Lindelöf. Let  $F$  be a compact subspace of  $\beta X$  with  $F \subset \beta X - X$  and let  $X_1 = X_2 = X$ . Denote  $F$  in  $\beta X_i$  by  $F_i$ ,  $i = 1, 2$ . Let  $Y = (\beta X_1 \oplus \beta X_2) / (F_1 \cup F_2)$  be the quotient space obtained from  $\beta X_1 \oplus \beta X_2$  identifying  $F_1 \cup F_2$  to a single point, and  $q : \beta X_1 \oplus \beta X_2 \rightarrow Y$  the natural quotient map. Since  $q(X_1)$  and  $q(X_2)$  are separated subsets in  $Y$ , by (\*), there exists disjoint cozero-sets  $U_1$  and  $U_2$  of  $Y$  such that  $q(X_i) \subset U_i$ ,  $i = 1, 2$ . We may assume  $q(F_1 \cup F_2) \not\subset U_1$ . Then,  $X_1 \subset q^{-1}(U_1) \cap \beta X_1 \subset \beta X_1 - F_1$  and  $q^{-1}(U_1) \cap \beta X_1$  is a cozero-set of  $\beta X_1$ . Hence  $X_1$  is (that is,  $X$  is) Lindelöf.

(2)  $\Rightarrow$  (1): Easy. □

It should be noted in Theorem 1 (resp. Theorem 3) that  $X$  may be equivalently assumed to be  $C$  or  $C^*$  (resp.  $z$ -)embedded in every Tychonoff space in which  $X$  is embedded as a closed subset ([4], [14], [16]). This can be proved similarly to the above proof showing (\*).

Yajima has given some generalizations of Theorem 4 and characterizations of paracompactness [18].

Next we give some results related to Theorem 4. The technique of making the adjunction space determined by a space and the Tychonoff plank, which is popular in the theory of relative topological properties (cf. [2]), will be used.

**Theorem 5.** *For a Tychonoff space  $X$ , the following statements are equivalent:*

- (1) *if a Tychonoff space  $Y$  contains two disjoint closed copies  $X_1$  and  $X_2$  of  $X$ , then these copies can be completely separated in  $Y$ ;*
- (2)  *$X$  is compact.*

PROOF: (1)  $\Rightarrow$  (2): Assume (1). By Theorem 4,  $X$  is Lindelöf. It suffices to show that every closed discrete set of  $X$  is finite. To prove this, assume the contrary and let  $\{x_n : n < \omega\}$  be a closed discrete set in  $X$  consisting of distinct points.

Let  $X_1 = X_2 = X$  and denote  $\{x_n : n < \omega\}$  in  $X_i$  by  $\{x_n^i : n < \omega\}$ ,  $i = 1, 2$ . Let  $Z = (\omega_1 + 1) \times (\omega + 1) - \{(\omega_1, \omega)\}$  be the Tychonoff plank. Let  $Z_1 = Z_2 = Z$ , and denote the right edge  $\{(\omega_1, n) : n < \omega\}$  of  $Z_i$  by  $\{(\omega_1, n)^i : n < \omega\}$ , and the top edge  $\{(\alpha, \omega) : \alpha < \omega_1\}$  of  $Z_i$  by  $\{(\alpha, \omega)^i : \alpha < \omega_1\}$ ,  $i = 1, 2$ . For  $i = 1, 2$ , define a map  $f_i : \{x_n^i : n < \omega\} \rightarrow Z_i$  by  $f_i(x_n^i) = (\omega_1, n)^i$ ,  $n < \omega$ . Consider the adjunction spaces  $X_i \cup_{f_i} Z_i$ ,  $i = 1, 2$  (see [10, p. 93]). Define a map  $g : \{(\alpha, \omega)^1 : \alpha < \omega_1\} \rightarrow X_2 \cup_{f_2} Z_2$  by  $g((\alpha, \omega)^1) = (\alpha, \omega)^2$ ,  $\alpha < \omega_1$ . Let  $Y$  be the adjunction space  $(X_1 \cup_{f_1} Z_1) \cup_g (X_2 \cup_{f_2} Z_2)$ . Since  $Y$  is a Tychonoff space and  $X_1$  and  $X_2$  are closed subsets of  $Y$ , by the assumption,  $X_1$  and  $X_2$  must be completely separated in  $Y$ . But this is a contradiction.

(2)  $\Rightarrow$  (1): Easy. □

The following result should be compared with Theorems 4 and 5; this is probably known and is proved similarly to Theorem 5.

**Theorem 6.** *For a Tychonoff space  $X$ , the following statements are equivalent:*

- (1) *if a Tychonoff space  $Y$  contains a copy  $X_1$  of  $X$  and a closed subset  $F$  disjoint from  $X_1$ , then  $X_1$  and  $F$  can be completely separated in  $Y$ ;*
- (2) *if a Tychonoff space  $Y$  contains a closed copy  $X_1$  of  $X$  and a closed subset  $F$  disjoint from  $X_1$ , then  $X_1$  and  $F$  can be separated in  $Y$  by open sets;*
- (3)  *$X$  is compact.*

Analogously, the following holds: (1)  $\Leftrightarrow$  (3) is due to Blair-Hager [8, Proposition 4.3], (2)  $\Leftrightarrow$  (3) is due to Aull [5, Theorem 1(b)]. Here, we give a direct proof of (2)  $\Rightarrow$  (3).

**Theorem 7** (Blair-Hager [8], Aull [5]). *For a Tychonoff space  $X$ , the following statements are equivalent:*

- (1) *if a Tychonoff space  $Y$  contains a copy  $X_1$  of  $X$  and a zero-set  $F$  disjoint from  $X_1$ , then  $X_1$  and  $F$  can be completely separated in  $Y$ ;*
- (2) *if a Tychonoff space  $Y$  contains a closed copy  $X_1$  of  $X$  and a zero-set  $F$  disjoint from  $X_1$ , then  $X_1$  and  $F$  can be separated in  $Y$  by open sets;*
- (3)  *$X$  is pseudocompact.*

PROOF: (1)  $\Rightarrow$  (2): Obvious.

(2)  $\Rightarrow$  (3): Assume (2) and assume  $X$  is not pseudocompact. Let  $f : X \rightarrow (0, +\infty)$  be a positive unbounded continuous function. Then, there exists a closed discrete subset  $\{x_n : n < \omega\}$  of  $X$  such that  $n \leq f(x_n)$  for every  $n < \omega$ . Consider the adjunction space  $X \cup_g Z$ , where  $Z$  is the Tychonoff plank, and  $g : \{x_n : n < \omega\} \rightarrow Z$  is a map defined by  $g(x_n) = (\omega_1, n)$ ,  $n < \omega$ . Then, notice that the top edge of  $Z$  is a zero-set of  $X \cup_g Z$ . Indeed, define a continuous function  $h : X \cup_g Z \rightarrow [0, +\infty)$  by  $h(x) = 1/f(x)$  if  $x \in X$ ;  $h(x) = 1/f(x_n)$  if  $x = (\alpha, n) \in Z$ ,  $\alpha \leq \omega_1$ ,  $n < \omega$ ;  $h(x) = 0$  if  $x = (\alpha, \omega) \in Z$ ,  $\alpha < \omega_1$ . Since the top edge of  $Z$  is  $h^{-1}(\{0\})$ , the condition (2) raises a contradiction.

(3)  $\Rightarrow$  (1): See [8]. □

By [8, Corollary 3.6] (see also [1]), a subspace  $A$  of a space  $X$  is  $C$ -embedded in  $X$  if and only if  $A$  is  $z$ -embedded and well-embedded in  $X$ , where  $A$  is said to be *well-embedded* in  $X$  if any zero-set of  $X$  disjoint from  $A$  can be completely separated in  $X$ . Hence, it should be noted that Theorems 3 and 7 induce Theorem 1.

#### REFERENCES

- [1] Alb R.A., Shapiro H.L., *Normal Topological Spaces*, Cambridge University Press, Cambridge, 1974.
- [2] Arhangel'skii A.V., *Relative topological properties and relative topological spaces*, *Topology Appl.* **70** (1996), 87–99.
- [3] Arhangel'skii A.V., Tartir J., *A characterization of compactness by relative separation property*, *Questions Answers Gen. Topology* **14** (1996), 49–52.
- [4] Aull C.E., *Some embeddings related to  $C^*$ -embeddings*, *J. Austral. Math. Soc. (Series A)* **44** (1988), 88–104.
- [5] Aull C.E., *On well-embedding*, *General Topology and Applications* (Middletown, CT, 1988), pp. 1–5; *Lecture Notes in Pure and Appl. Math.*, 123, Dekker, New York, 1990.
- [6] Bella A., Yaschenko I.V., *Lindelöf property and absolute embeddings*, *Proc. Amer. Math. Soc.* **127** (1999), 907–913.
- [7] Blair R.L., *On  $v$ -embedded sets in topological spaces*, *TOPO* 72 — *General Topology and its Applications* (Proc. Second Pittsburgh Internat. Conf., Pittsburgh, Pa., 1972; dedicated to the memory of Johannes H. de Groot), pp. 46–79; *Lecture Notes in Math.*, Vol. 378, Springer, Berlin, 1974.
- [8] Blair R.L., Hager A.W., *Extensions of zero-sets and of real-valued functions*, *Math. Z.* **136** (1974), 41–52.
- [9] Doss R., *On uniform spaces with a unique structure*, *Amer. J. Math.* **71** (1949), 19–23.
- [10] Engelking R., *General Topology*, Heldermann Verlag, Berlin, 1989.
- [11] Gillman L., Jerison M., *Rings of Continuous Functions*, Van Nostrand, Princeton, 1960.
- [12] Hager A.W., Johnson D.G., *A note on certain subalgebras of  $C(X)$* , *Canad. J. Math.* **20** (1968), 389–393.
- [13] Henriksen M., Johnson D.G., *On the structure of a class of archimedean lattice-ordered algebras*, *Fund. Math.* **50** (1961), 73–94.
- [14] Hewitt E., *A note on extensions of continuous functions*, *An. Acad. Brasil. Ci.* **21** (1949), 175–179.
- [15] Noble N.,  *$C$ -embedded subsets of products*, *Proc. Amer. Math. Soc.* **31** (1972), 613–614.
- [16] Smirnov Y., *Mappings of systems of open sets* (in Russian), *Mat. Sb.* **31** (1952), 152–166.
- [17] Terada T., *Note on  $z$ -,  $C^*$ -, and  $C$ -embedding*, *Sci. Rep. Tokyo Kyoiku Daigaku Sect. A.* **13** (1975), 129–132.
- [18] Yajima Y., *Characterizations of paracompactness and Lindelöfness by separation property*, preprint.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF TSUKUBA, TSUKUBA, IBARAKI 305-8571, JAPAN  
*E-mail:* kaori@math.tsukuba.ac.jp

(Received May 9, 2001, revised November 22, 2001)