

Raushan Z. Buzyakova

On D -property of strong Σ spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 43 (2002), No. 3, 493--495

Persistent URL: <http://dml.cz/dmlcz/119338>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On D -property of strong Σ spaces

RAUSHAN Z. BUZYAKOVA

Abstract. It is shown that every strong Σ space is a D -space. In particular, it follows that every paracompact Σ space is a D -space.

Keywords: strong Σ space, D -space

Classification: 54D20, 54F99

In this paper we will show that any strong Σ space is a D -space. This result positively answers Borges and Matveev's question whether any paracompact Σ space is a D -space. The notion of D -space was introduced by Eric van Douwen [6].

A *neighborhood assignment* for a space X is a function φ from X to the topology of X such that $x \in \varphi(x)$ for any $x \in X$. A space X is a D -space, if for any neighborhood assignment φ for X there exists a closed discrete subset D of X such that $X = \bigcup_{d \in D} \varphi(d)$.

It is natural to ask which spaces possess the D -property. It is known that σ -compact spaces, metrizable spaces, semi-stratifiable spaces, and paracompact p -spaces are all D -spaces (see [5], [2]). In [5], DeCaux showed that every finite product of copies of the Sorgenfrey line is a D -space. The D -property of subspaces of generalized ordered spaces was studied in [8]. In a recent paper [10] of Fleissner and Stanley, the authors give conditions under which a subspace of a product of finitely many ordinals is a D -space. Several interesting questions on D -spaces were raised by E. van Douwen and W.F. Pfeffer in [7], which was the first published paper that contained results on D -spaces. Some other results and questions on D -spaces can be found in [5], [2], [3], [4], [8], [10].

The result in this article is obtained in an attempt to answer E.K. van Douwen's question whether each Lindelöf space is a D -space. However, this question remains unanswered. And, one of approaches to solve this problem could be to consider continuous images of Lindelöf D -spaces.

Question (A. V. Arhangel'skii). *Is it true that a continuous image of a Lindelöf D -space is a D -space?*

We consider only Tychonoff spaces. In notation and terminology, we will follow [9].

A space X is a *strong* Σ space if there exist a σ -locally-finite family γ of closed sets in X and a cover \mathcal{K} of X by compact subsets, such that for any open set U containing an element K of \mathcal{K} , $K \subseteq \Gamma \subseteq U$ for some $\Gamma \in \gamma$.

The class of strong Σ spaces is wide and it contains all metrizable spaces, σ -compact spaces, Lindelöf Σ spaces, paracompact Σ spaces, paracompact p -spaces, Moore spaces, spaces with countable network, as well as spaces with σ -discrete network (σ spaces). Thus, our result implies that the mentioned spaces are all D -spaces. In addition, a finite (countable) product of strong Σ spaces is a D -space as well, since the class of strong Σ spaces is closed with respect to countable products. Therefore, in particular, the product of a Lindelöf Σ space with a Moore space is still a D -space. However, as shown in [4], in general case the product of two D -spaces need not be a D -space.

Theorem. *Every strong Σ space X is a D -space.*

PROOF: Let \mathcal{K} and γ be the families from the definition of a strong Σ space. Represent γ as $\bigcup\{\gamma_n\}$, where each γ_n is a locally-finite family of closed sets in X and $\gamma_n \subseteq \gamma_{n+1}$. Enumerate each $\gamma_n = \{\Gamma_\alpha^n\}$, where α ranges through some ordinal number.

Let φ be an arbitrary neighborhood assignment. We need to find a discrete closed subset D in X such that $X = \bigcup_{d \in D} \varphi(d)$. Recursively, we will define closed discrete sets D_n such that $D = \bigcup D_n$.

Step 0. Set $D_0 = \emptyset$.

Assume D_m is defined for all $0 < m < n$.

Step n. Recursively, we will define finite sets D_α^n such that $D_n = (\bigcup D_\alpha^n) \cup D_{n-1}$.

Sub-step 0. Set $D_0^n = \emptyset$.

Assume D_β^n is defined for all $0 < \beta < \alpha$.

Sub-step α . Let $U = \bigcup\{\varphi(d) : d \in (\bigcup_{\beta < \alpha} D_\beta^n) \cup D_{n-1}\}$. Take the first Γ in γ_n that satisfies the following requirement.

Requirement R_α^n : there exists $K \in \mathcal{K}$ which is not fully covered by U . And there exist $x_1, \dots, x_k \in K \setminus U$ such that $K \setminus U \subseteq \Gamma \setminus U \subseteq \varphi(x_1) \cup \dots \cup \varphi(x_k)$.

If no such Γ exists, sub-recursion stops. Put $D_\alpha^n = \{x_1, \dots, x_k\}$.

Let $D_n = (\bigcup D_\alpha^n) \cup D_{n-1}$. We need to show that D_n is closed and discrete in X . Take an arbitrary $x \in X$. We need to separate x from $D_n \setminus \{x\}$ by a neighborhood. Consider the family

$$\gamma'_n = \{\Gamma_\beta : \Gamma_\beta \text{ is the first in } \gamma_n \text{ satisfying Requirement } R_\alpha^n \text{ for some } \alpha\}.$$

Since $\gamma'_n \subseteq \gamma_n$, γ'_n is locally-finite too. Therefore, there exists a neighborhood of x that intersects only a finite number of elements in γ'_n , and therefore, only

finite number of sets D_α^n 's. Since the D_α^n 's are finite, x is not in the closure of $(\bigcup D_\alpha^n) \setminus \{x\}$. And x can be separated from $D_{n-1} \setminus \{x\}$ since the latter is closed and discrete by assumption.

The construction is complete. Put $D = \bigcup D_n$.

Let us show that $X = \bigcup_{d \in D} \varphi(d)$. Assume the contrary. Then there exists a K in \mathcal{K} such that $K' = K \setminus \bigcup_{d \in D} \varphi(d) \neq \emptyset$. Since K' is compact we can find $x_1, \dots, x_k \in K'$ such that $K' \subseteq \varphi(x_1) \cup \dots \cup \varphi(x_k)$. Consider a compactum $K'' = K \setminus (\varphi(x_1) \cup \dots \cup \varphi(x_k))$. Find the smallest n such that $K'' \subseteq \bigcup_{d \in D_n} \varphi(d)$. Now take the first γ_l containing such a Γ that

$$K \subseteq \Gamma \subseteq \varphi(x_1) \cup \dots \cup \varphi(x_k) \cup \left(\bigcup_{d \in D_n} \varphi(d) \right).$$

Let $m = \max\{n, l\}$. Then $\gamma_l \subseteq \gamma_{m+1}$, and therefore, $\Gamma \in \gamma_{m+1}$. By the choice of n and l , Γ satisfies the *Requirement* starting not later than from Sub-step 1 of Step $m+1$. And eventually, Γ will be the first in γ_{m+1} satisfying the *Requirement*. Therefore, Γ must be covered by $\bigcup_{d \in D} \varphi(d)$, and so must K .

Let us show now that D is closed and discrete. Take an arbitrary $x \in X$. We need to show that x can be separated from $D \setminus \{x\}$ by a neighborhood of x . There exists an n such that $x \in \bigcup_{d \in D_n} \varphi(d)$. This means that x is separated from $D \setminus D_n$ by $\bigcup_{d \in D_n} \varphi(d)$ (follows from the construction of D_n 's). And x can be separated from $D_n \setminus \{x\}$, since D_n is closed and discrete. \square

REFERENCES

- [1] Arhangel'skii A., *private communications*, 2001.
- [2] Borges C.R., Wehrly A.C., *A study of D-spaces*, Topology Proc. **16** (1991), 7–15.
- [3] Borges C.R., Wehrly A.C., *Another study of D-spaces*, Questions Answers Gen. Topology **14:1** (1996), 73–76.
- [4] Borges C.R., Wehrly A.C., *Correction: another study of D-spaces*, Questions Answers Gen. Topology **16:1** (1998), 77–78.
- [5] DeCaux P., *Yet another property of the Sorgenfrey line*, Topology Proc. **6:1** (1981), 31–43.
- [6] van Douwen E.K., *Simultaneous extension of continuous functions*, Thesis, Free University, Amsterdam, 1975.
- [7] van Douwen E.K., Pfeffer W.F., *Some properties of the Sorgenfrey line and related spaces*, Pacific J. Math. **81** (1979), 371–377.
- [8] van Douwen E.K., Lutzer D.J., *A note on paracompactness in generalized ordered spaces*, Proc. Amer. Math. Soc. **125** (1997), 1237–1245.
- [9] Engelking R., *General Topology*, Sigma Series in Pure Mathematics, 6, Heldermann, Berlin, revised ed., 1989.
- [10] Fleissner W.G., Stanley A.M., *D-spaces*, Topology Appl. **114** (2001), 261–271.

MATHEMATICS DEPARTMENT, BROOKLYN COLLEGE, BROOKLYN, 11210 USA

E-mail: RaushanB@brooklyn.cuny.edu

(Received December 20, 2001)