

Abraham A. Ungar

The hyperbolic triangle centroid

Commentationes Mathematicae Universitatis Carolinae, Vol. 45 (2004), No. 2, 355--369

Persistent URL: <http://dml.cz/dmlcz/119464>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2004

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

The hyperbolic triangle centroid

ABRAHAM A. UNGAR

Abstract. Some gyrocommutative gyrogroups, also known as Bruck loops or K-loops, admit scalar multiplication, turning themselves into gyrovector spaces. The latter, in turn, form the setting for hyperbolic geometry just as vector spaces form the setting for Euclidean geometry. In classical mechanics the centroid of a triangle in velocity space is the velocity of the center of momentum of three massive objects with equal masses located at the triangle vertices. Employing gyrovector space techniques we find in this article that, in full analogy, the centroid of a hyperbolic triangle in relativity velocity space is the velocity of the center of momentum of three massive objects with equal rest masses located at the triangle vertices. Being guided by the relativistic mass correction of moving massive objects in special relativity theory, we express the hyperbolic triangle centroid in terms of the triangle vertices, resulting in a novel hyperbolic triangle centroid identity that captures remarkable analogies with its Euclidean counterpart.

Keywords: loops, gyrogroups, gyrovector spaces, hyperbolic geometry, Einstein addition, Möbius transformation

Classification: 20N05, 51P05, 83A05

1. Introduction

Einstein addition of relativistic velocities in special relativity theory and Möbius addition in the theory of complex functions are isomorphic binary operations in a grouplike structure that turns out to be a loop. Einstein addition is straightforwardly generalized in Section 5 into a binary operation \oplus_E in the open unit ball \mathbb{B} of any real inner product space \mathbb{V} , giving rise to the Einstein groupoid (\mathbb{B}, \oplus_E) , presented in Section 5. We recall that a groupoid is a nonempty set with a binary operation. Möbius addition in the complex open unit disc is a Möbius transformation of the disc without rotation. It is generalized in [21] into a binary operation, \oplus_M , in the open unit ball \mathbb{B} of any real inner product space \mathbb{V} . It gives rise to the Möbius groupoid (\mathbb{B}, \oplus_M) , presented in Section 3.

Both Möbius addition and Einstein addition are neither commutative nor associative. Accordingly, the groupoids of Einstein and Möbius do not form groups. These two groupoids share a common grouplike structure called a *gyrocommutative gyrogroup*. The latter turns out to be equivalent to the Bruck loop. Unlike the Bruck loop definition [12], however, the gyrocommutative gyrogroup definition, Section 2, emphasizes analogies with groups.

Endowing a gyrocommutative gyrogroup with scalar multiplication and inner product, one obtains a gyrovector space. Remarkably, gyrovector spaces form the setting for hyperbolic geometry just as vector spaces form the setting for Euclidean geometry. In particular, the resulting Möbius and Einstein gyrovector spaces are studied in this article in order to set the stage for the study of hyperbolic geometry analytically, allowing the hyperbolic triangle centroid to be determined analytically in Sections 6 and 7. An earlier study of the hyperbolic triangle centroid is found in [2].

Möbius gyrovector spaces form the setting for the Poincaré ball model of hyperbolic geometry, as shown in Sections 3 and 4. Similarly, Einstein gyrovector spaces form the setting for the Beltrami (also known as Klein) ball model of hyperbolic geometry, as shown in Sections 5 and 6.

The study of hyperbolic geometry in terms of its two isomorphic models of Poincaré and Beltrami is particularly useful. On the one hand, the Poincaré model is conformal, so that hyperbolic angles between intersecting geodesics have the same measure as Euclidean angles between corresponding intersecting tangent lines [23, Figure 6.14]. On the other hand, geodesics in the Beltrami model are Euclidean straight lines, allowing one to employ linear algebra in the determination of points of intersection of geodesics. Indeed, in this article we use linear algebra to obtain the triangle centroid as the point of intersection of the triangle medians in the Beltrami model, Figure 3, and translate the result into the Poincaré model, Figure 4.

2. Gyrogroups and gyrovector spaces

Definition 1 (Gyrogroups). The groupoid (G, \oplus) is a *gyrogroup* if its binary operation satisfies the following axioms. In G there is at least one element, 0 , called a left identity, satisfying

$$(G1) \quad 0 \oplus a = a \qquad \text{Left Identity}$$

for all $a \in G$. There is an element $0 \in G$ satisfying axiom (G1) such that for each a in G there is an element $\ominus a$ in G , called a left inverse of a , with

$$(G2) \quad \ominus a \oplus a = 0. \qquad \text{Left Inverse}$$

Moreover, for any $a, b, z \in G$ there exists a unique element $\text{gyr}[a, b]z \in G$ such that

$$(G3) \quad a \oplus (b \oplus z) = (a \oplus b) \oplus \text{gyr}[a, b]z. \qquad \text{Left Gyroassociative Law}$$

If $\text{gyr}[a, b]$ denotes the map $\text{gyr}[a, b] : G \rightarrow G$ given by $z \mapsto \text{gyr}[a, b]z$ then

$$(G4) \quad \text{gyr}[a, b] \in \text{Aut}(G, \oplus) \qquad \text{Gyroautomorphism}$$

and $\text{gyr}[a, b]$ is called the *Thomas gyration*, or the *gyroautomorphism* of G , generated by $a, b \in G$. The operation $\text{gyr} : G \times G \rightarrow \text{Aut}(G, \oplus)$ is called the

gyrooperation of G . Finally, the gyroautomorphism $\text{gyr}[a, b]$ generated by any $a, b \in G$ satisfies

$$(G5) \quad \text{gyr}[a, b] = \text{gyr}[a \oplus b, b]. \quad \text{Left Loop Property}$$

Various gyrogroup theorems are presented in [23]. Thus, for instance, any gyrogroup possesses a right identity and a right inverse as well, which are identical to their left counterparts. Furthermore, the resulting identity element and the inverse of any given element are unique. In full analogy with groups, gyrogroups are classified into gyrocommutative and non-gyrocommutative gyrogroups.

Definition 2 (Gyrocommutative gyrogroups). The gyrogroup (G, \oplus) is *gyrocommutative* if for all $a, b \in G$

$$(G6) \quad a \oplus b = \text{gyr}[a, b](b \oplus a). \quad \text{Gyrocommutative Law}$$

A gyrogroup is a loop [23], and the gyrocommutative gyrogroup is equivalent to the Bruck loop. Following Ungar [12], [20], the latter is also known as a K-loop. Furthermore, a gyrocommutative gyrogroup is also a special Bol loop that possesses the automorphic inverse property, $\ominus(a \oplus b) = \ominus a \ominus b$ [18], [10]. Gyrocommutative gyrogroups result from transversals to subgroups, as shown in [8], [9]. Indeed, transversals to subgroups [4] and transversals in loops [14] are important in loop theory.

Definition 3 (Inner product gyrovectors spaces). A(n inner product) gyrovectors space (G, \oplus, \otimes) is a gyrocommutative gyrogroup (G, \oplus) that admits:

- (1) Inner product \cdot, \cdot , (i) which gives rise to a positive definite norm $\|\mathbf{v}\|$, that is, $\|\mathbf{v}\|^2 = \mathbf{v} \cdot \mathbf{v}$, $\|\mathbf{v}\| \geq 0$ and $\|\mathbf{v}\| = 0$ if and only if $\mathbf{v} = \mathbf{0}$, $|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\| \|\mathbf{v}\|$; and (ii) which is invariant under gyroautomorphisms, that is,

$$\text{gyr}[\mathbf{a}, \mathbf{b}]\mathbf{u} \cdot \text{gyr}[\mathbf{a}, \mathbf{b}]\mathbf{v} = \mathbf{u} \cdot \mathbf{v}$$

for all gyrovectors $\mathbf{a}, \mathbf{b}, \mathbf{u}, \mathbf{v} \in G$;

- (2) Scalar multiplication, \otimes , satisfying the following properties. For all real numbers $r, r_1, r_2 \in \mathbb{R}$ and all gyrovectors $\mathbf{a}, \mathbf{b}, \mathbf{v} \in G$:

- (V1) $1 \otimes \mathbf{v} = \mathbf{v}$,
- (V2) $(r_1 + r_2) \otimes \mathbf{v} = r_1 \otimes \mathbf{v} \oplus r_2 \otimes \mathbf{v}$, Scalar Distributive Law
- (V3) $(r_1 r_2) \otimes \mathbf{v} = r_1 \otimes (r_2 \otimes \mathbf{v})$, Scalar Associative Law
- (V4) $\frac{|r| \otimes \mathbf{v}}{\|r \otimes \mathbf{v}\|} = \frac{\mathbf{v}}{\|\mathbf{v}\|}$, Scaling Property
- (V5) $\text{gyr}[\mathbf{a}, \mathbf{b}](r \otimes \mathbf{v}) = r \otimes \text{gyr}[\mathbf{a}, \mathbf{b}]\mathbf{v}$, Gyroautomorphism Property
- (V6) $\text{gyr}[r_1 \otimes \mathbf{v}, r_2 \otimes \mathbf{v}] = I$; Identity Automorphism

- (3) Real vector space structure $(\|G\|, \oplus, \otimes)$ for the set $\|G\|$ of one-dimensional ‘vectors’

$$\|G\| = \{\pm \|\mathbf{v}\| : \mathbf{v} \in G\} \subset \mathbb{R}$$

with vector addition \oplus and scalar multiplication \otimes , such that for all $r \in \mathbb{R}$ and $\mathbf{u}, \mathbf{v} \in G$,

$$\begin{aligned} (V7) \quad & \|r \otimes \mathbf{v}\| = |r| \otimes \|\mathbf{v}\|, && \text{Homogeneity Property} \\ (V8) \quad & \|\mathbf{u} \oplus \mathbf{v}\| \leq \|\mathbf{u}\| \oplus \|\mathbf{v}\|. && \text{Gyrotriangle inequality} \end{aligned}$$

A gyrovector space $G = (G, \oplus, \otimes)$ is a gyrometric space, with gyrometric given by the distance function

$$(1) \quad d(\mathbf{u}, \mathbf{v}) = \|\ominus \mathbf{u} \oplus \mathbf{v}\| = \|\mathbf{v} \ominus \mathbf{u}\|$$

satisfying the gyrotriangle inequality

$$(2) \quad \|\ominus \mathbf{u} \oplus \mathbf{w}\| \leq \|\ominus \mathbf{u} \oplus \mathbf{v}\| \oplus \|\ominus \mathbf{v} \oplus \mathbf{w}\|$$

verified below.

By a gyrogroup identity [23] we have

$$(3) \quad \ominus \mathbf{u} \oplus \mathbf{w} = (\ominus \mathbf{u} \oplus \mathbf{v}) \oplus \text{gyr}[\mathbf{u}, \ominus \mathbf{v}](\ominus \mathbf{v} \oplus \mathbf{w}).$$

Hence, by the gyrotriangle inequality (V8) we have

$$\begin{aligned} (4) \quad \|\ominus \mathbf{u} \oplus \mathbf{w}\| &= \|(\ominus \mathbf{u} \oplus \mathbf{v}) \oplus \text{gyr}[\mathbf{u}, \ominus \mathbf{v}](\ominus \mathbf{v} \oplus \mathbf{w})\| \\ &\leq \|\ominus \mathbf{u} \oplus \mathbf{v}\| \oplus \|\text{gyr}[\mathbf{u}, \ominus \mathbf{v}](\ominus \mathbf{v} \oplus \mathbf{w})\| \\ &= \|\ominus \mathbf{u} \oplus \mathbf{v}\| \oplus \|\ominus \mathbf{v} \oplus \mathbf{w}\|. \end{aligned}$$

Our ambiguous use of \oplus and \otimes , Definition 3, as operations in the gyrovector space (G, \oplus, \otimes) and in the vector space $(\|G\|, \oplus, \otimes)$ should raise no confusion, since the sets in which these operations operate are always clear from the context. The operations in the former are nonassociative-nondistributive gyrovector space operations, and in the latter are associative-distributive vector space operations. Additionally, the gyro-addition \oplus is gyrocommutative in the former and commutative in the latter.

An inner product gyrovector space possesses a weak form of a distributive law,

$$(5) \quad r \otimes (r_1 \otimes \mathbf{v} \oplus r_2 \otimes \mathbf{v}) = r \otimes (r_1 \otimes \mathbf{v}) \oplus r \otimes (r_2 \otimes \mathbf{v})$$

called the *monodistributive law*, which follows from (V2) and (V3),

$$\begin{aligned} (6) \quad r \otimes (r_1 \otimes \mathbf{v} \oplus r_2 \otimes \mathbf{v}) &= r \otimes \{(r_1 + r_2) \otimes \mathbf{v}\} \\ &= (r(r_1 + r_2)) \otimes \mathbf{v} \\ &= (rr_1 + rr_2) \otimes \mathbf{v} \\ &= (rr_1) \otimes \mathbf{v} \oplus (rr_2) \otimes \mathbf{v} \\ &= r \otimes (r_1 \otimes \mathbf{v}) \oplus r \otimes (r_2 \otimes \mathbf{v}). \end{aligned}$$

3. Möbius gyrovector spaces

Definition 4 (Möbius addition). Let \mathbb{V} be a real inner product space, and let $\mathbb{B} = \{\mathbf{v} \in \mathbb{V} : \|\mathbf{v}\| < 1\}$ be the open unit ball of \mathbb{V} . Möbius addition \oplus_M in the ball \mathbb{B} is a binary operation in \mathbb{B} given by the equation

$$(7) \quad \mathbf{u} \oplus_M \mathbf{v} = \frac{(1 + 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{v}\|^2)\mathbf{u} + (1 - \|\mathbf{u}\|^2)\mathbf{v}}{1 + 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{u}\|^2\|\mathbf{v}\|^2}$$

where \cdot and $\|\cdot\|$ are the inner product and norm that the ball \mathbb{B} inherits from its space \mathbb{V} .

To justify calling \oplus_M in Definition 4 a Möbius addition we note that it is a natural extension of a special Möbius transformation of the complex open unit disc, as explained in [21], [23]. In earlier studies by Ahlfors [1] and Ratcliffe [17], Möbius addition is treated as a hyperbolic translation. Möbius translation became Möbius addition in [21] following the discovery of the analogies it shares, as a gyrocommutative gyrogroup operation, with ordinary vector addition. Applications of Möbius addition and its hyperbolic geometry in quantum mechanics are found in [3], [15], [16], [24].

The groupoid (\mathbb{B}, \oplus_M) is a gyrocommutative gyrogroup, as demonstrated in [23], called a Möbius gyrogroup. Furthermore, it admits scalar multiplication \otimes , turning itself into the Möbius gyrovector space $(\mathbb{B}, \oplus_M, \otimes)$.

Definition 5 (Möbius scalar multiplication). Let (\mathbb{B}, \oplus_M) be a Möbius gyrogroup. The Möbius scalar multiplication $r \otimes \mathbf{v} = \mathbf{v} \otimes r$ in \mathbb{B} is given by the equation

$$(8) \quad \begin{aligned} r \otimes \mathbf{v} &= \frac{(1 + \|\mathbf{v}\|)^r - (1 - \|\mathbf{v}\|)^r}{(1 + \|\mathbf{v}\|)^r + (1 - \|\mathbf{v}\|)^r} \frac{\mathbf{v}}{\|\mathbf{v}\|} \\ &= \tanh(r \tanh^{-1} \|\mathbf{v}\|) \frac{\mathbf{v}}{\|\mathbf{v}\|} \end{aligned}$$

where $r \in \mathbb{R}$, $\mathbf{v} \in \mathbb{B}$, $\mathbf{v} \neq \mathbf{0}$; and $r \otimes \mathbf{0} = \mathbf{0}$.

As an example we present the Möbius half,

$$(9) \quad \frac{1}{2} \otimes \mathbf{v} = \frac{\gamma_{\mathbf{v}}}{1 + \gamma_{\mathbf{v}}} \mathbf{v}$$

where $\gamma_{\mathbf{v}} = (1 - \|\mathbf{v}\|^2)^{-1/2}$. In accordance with the scalar associative law of gyrovector spaces we have

$$(10) \quad \begin{aligned} 2 \otimes (\frac{1}{2} \otimes \mathbf{v}) &= 2 \otimes \frac{\gamma_{\mathbf{v}}}{1 + \gamma_{\mathbf{v}}} \mathbf{v} \\ &= \frac{\gamma_{\mathbf{v}}}{1 + \gamma_{\mathbf{v}}} \mathbf{v} \oplus_M \frac{\gamma_{\mathbf{v}}}{1 + \gamma_{\mathbf{v}}} \mathbf{v} \\ &= \mathbf{v}. \end{aligned}$$




Figure 1: A Möbius triangle Δabc in the Möbius gyrovector plane $(\mathbb{B}, \oplus, \otimes)$, $\oplus = \oplus_M$ is shown. Its sides are formed by geodesic segments that link its vertices \mathbf{a} , \mathbf{b} and \mathbf{c} , having the hyperbolic lengths a, b and c . The cosine of its angles are given by an identity that is fully analogous to its Euclidean counterpart. The Möbius gyrovector plane forms the setting for the Poincaré disc model of hyperbolic geometry just as the common vector plane forms the setting for the standard model of Euclidean plane geometry [13],[23]. Unlike the Euclidean triangle, the sides of the hyperbolic triangle are uniquely determined by its angles.

4. The Poincaré ball model of hyperbolic geometry

Möbius gyrovector spaces form the setting for the Poincaré ball model of hyperbolic geometry, as demonstrated in [13], [23], just as vector spaces form the setting for the standard model of Euclidean geometry. Thus, the unique geodesic passing through the points $\mathbf{a}, \mathbf{b} \in \mathbb{B}$ in a Möbius gyrovector space $(\mathbb{B}, \oplus_M, \otimes)$ is given by the equation

$$(11) \quad \mathbf{a} \oplus_M (\ominus_M \mathbf{a} \oplus_M \mathbf{b}) \otimes t$$

with the real parameter $t \in \mathbb{R}$. It passes through the point \mathbf{a} at “time” $t = 0$ and, owing to the left cancellation law of gyrogroup theory, it passes through the

point \mathbf{b} at “time” $t = 1$. Several geodesics in the Poincaré disc model, generated by (11), are shown in Figures 1, 2 and 4.




Figure 2: The Hyperbolic Pythagorean Theorem in the Poincaré disc model of hyperbolic geometry that is, equivalently, in the Möbius gyrovector plane $(\mathbb{B}, \oplus, \otimes)$, $\oplus = \oplus_M$ [22],[23].

The hyperbolic midpoint \mathbf{m}_{ab}^P of \mathbf{a} and \mathbf{b} in the Poincaré model is obtained from (11) by selecting $t = 1/2$. It is the midpoint in the sense that $d(\mathbf{a}, \mathbf{m}_{ab}^P) = d(\mathbf{b}, \mathbf{m}_{ab}^P)$ where $d(\mathbf{a}, \mathbf{b})$ is the hyperbolic distance function in the Poincaré model, given by the equation $d(\mathbf{a}, \mathbf{b}) = \|\mathbf{a} \ominus_M \mathbf{b}\|$.

The cosine of the hyperbolic angle generated by two geodesics passing, respectively, through the points \mathbf{a}, \mathbf{b} and \mathbf{a}, \mathbf{c} in the Möbius gyrovector space $(\mathbb{B}, \oplus_M, \otimes)$, Figure 1, is given by the equation

$$(12) \quad \cos \alpha = \frac{\ominus \mathbf{a} \oplus_M \mathbf{b}}{\|\ominus \mathbf{a} \oplus_M \mathbf{b}\|} \cdot \frac{\ominus \mathbf{a} \oplus_M \mathbf{c}}{\|\ominus \mathbf{a} \oplus_M \mathbf{c}\|}$$

in full analogy with its Euclidean counterpart.

Three geodesic segments that form a triangle, and the triangle angles in the Möbius gyrovector plane are shown in Figure 1. The hyperbolic Pythagorean

theorem in the Möbius gyrovector plane is shown in Figure 2. It shares visual analogies with its Euclidean counterpart, demonstrating the union of hyperbolic and Euclidean geometry [25]. The task we face in this article is to determine the centroid of the hyperbolic triangle in Figure 1.

5. Einstein gyrovector spaces

Definition 6 (Einstein addition). Let \mathbb{V} be a real inner product space, and let $\mathbb{B} = \{\mathbf{v} \in \mathbb{V} : \|\mathbf{v}\| < 1\}$ be the open unit ball of \mathbb{V} . Einstein addition $\oplus_{\mathbb{E}}$ in the ball \mathbb{B} is a binary operation in \mathbb{B} given by the equation ([5], [6], [7], [19], [23], [3], [26])

$$(13) \quad \mathbf{u} \oplus \mathbf{v} = \frac{1}{1 + \mathbf{u} \cdot \mathbf{v}} \left\{ \mathbf{u} + \frac{1}{\gamma_{\mathbf{u}}} \mathbf{v} + \frac{\gamma_{\mathbf{u}}}{1 + \gamma_{\mathbf{u}}} (\mathbf{u} \cdot \mathbf{v}) \mathbf{u} \right\}$$

where the vacuum speed of light is normalized to $c = 1$, where \cdot and $\|\cdot\|$ are the inner product and norm that the ball \mathbb{B} inherits from its space \mathbb{V} , and where $\gamma_{\mathbf{v}}$ is the Lorentz factor of \mathbf{v} ,

$$(14) \quad \gamma_{\mathbf{v}} = \frac{1}{\sqrt{1 - \|\mathbf{v}\|^2}}.$$

The groupoid $(\mathbb{B}, \oplus_{\mathbb{E}})$ is a gyrocommutative gyrogroup, as demonstrated in [23], called an Einstein gyrogroup. Furthermore, it admits scalar multiplication \otimes , turning itself into an Einstein gyrovector space $(\mathbb{B}, \oplus_{\mathbb{E}}, \otimes)$.

Definition 7 (Einstein scalar multiplication). Let $(\mathbb{B}, \oplus_{\mathbb{E}})$ be an Einstein gyrogroup. The Einstein scalar multiplication $r \otimes \mathbf{v} = \mathbf{v} \otimes r$ in \mathbb{B} is given by the equation

$$(15) \quad \begin{aligned} r \otimes \mathbf{v} &= \frac{(1 + \|\mathbf{v}\|)^r - (1 - \|\mathbf{v}\|)^r}{(1 + \|\mathbf{v}\|)^r + (1 - \|\mathbf{v}\|)^r} \frac{\mathbf{v}}{\|\mathbf{v}\|} \\ &= \tanh(r \tanh^{-1} \|\mathbf{v}\|) \frac{\mathbf{v}}{\|\mathbf{v}\|} \end{aligned}$$

where $r \in \mathbb{R}$, $\mathbf{v} \in \mathbb{B}$, $\mathbf{v} \neq \mathbf{0}$; and $r \otimes \mathbf{0} = \mathbf{0}$.

Useful identities that relate the Einstein scalar multiplication to the Lorentz factor (14) are [23, Chapter 3]

$$(16) \quad \gamma_{r \otimes \mathbf{v}} = \frac{1}{2} \gamma_{\mathbf{v}}^r \{ (1 + \|\mathbf{v}\|)^r + (1 - \|\mathbf{v}\|)^r \}$$

and

$$(17) \quad \gamma_{r \otimes \mathbf{v}} r \otimes \mathbf{v} = \frac{1}{2} \gamma_{\mathbf{v}}^r \{ (1 + \|\mathbf{v}\|)^r - (1 - \|\mathbf{v}\|)^r \} \frac{\mathbf{v}}{\|\mathbf{v}\|}$$

for $\mathbf{v} \neq \mathbf{0}$, of which the special case of $r = 2$ is of particular interest in this article,

$$(18) \quad \gamma_{2 \otimes \mathbf{v}} = 2\gamma_{\mathbf{v}}^2 - 1$$

and

$$(19) \quad \gamma_{2 \otimes \mathbf{v}} 2 \otimes \mathbf{v} = 2\gamma_{\mathbf{v}}^2 \mathbf{v}.$$

Interestingly, the scalar multiplication that Möbius and Einstein addition admit coincide. This is compatible with the fact that for parallel vectors in the ball, Möbius addition and Einstein addition coincide as well.

6. The Beltrami ball model of hyperbolic geometry

Einstein gyrovector spaces form the setting for the Beltrami ball model of hyperbolic geometry just as vector spaces form the setting for the standard model of Euclidean geometry [23]. Thus, the unique geodesic passing through the points $\mathbf{a}, \mathbf{b} \in \mathbb{B}$ in an Einstein gyrovector space $(\mathbb{B}, \oplus_E, \otimes)$ is given by the equation

$$(20) \quad \mathbf{a} \oplus_E (\ominus_E \mathbf{a} \oplus_M \mathbf{b}) \otimes t$$

with the real parameter $t \in \mathbb{R}$. It passes through the point \mathbf{a} at “time” $t = 0$ and, owing to the left cancellation law of gyrogroup theory, it passes through the point \mathbf{b} at “time” $t = 1$. Several geodesics in the Beltrami disc model, generated by (20), are shown in Figure 3.

The hyperbolic midpoint $\mathbf{m}_{\mathbf{ab}}^B$ of \mathbf{a} and \mathbf{b} in the Beltrami model is obtained from (20) by selecting $t = 1/2$. It is the midpoint in the sense that $d(\mathbf{a}, \mathbf{m}_{\mathbf{ab}}^B) = d(\mathbf{b}, \mathbf{m}_{\mathbf{ab}}^B)$ where $d(\mathbf{a}, \mathbf{b})$ is the hyperbolic distance function in the Beltrami model, given by the equation $d(\mathbf{a}, \mathbf{b}) = \|\mathbf{a} \ominus_E \mathbf{b}\|$.

Interestingly, the hyperbolic midpoint $\mathbf{m}_{\mathbf{ab}}^B$ can be written in terms of ordinary, rather than Einstein, vector addition as

$$(21) \quad \mathbf{m}_{\mathbf{uv}}^B = \mathbf{a} \oplus_E (\ominus_E \mathbf{a} \oplus_M \mathbf{b}) \otimes \frac{1}{2} = \frac{\gamma_{\mathbf{u}} \mathbf{u} + \gamma_{\mathbf{v}} \mathbf{v}}{\gamma_{\mathbf{u}} + \gamma_{\mathbf{v}}}.$$

The derivation of (21) follows from [23, Equation 3.41] and [23, Equation 1.40].

The hyperbolic midpoint $\mathbf{m}_{\mathbf{uv}}^B$, (21), in the Beltrami ball model of hyperbolic geometry is interesting. It can be interpreted, Figure 3, as the Newtonian velocity of the center of momentum of two objects with relativistically corrected masses $m\gamma_{\mathbf{u}}$ and $m\gamma_{\mathbf{v}}$, that move respectively with velocities \mathbf{u} and \mathbf{v} relative to some inertial frame,

$$(22) \quad \mathbf{m}_{\mathbf{uv}}^B = \frac{m\gamma_{\mathbf{u}} \mathbf{u} + m\gamma_{\mathbf{v}} \mathbf{v}}{m\gamma_{\mathbf{u}} + m\gamma_{\mathbf{v}}} = \frac{\gamma_{\mathbf{u}} \mathbf{u} + \gamma_{\mathbf{v}} \mathbf{v}}{\gamma_{\mathbf{u}} + \gamma_{\mathbf{v}}}.$$




Figure 3: Hyperbolic midpoints, medians and a centroid of a triangle and its sides in the Beltrami model and its underlying Einstein gyrovector space. They possess a relativistic mechanical interpretation, analogous to the classical mechanical interpretation of their Euclidean counterparts in [11],[26].

The two masses $m\gamma_{\mathbf{u}}$ and $m\gamma_{\mathbf{v}}$ in (22), shown in Figure 3, are just the common relativistic masses of two objects with equal rest masses m and respective relative velocities \mathbf{u} and \mathbf{v} .

Having identified the hyperbolic midpoint $\mathbf{m}_{\mathbf{u}\mathbf{v}}^B$, (21), with the Newtonian velocity of the center of momentum of the two relativistically corrected masses, $m\gamma_{\mathbf{u}}$ and $m\gamma_{\mathbf{v}}$, (22), it is clear that the Newtonian velocity $\mathbf{m}_{\mathbf{u}\mathbf{v}\mathbf{w}}^B$ of the center of momentum of the three relativistically corrected masses $m\gamma_{\mathbf{u}}$, $m\gamma_{\mathbf{v}}$, and $m\gamma_{\mathbf{w}}$,

$$(23) \quad \mathbf{m}_{\mathbf{u}\mathbf{v}\mathbf{w}}^B = \frac{m\gamma_{\mathbf{u}}\mathbf{u} + m\gamma_{\mathbf{v}}\mathbf{v} + m\gamma_{\mathbf{w}}\mathbf{w}}{m\gamma_{\mathbf{u}} + m\gamma_{\mathbf{v}} + m\gamma_{\mathbf{w}}} = \frac{\gamma_{\mathbf{u}}\mathbf{u} + \gamma_{\mathbf{v}}\mathbf{v} + \gamma_{\mathbf{w}}\mathbf{w}}{\gamma_{\mathbf{u}} + \gamma_{\mathbf{v}} + \gamma_{\mathbf{w}}}$$

lies on the median connecting the point $\mathbf{m}_{\mathbf{u}\mathbf{v}}^B$ to the point \mathbf{w} of triangle $\Delta\mathbf{u}\mathbf{v}\mathbf{w}$ of Figure 3. By symmetry considerations, the Newtonian velocity $\mathbf{m}_{\mathbf{u}\mathbf{v}\mathbf{w}}^B$ of the center of momentum of the three relativistically corrected masses $m\gamma_{\mathbf{u}}$, $m\gamma_{\mathbf{v}}$, and $m\gamma_{\mathbf{w}}$ lies on the other two hyperbolic medians of triangle $\Delta\mathbf{u}\mathbf{v}\mathbf{w}$ as well. Hence, the point $\mathbf{m}_{\mathbf{u}\mathbf{v}\mathbf{w}}^B$ coincides with the centroid of the hyperbolic triangle $\Delta\mathbf{u}\mathbf{v}\mathbf{w}$, as

shown in Figure 3. Hence, by elementary linear algebra, the hyperbolic centroid $C_{\mathbf{uvw}}^B$ of the hyperbolic triangle $\Delta\mathbf{uvw}$ in the Beltrami ball model of hyperbolic geometry, Figure 3, is equal to the velocity $\mathbf{m}_{\mathbf{uvw}}^B$ in (23), that is,

$$(24) \quad C_{\mathbf{uvw}}^B = \frac{\gamma_{\mathbf{u}}\mathbf{u} + \gamma_{\mathbf{v}}\mathbf{v} + \gamma_{\mathbf{w}}\mathbf{w}}{\gamma_{\mathbf{u}} + \gamma_{\mathbf{v}} + \gamma_{\mathbf{w}}}.$$

Formalizing our result in (24), and noting that an Einstein gyrovector space underlies the Beltrami ball model of hyperbolic geometry, we have the following

Theorem 8. *Let $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{B}$ be any three non-gyrocollinear points of a Beltrami ball model, \mathbb{B} , of hyperbolic geometry, where \mathbb{B} is the ball of a real inner product space \mathbb{V} . The centroid $C_{\mathbf{abc}}^B$ of the hyperbolic triangle $\Delta\mathbf{abc}$ in \mathbb{B} is given by the equation*

$$(25) \quad C_{\mathbf{abc}}^B = \frac{\gamma_{\mathbf{a}}\mathbf{a} + \gamma_{\mathbf{b}}\mathbf{b} + \gamma_{\mathbf{c}}\mathbf{c}}{\gamma_{\mathbf{a}} + \gamma_{\mathbf{b}} + \gamma_{\mathbf{c}}}.$$

7. Triangle centroids in the Poincaré ball model of hyperbolic geometry

We wish, in this section, to translate the expression (25) of the centroid of a hyperbolic triangle in an Einstein gyrovector space and its associated Beltrami ball model of hyperbolic geometry into an expression describing the centroid of a hyperbolic triangle in a Möbius gyrovector space and its associated Poincaré ball model of hyperbolic geometry.

Let $G_e = (\mathbb{B}, \oplus_E, \otimes)$ and $G_m = (\mathbb{B}, \oplus_M, \otimes)$ be, respectively, the Einstein and the Möbius gyrovector spaces of the ball \mathbb{B} of a real inner product space \mathbb{V} . They are gyrovector space isomorphic, with the isomorphism and its inverse isomorphism from G_m into G_e given by the equations [23]

$$(26) \quad \begin{aligned} \mathbf{v}_e &= 2 \otimes \mathbf{v}_m, \\ \mathbf{v}_m &= \frac{1}{2} \otimes \mathbf{v}_e, \end{aligned}$$

$\mathbf{v}_e \in G_e, \mathbf{v}_m \in G_m$. Accordingly, the gyrogroup operations \oplus_E and \oplus_M in G_e and G_m are related to each other by the equation

$$(27) \quad \begin{aligned} \mathbf{u}_m \oplus_M \mathbf{v}_m &= \frac{1}{2} \otimes (2 \otimes \mathbf{u}_m \oplus_E 2 \otimes \mathbf{v}_m), \\ \mathbf{u}_e \oplus_E \mathbf{v}_e &= 2 \otimes (\frac{1}{2} \otimes \mathbf{u}_e \oplus_M \frac{1}{2} \otimes \mathbf{v}_e). \end{aligned}$$

Following (26) and (18) we have

$$(28) \quad \gamma_{\mathbf{v}_e} = \gamma_{2 \otimes \mathbf{v}_m} = 2\gamma_{\mathbf{v}_m}^2 - 1.$$

Similarly, following (26) and (19) we have

$$(29) \quad \gamma_{\mathbf{v}_e} \mathbf{v}_e = \gamma_{2 \otimes \mathbf{v}_m} 2 \otimes \mathbf{v}_m = 2\gamma_{\mathbf{v}_m}^2 \mathbf{v}_m .$$

Hence, by (21), (28) and (29) we have

$$(30) \quad \begin{aligned} \mathbf{m}_{\mathbf{u}_e \mathbf{v}_e}^B &= \frac{\gamma_{\mathbf{u}_e} \mathbf{u}_e + \gamma_{\mathbf{v}_e} \mathbf{v}_e}{\gamma_{\mathbf{u}_e} + \gamma_{\mathbf{v}_e}} \\ &= \frac{2\gamma_{\mathbf{u}_m}^2 \mathbf{u}_m + 2\gamma_{\mathbf{v}_m}^2 \mathbf{v}_m}{(2\gamma_{\mathbf{u}_m}^2 - 1) + (2\gamma_{\mathbf{v}_m}^2 - 1)} \\ &= \frac{\gamma_{\mathbf{u}_m}^2 \mathbf{u}_m + \gamma_{\mathbf{v}_m}^2 \mathbf{v}_m}{\gamma_{\mathbf{u}_m}^2 + \gamma_{\mathbf{v}_m}^2 - 1} \end{aligned}$$

so that

$$(31) \quad \begin{aligned} \mathbf{m}_{\mathbf{u}_m \mathbf{v}_m}^P &= \frac{1}{2} \otimes \mathbf{m}_{\mathbf{u}_e \mathbf{v}_e}^B \\ &= \frac{1}{2} \otimes \frac{\gamma_{\mathbf{u}_m}^2 \mathbf{u}_m + \gamma_{\mathbf{v}_m}^2 \mathbf{v}_m}{\gamma_{\mathbf{u}_m}^2 + \gamma_{\mathbf{v}_m}^2 - 1} . \end{aligned}$$

We have thus obtained in (31) the following

Theorem 9. *Let $\mathbf{a}, \mathbf{b} \in \mathbb{B}$ be any two points of a Poincaré ball model, \mathbb{B} , of hyperbolic geometry, where \mathbb{B} is the ball of a real inner product space \mathbb{V} . The midpoint $\mathbf{m}_{\mathbf{ab}}^P$ of the hyperbolic segment \mathbf{ab} joining the points \mathbf{a} and \mathbf{b} in \mathbb{B} is given by the equation*

$$(32) \quad \mathbf{m}_{\mathbf{ab}}^P = \frac{1}{2} \otimes \frac{\gamma_{\mathbf{a}}^2 \mathbf{a} + \gamma_{\mathbf{b}}^2 \mathbf{b}}{\gamma_{\mathbf{a}}^2 + \gamma_{\mathbf{b}}^2 - 1} .$$

In the same way we obtained (30) and (31) it follows, by (25), (28) and (29), that

$$(33) \quad \begin{aligned} C_{\mathbf{u}_e \mathbf{v}_e \mathbf{w}_e}^B &= \frac{\gamma_{\mathbf{u}_e} \mathbf{u}_e + \gamma_{\mathbf{v}_e} \mathbf{v}_e + \gamma_{\mathbf{w}_e} \mathbf{w}_e}{\gamma_{\mathbf{u}_e} + \gamma_{\mathbf{v}_e} + \gamma_{\mathbf{w}_e}} \\ &= \frac{2\gamma_{\mathbf{u}_m}^2 \mathbf{u}_m + 2\gamma_{\mathbf{v}_m}^2 \mathbf{v}_m + 2\gamma_{\mathbf{w}_m}^2 \mathbf{w}_m}{(2\gamma_{\mathbf{u}_m}^2 - 1) + (2\gamma_{\mathbf{v}_m}^2 - 1) + (2\gamma_{\mathbf{w}_m}^2 - 1)} \\ &= \frac{\gamma_{\mathbf{u}_m}^2 \mathbf{u}_m + \gamma_{\mathbf{v}_m}^2 \mathbf{v}_m + \gamma_{\mathbf{w}_m}^2 \mathbf{w}_m}{\gamma_{\mathbf{u}_m}^2 + \gamma_{\mathbf{v}_m}^2 + \gamma_{\mathbf{w}_m}^2 - \frac{3}{2}} \end{aligned}$$




Figure 4: A triangle Δuvw in the Poincaré disc model of hyperbolic geometry is shown with the midpoints \mathbf{m}_{uv}^P , \mathbf{m}_{uw}^P and \mathbf{m}_{vw}^P of its sides, and its medians, and centroid \mathbf{C}_{uvw}^P .

so that

$$\begin{aligned}
 (34) \quad \mathbf{C}_{\mathbf{u}_m \mathbf{v}_m \mathbf{w}_m}^P &= \frac{1}{2} \otimes \mathbf{C}_{\mathbf{u}_e \mathbf{v}_e \mathbf{w}_e}^B \\
 &= \frac{1}{2} \otimes \frac{\gamma_{\mathbf{u}_m}^2 \mathbf{u}_m + \gamma_{\mathbf{v}_m}^2 \mathbf{v}_m + \gamma_{\mathbf{w}_m}^2 \mathbf{w}_m}{\gamma_{\mathbf{u}_m}^2 + \gamma_{\mathbf{v}_m}^2 + \gamma_{\mathbf{w}_m}^2 - \frac{3}{2}}
 \end{aligned}$$

for any $\mathbf{u}_m, \mathbf{v}_m, \mathbf{w}_m \in G_m$.

We have thus obtained in (34) the following

Theorem 10. *Let $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{B}$ be any three non-gyrocollinear points of a Poincaré ball model, \mathbb{B} , of hyperbolic geometry, where \mathbb{B} is the ball of a real inner product space \mathbb{V} . The centroid $\mathbf{C}_{\mathbf{abc}}^P$ of the hyperbolic triangle $\Delta \mathbf{abc}$ in \mathbb{B} is given by the equation*

$$(35) \quad \mathbf{C}_{\mathbf{abc}}^P = \frac{1}{2} \otimes \frac{\gamma_{\mathbf{a}}^2 \mathbf{a} + \gamma_{\mathbf{b}}^2 \mathbf{b} + \gamma_{\mathbf{c}}^2 \mathbf{c}}{\gamma_{\mathbf{a}}^2 + \gamma_{\mathbf{b}}^2 + \gamma_{\mathbf{c}}^2 - \frac{3}{2}}.$$

The hyperbolic triangle centroid was also studied by O. Bottema [2]. The centroid of the hyperbolic triangle Δabc in the Poincaré disc model, as determined by (35), is shown in Figure 4.

REFERENCES

- [1] Ahlfors L.V., *Möbius transformations in several dimensions*, Univ. of Minnesota School of Mathematics, Minneapolis, Minnesota, 1981.
- [2] Bottema O., *On the medians of a triangle in hyperbolic geometry*, *Canad. J. Math.* **10** (1958), 502–506.
- [3] Chen J.-L., Ungar A.A., *The Bloch gyrovector*, *Found. Phys.* **32** (2002), no. 4, 531–565.
- [4] Csörgő P., *H-connected transversals to abelian subgroups*, preprint.
- [5] Einstein A., *Zur Elektrodynamik Bewegter Körper [On the electrodynamics of moving bodies]*, *Ann. Physik (Leipzig)* **17** (1905), 891–921.
- [6] Einstein A., *Einstein's Miraculous Years: Five Papers that Changed the Face of Physics*, Princeton, Princeton, NJ, 1998. Edited and introduced by John Stachel. Includes bibliographical references. Einstein's dissertation on the determination of molecular dimensions – Einstein on Brownian motion – Einstein on the theory of relativity – Einstein's early work on the quantum hypothesis. A new English translation of Einstein's 1905 paper on pp. 123–160.
- [7] Fock V., *The Theory of Space, Time and Gravitation*, The Macmillan Co., New York, 1964. Second revised edition. Translated from the Russian by N. Kemmer. A Pergamon Press Book.
- [8] Foguel T., Ungar A.A., *Involutory decomposition of groups into twisted subgroups and subgroups*, *J. Group Theory* **3** (1) (2000), 27–46.
- [9] Foguel T., Ungar A.A., *Gyrogroups and the decomposition of groups into twisted subgroups and subgroups*, *Pacific J. Math.* **197:1** (2001), 1–11.
- [10] Foguel T., Kinyon M.K., Phillip J.D., *On twisted subgroups and Bol loops of odd order*, *Rocky Mountain J. Math.*, in print.
- [11] Hausner M., *A Vector Space Approach to Geometry*, Dover Publications Inc., Mineola, NY, 1998. Reprint of the 1965 original.
- [12] Kiechle H., *Theory of K-loops*, Springer-Verlag, Berlin, 2002.
- [13] Kinyon M.K., Ungar A.A., *The gyro-structure of the complex unit disk*, *Math. Mag.* **73:4** (2000), 273–284.
- [14] Kuznetsov E., *Transversals in loops*, preprint.
- [15] Lévay P., *The geometry of entanglement: metrics, connections and the geometric phase*, [arXiv:quant-ph/0306115](https://arxiv.org/abs/quant-ph/0306115) v1 2003.
- [16] Lévay P., *Mixed state geometric phase from Thomas rotations*, [arXiv:quant-ph/0312023](https://arxiv.org/abs/quant-ph/0312023) v1 2003.
- [17] Ratcliffe J.G., *Foundations of hyperbolic manifolds*, vol. 149 of *Graduate Texts in Mathematics*, Springer-Verlag, New York, 1994.
- [18] Sabinin L.V., Sabinina L.L., Sbitneva L.V., *On the notion of gyrogroup*, *Aequationes Math.* **56** (1-2) (1998), 11–17.
- [19] Sexl R.U., Urbantke H.K., *Relativity, Groups, Particles*, Springer-Verlag, Vienna, 2001. Special relativity and relativistic symmetry in field and particle physics; revised and translated from the third German (1992) edition by Urbantke.
- [20] Ungar A.A., *The relativistic noncommutative nonassociative group of velocities and the Thomas rotation*, *Resultate Math.* **16** (1-2) (1989), 168–179. The term “K-loop” is coined here.

- [21] Ungar A.A., *Extension of the unit disk gyrogroup into the unit ball of any real inner product space*, J. Math. Anal. Appl. **202:3** (1996), 1040–1057.
- [22] Ungar A.A., *The hyperbolic Pythagorean theorem in the Poincaré disc model of hyperbolic geometry*, Amer. Math. Monthly **106:8** (1999), 759–763.
- [23] Ungar A.A., *Beyond the Einstein addition law and its gyroscopic Thomas precession*, volume 117 of *Fundamental Theories of Physics*, Kluwer Academic Publishers Group, Dordrecht, 2001. The theory of gyrogroups and gyrovector spaces.
- [24] Ungar A.A., *The hyperbolic geometric structure of the density matrix for mixed state qubits*, Found. Phys. **32** (2002), no. 11, 1671–1699.
- [25] Ungar A.A., *On the unification of hyperbolic and Euclidean geometry*, Complex Variables Theory Appl. **49** (2004), 197–213.
- [26] Ungar A.A., *Einstein's special relativity: Unleashing the power of its hyperbolic geometry*, preprint, 2004.

DEPARTMENT OF MATHEMATICS, NORTH DAKOTA STATE UNIVERSITY, FARGO, ND 58105,
USA

E-mail: abraham.ungar@ndsu.nodak.edu

(Received September 15, 2003, revised January 13, 2004)