

Yan-Kui Song

Embedding into discretely absolutely star-Lindelöf spaces

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 48 (2007), No. 2, 303--309

Persistent URL: <http://dml.cz/dmlcz/119660>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2007

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Embedding into discretely absolutely star-Lindelöf spaces

YAN-KUI SONG

*Abstract.* A space  $X$  is *discretely absolutely star-Lindelöf* if for every open cover  $\mathcal{U}$  of  $X$  and every dense subset  $D$  of  $X$ , there exists a countable subset  $F$  of  $D$  such that  $F$  is discrete closed in  $X$  and  $\text{St}(F, \mathcal{U}) = X$ , where  $\text{St}(F, \mathcal{U}) = \bigcup\{U \in \mathcal{U} : U \cap F \neq \emptyset\}$ . We show that every Hausdorff star-Lindelöf space can be represented in a Hausdorff discretely absolutely star-Lindelöf space as a closed subspace.

*Keywords:* normal, star-Lindelöf, centered-Lindelöf

*Classification:* 54D20, 54G20

### 1. Introduction

By a space, we mean a topological space. A space  $X$  is *absolutely star-Lindelöf* (see [1]) (*discretely absolutely star-Lindelöf*) (see [10], [11]) if for every open cover  $\mathcal{U}$  of  $X$  and every dense subset  $D$  of  $X$ , there exists a countable subset  $F$  of  $D$  such that  $\text{St}(F, \mathcal{U}) = X$  ( $F$  is discrete and closed in  $X$  and  $\text{St}(F, \mathcal{U}) = X$ , respectively), where  $\text{St}(F, \mathcal{U}) = \bigcup\{U \in \mathcal{U} : U \cap F \neq \emptyset\}$ .

A space  $X$  is *star-Lindelöf* (see [4], [7] under different names) (*discretely star-Lindelöf*) (see [9], [15]) if for every open cover  $\mathcal{U}$  of  $X$ , there exists a countable subset (a countable discrete closed subset, respectively)  $F$  of  $X$  such that  $\text{St}(F, \mathcal{U}) = X$ . It is clear that every separable space and every discretely star-Lindelöf space are star-Lindelöf as well as every space of countable extent (in particular, every countably compact space or every Lindelöf space).

A family of subsets is *centered* (*linked*) provided every finite subfamily (every two elements, respectively) has nonempty intersection and a family is called  $\sigma$ -*centered* ( $\sigma$ -*linked*) if it is the union of countably many centered subfamilies (linked subfamilies, respectively). A space  $X$  is *centered-Lindelöf* (*linked-Lindelöf*) (see [2], [3]) if every open cover  $\mathcal{U}$  of  $X$  has a  $\sigma$ -centered ( $\sigma$ -linked) subcover.

From the above definitions, it is not difficult to see that every discretely absolutely star-Lindelöf space is absolutely star-Lindelöf, every discretely absolutely star-Lindelöf space is discretely star-Lindelöf, every absolutely star-Lindelöf space is star-Lindelöf, every star-Lindelöf space is centered-Lindelöf, and every centered-Lindelöf space is linked-Lindelöf.

---

The author acknowledges support from the NSF of China Grant 10571081.

Bonanzinga and Matveev [2] proved that every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented a closed subspace in a Hausdorff (regular, Tychonoff, respectively) star-Lindelöf space. They asked if every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented a closed  $G_\delta$ -subspace in a Hausdorff (regular, Tychonoff, respectively) star-Lindelöf space. The author [10] gave a positive answer to their question. The author [11] showed that every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented a closed  $G_\delta$ -subspace in a Hausdorff (regular, Tychonoff, respectively) absolutely star-Lindelöf space. The author [12] showed that every separable Hausdorff (regular, Tychonoff, normal) star-Lindelöf space can be represented in a Hausdorff (regular, Tychonoff, normal, respectively) discretely absolutely star-Lindelöf space as a closed  $G_\delta$ -subspace. Thus, it is natural for us to consider the following question:

**Question.** *Is it true that every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented a closed subspace in a Hausdorff (regular, Tychonoff, respectively) discretely absolutely star-Lindelöf space? And can it be embedded as a closed  $G_\delta$ -subspace?*

The purpose of this note is to give a construction showing that every Hausdorff linked-Lindelöf space can be represented in a Hausdorff discretely absolutely star-Lindelöf space as a closed subspace, which gives a positive answer to the question in the class of Hausdorff spaces.

Throughout this paper, the cardinality of a set  $A$  is denoted by  $|A|$ . Let  $\omega$  denote the first infinite cardinal. For a cardinal  $\kappa$ , let  $\kappa^+$  be the smallest cardinal greater than  $\kappa$ . As usual, a cardinal is the initial ordinal and an ordinal is the set of smaller ordinals. When viewed as a space, every cardinal has the usual order topology. For each pair of ordinals  $\alpha, \beta$  with  $\alpha < \beta$ , we write  $[\alpha, \beta] = \{\gamma : \alpha \leq \gamma \leq \beta\}$  and  $(\alpha, \beta) = \{\gamma : \alpha < \gamma < \beta\}$ . Other terms and symbols that we do not define will be used as in [5].

## 2. Embedding into discretely absolutely star-Lindelöf spaces

First, we show that every Hausdorff star-Lindelöf space can be represented in a Hausdorff discretely absolutely star-Lindelöf space as a closed subspace.

Recall the definition of the Alexandorff duplicate  $A(X)$  of a space  $X$ . The underlying set of  $A(X)$  is  $X \times \{0, 1\}$ ; each point of  $X \times \{1\}$  is isolated and a basic neighborhood of a point  $\langle x, 0 \rangle \in X \times \{0\}$  is of the form  $(U \times \{0\}) \cup ((U \times \{1\}) \setminus \{\langle x, 1 \rangle\})$ , where  $U$  is a neighborhood of  $x$  in  $X$ . It is well-known that  $A(X)$  is Hausdorff (regular, Tychonoff, normal) iff  $X$  is,  $A(X)$  is compact iff  $X$  is and  $A(X)$  is Lindelöf iff  $X$  is.

Recall from [6] that a space  $X$  is *absolutely countably compact* ( $=$  acc) if for every open cover  $\mathcal{U}$  of  $X$  and every dense subset  $D$  of  $X$ , there exists a finite subset  $F$  of  $D$  such that  $\text{St}(F, \mathcal{U}) = X$ . It is not difficult to show that every

Hausdorff absolutely countably compact space is countably compact (see [6]). In our construction, we use the following lemma.

**Lemma 2.1** ([8], [14]). *If  $X$  is countably compact, then  $A(X)$  is acc. Moreover, for any open cover  $\mathcal{U}$  of  $A(X)$ , there exists a finite subset  $F$  of  $X \times \{1\}$  such that  $A(X) \setminus \text{St}(F, \mathcal{U}) \subseteq X \times \{0\}$  is a finite subset consisting of isolated points of  $X \times \{0\}$ .*

**Theorem 2.2.** *Every Hausdorff star-Lindelöf space can be represented in a Hausdorff discretely absolutely star-Lindelöf space as a closed subspace.*

PROOF: If  $|X| \leq \omega$ , then  $X$  is separable; the author [12] showed that every separable Hausdorff (regular, Tychonoff, normal) space can be represented in Hausdorff (regular, Tychonoff, normal respectively) discretely absolutely star-Lindelöf space as a closed  $G_\delta$ -subspace.

Let  $X$  be a star-Lindelöf space with  $|X| > \omega$ , let  $T$  be  $X$  with the discrete topology and let

$$Y = T \cup \{\infty\}, \text{ where } \infty \notin T$$

be the one-point Lindelöfication of  $T$ . Pick a cardinal  $\kappa$  with  $\kappa \geq |X|$ . Define

$$S(X, \kappa) = X \cup (Y \times \kappa^+).$$

We topologize  $S(X, \kappa)$  as follows:  $Y \times \kappa^+$  has the usual product topology and is an open subspace of  $S(X, \kappa)$ , and a basic neighborhood of a point  $x$  of  $X$  takes the form

$$G(U, \alpha) = U \cup (U \times (\alpha, \kappa^+)),$$

where  $U$  is a neighborhood of  $x$  in  $X$  and  $\alpha < \kappa^+$ . Then, it is easy to see that  $X$  is a closed subset of  $S(X, \kappa)$  and  $S(X, \kappa)$  is Hausdorff if  $X$  is Hausdorff.

Let

$$\mathcal{R}(X) = A(S(X, \kappa)) \setminus (X \times \{1\}).$$

Then  $\mathcal{R}(X)$  is Hausdorff if  $X$  is Hausdorff.

We show that  $\mathcal{R}(X)$  is discretely absolutely star-Lindelöf. To this end, let  $\mathcal{U}$  be an open cover of  $\mathcal{R}(X)$ . Without loss of generality, we assume that  $\mathcal{U}$  consists of basic open sets of  $\mathcal{R}(X)$ . Let  $S$  be the set of all isolated points of  $\kappa^+$  and let

$$D_1 = ((T \times S) \times \{0\}) \cup ((T \times \kappa^+) \times \{1\}) \text{ and } D_2 = (\{\infty\} \times \kappa^+) \times \{1\}.$$

Set  $D = D_1 \cup D_2$ . Then, every element of  $D$  is isolated in  $\mathcal{R}(X)$ , and so every dense subset of  $\mathcal{R}(X)$  contains  $D$ . Thus, it is sufficient to show that there exists a countable subset  $F$  of  $D$  such that  $F$  is discrete closed in  $\mathcal{R}(X)$  and  $\text{St}(F, \mathcal{U}) = \mathcal{R}(X)$ .

For each  $x \in X$ , there exists a  $U_x \in \mathcal{U}$  such that  $\langle x, 0 \rangle \in U_x$ . Hence there exist  $\alpha_x < \kappa^+$  and an open neighborhood  $V_x$  of  $x$  in  $X$  such that

$$(V_x \times \{0\}) \cup A(V_x \times (\alpha_x, \kappa^+)) \subseteq U_x.$$

If we put  $\mathcal{V} = \{V_x : x \in X\}$ , then  $\mathcal{V}$  is an open cover of  $X$ , hence there exists a countable subset  $F'_1$  of  $X$  such that  $X = \text{St}(F'_1, \mathcal{V})$ , since  $X$  is star-Lindelöf. We pick  $\alpha_0 > \sup\{\alpha_x : x \in X\}$ . Let

$$\begin{aligned} X_1 &= (X \times \{0\}) \cup A(T \times [\alpha_0, \kappa^+)); \\ X_2 &= A(T \times [0, \alpha_0]) \quad \text{and} \quad X_3 = A(\{\infty\} \times \kappa^+). \end{aligned}$$

Then,

$$X = X_1 \cup X_2 \cup X_3.$$

Let

$$F_1 = (F'_1 \times \{0\}) \times \{1\}.$$

Then,  $F_1$  is a countable subset of  $D_1$  and

$$X_1 \subseteq \text{St}(F_1, \mathcal{U}),$$

since  $U_x \cap F_1 \neq \emptyset$  for each  $x \in X$ . Since  $F_1 \subseteq D_1$  and  $F_1$  is countable,  $F_1$  is closed in  $\mathcal{R}(X)$  by the construction of the topology of  $\mathcal{R}(X)$ .

On the other hand, since  $Y$  is Lindelöf and  $[0, \alpha_0]$  is compact,  $Y \times [0, \alpha_0]$  is Lindelöf, hence  $X_1 = A(Y \times [0, \alpha_0])$  is Lindelöf. For each  $\alpha \leq \alpha_0$  there exists a  $U_\alpha \in \mathcal{U}$  such that  $\langle \langle \infty, \alpha \rangle, 0 \rangle \in U_\alpha$ , hence there exists an open neighborhood  $V_\alpha$  of  $\alpha$  in  $\kappa^+$  and an open neighborhood  $V'_\alpha$  of  $\infty$  in  $Y$  such that

$$A(V'_\alpha \times V_\alpha) \setminus (\langle \langle \infty, \alpha \rangle, 1 \rangle) \subseteq U_\alpha.$$

Let  $\mathcal{V}' = \{V_\alpha : \alpha \leq \alpha_0\}$ . Then,  $\mathcal{V}'$  is an open cover of  $[0, \alpha_0]$ . Hence, there exists a finite subcover  $V_{\alpha_1}, V_{\alpha_2}, \dots, V_{\alpha_n}$ , since  $[0, \alpha_0]$  is compact. Let

$$T_1 = \bigcup \{T \setminus V'_{\alpha_i} : i \leq n\}.$$

Then,  $T_1$  is a countable subset of  $T$ . For each  $i \leq n$ , we pick  $x_i \in D_1 \cap U_{\alpha_i}$ . Let  $F'_2 = \{x_i : i \leq n\}$ . Then,  $F'_2$  is a finite subset of  $D_1$  and

$$((\{\infty\} \times [0, \alpha_0]) \times \{0\}) \cup A((T \setminus T_1) \times [0, \alpha_0]) \subseteq \text{St}(F'_2, \mathcal{U}).$$

For each  $t \in T_1$ , since  $\{t\} \times [0, \alpha_0]$  is compact,  $A(\{t\} \times [0, \alpha_0])$  is compact as well, hence there exists a finite subset  $F_t$  of  $D_1$  such that

$$A(\{t\} \times [0, \alpha_0]) \subseteq \text{St}(F_t, \mathcal{U}).$$

Let  $F_2'' = \bigcup\{F_t : t \in T_1\}$ . Then,  $F_2''$  is countable, since  $T_1$  is countable. Since  $F_2'' \cap A(\{\alpha\} \times Y)$  is countable for each  $\alpha < \kappa^+$  and  $F_2'' \cap A(\{\kappa^+\} \times \{t\})$  is finite for each  $t \in T$ ,  $F_2''$  is closed in  $\mathcal{R}(X)$  by the construction of the topology of  $\mathcal{R}(X)$ . By the definition of  $F_2''$ , we have

$$A(T_1 \times [0, \alpha_0]) \subseteq \text{St}(F_2'', \mathcal{U}).$$

Then,  $F_2 = F_2' \cup F_2''$  is a countable closed subset of  $D_2$ , since  $F_1'$  and  $F_2''$  are closed in  $\mathcal{R}(X)$ , and

$$X_2 \cup ((\{\infty\} \times [0, \alpha_0]) \times \{0\}) \subseteq \text{St}(F_2, \mathcal{U}).$$

Finally, we show that there exists a finite subset  $F_3$  of  $D$  such that  $X_3 \subseteq \text{St}(F_3, \mathcal{U})$ . Since  $\{\infty\} \times \kappa^+$  is countably compact, then, by Lemma 2.1, there exists a finite subset  $F_3' \subseteq (\{\infty\} \times \kappa^+) \times \{1\}$  such that

$$E = X_3 \setminus \text{St}(F_3', \mathcal{U}) \subseteq (\{\infty\} \times \kappa^+) \times \{0\} \text{ is a finite subset}$$

and each point of  $E$  is an isolated point of  $(\{\infty\} \times \kappa^+) \times \{0\}$ . For each point  $x \in E$ , there exists a  $U_x \in \mathcal{U}$  such that  $x \in U_x$ . For each point  $x \in E$ , pick  $d_x \in D \cap U_x$ . Let  $F_3'' = \{d_x : x \in E\}$ ; then  $F_3''$  is a finite subset of  $D$  and  $E \subseteq \text{St}(F_3'', \mathcal{U})$ . If we put  $F_3 = F_3' \cup F_3''$ , then  $F_3$  is a finite subset of  $D$  and

$$X_3 \subseteq \text{St}(F_3, \mathcal{U}).$$

If we put  $F = F_1 \cup F_2 \cup F_3$ , then  $F$  is a countable subset of  $D$  such that  $\text{St}(F, \mathcal{U}) = \mathcal{R}(X)$ . Since  $F_1$  and  $F_2$  are closed in  $\mathcal{R}(X)$ ,  $F_3$  is finite, and each point of  $F$  is isolated,  $F$  is discrete closed in  $X$ , which completes the proof.  $\square$

Since every discretely absolutely star-Lindelöf space is discretely star-Lindelöf, the next corollary follows from Theorem 2.2.

**Corollary 2.3.** *Every Hausdorff star-Lindelöf space can be represented in a Hausdorff discretely star-Lindelöf space as a closed subspace.*

Since every discretely absolutely star-Lindelöf space is absolutely star-Lindelöf, the next corollary follows from Theorem 2.2.

**Corollary 2.4.** *Every Hausdorff star-Lindelöf space can be represented in a Hausdorff absolutely star-Lindelöf space as a closed subspace.*

Bonanzinga and Matveev [2] proved that every Hausdorff (regular, Tychonoff) linked-Lindelöf space can be represented a closed subspace in Hausdorff (regular, Tychonoff, respectively) star-Lindelöf space. Thus, we have the following corollary.

**Corollary 2.5.** *Every Hausdorff linked-Lindelöf space can be represented in a Hausdorff discretely absolutely star-Lindelöf space as a closed subspace.*

We have the following two propositions on the separation of Theorem 2.2.

**Proposition 2.6.** *If  $X$  is locally countable (i.e., each point of  $X$  has a neighborhood  $U$  with  $|U| \leq \omega$ ) and Tychonoff, then  $S(X, \kappa)$  is Tychonoff (hence,  $\mathcal{R}(X)$  is Tychonoff).*

PROOF: Assume that  $X$  is locally countable, Tychonoff and let  $x \in X$ . Since a locally countable, Tychonoff space is zero-dimensional,  $x$  has a neighborhood base  $\mathcal{U}(x)$  in  $X$  consisting countable, open-closed sets in  $X$ . If  $U \in \mathcal{U}(x)$ , then the set

$$G(U, \alpha) = U \cup (U \times (\alpha, \kappa^+))$$

is open-closed in  $S(X, \kappa)$  for each  $\alpha < \kappa^+$ . Hence,  $x$  has a neighborhood base in  $S(X, \kappa)$  consisting of open-closed sets, which implies  $S(X, \kappa)$  is Tychonoff.  $\square$

**Proposition 2.7.** *If  $X$  is not locally countable, then  $S(X, \kappa)$  is not regular (hence,  $\mathcal{R}(X)$  is not regular).*

PROOF: If  $X$  is not locally countable, then there exists a point  $x \in X$  which has no countable neighborhood in  $X$ . Let  $\mathcal{U}(x)$  be a neighborhood base of  $x$  in  $X$ . If  $U \in \mathcal{U}$  and  $\alpha < \kappa^+$ , then the closure of  $G(U, \alpha)$  in  $S(X, \kappa)$  contains  $\langle \infty, \gamma \rangle$  for each  $\gamma > \alpha$  by construction of  $Y$ . This means that for any  $U, V \in \mathcal{U}(x)$  and any  $\alpha, \beta < \kappa^+$ ,

$$\langle \infty, \gamma \rangle \in \text{cl}_{S(X, \kappa)} G(U, \alpha) \setminus G(V, \beta)$$

for each  $\gamma > \alpha$ . Hence,  $S(X, \kappa)$  is not regular.  $\square$

By Theorem 2.2 and Proposition 2.6, we have the following theorem:

**Theorem 2.8.** *Every locally-countable, star-Lindelöf Tychonoff space can be represented in a discretely absolutely star-Lindelöf Tychonoff space as a closed subspace.*

*Remark 1.* In Theorem 2.2, even if  $X$  is locally-countable normal,  $\mathcal{R}(X)$  need not be normal. Indeed,  $X \times \{0\}$  and  $A(\{\infty\} \times \kappa^+)$  are disjoint closed subsets of  $\mathcal{R}(X)$  that cannot be separated by disjoint open subsets of  $\mathcal{R}(X)$ . Thus, the author does not know if every normal star-Lindelöf space can be represented in a normal discretely absolutely star-Lindelöf space as a closed subspace.

*Remark 2.* From Theorem 2.2, it is not difficult to see that  $X \times \{0\}$  is not a closed  $G_\delta$  subset of  $\mathcal{R}(X)$ . Thus, the author does not know if every Hausdorff (regular, Tychonoff) star-Lindelöf space can be represented in a Hausdorff (regular, Tychonoff, respectively) discretely absolutely star-Lindelöf space as a closed  $G_\delta$ -subspace.

**Acknowledgments.** The author is most grateful to Prof. H. Ohta for his kind help and valuable suggestions. The author is also most grateful to the referee for his kind help and valuable comments.

## REFERENCES

- [1] Bonanzinga M., *Star-Lindelöf and absolutely star-Lindelöf spaces*, Questions Answers Gen. Topology **16** (1998), 79–104.
- [2] Bonanzinga M., Matveev M.V., *Closed subspaces of star-Lindelöf and related spaces*, East-West J. Math. **2** (2000), no. 2, 171–179.
- [3] Bonanzinga M., Matveev M.V., *Products of star-Lindelöf and related spaces*, Houston J. Math. **27** (2001), 45–57.
- [4] van Douwen E.K., Reed G.M., Roscoe A.W., Tree I.J., *Star covering properties*, Topology Appl. **39** (1991), 710–103.
- [5] Engelking R., *General Topology*, revised and completed edition, Heldermann Verlag, Berlin, 1989.
- [6] Matveev M.V., *Absolutely countably compact spaces*, Topology Appl. **58** (1994), 81–92.
- [7] Matveev M.V., *A survey on star-covering properties*, Topological Atlas, no. 330, 1998.
- [8] Shi W.-X., Song Y.-K., Gao Y.-Z., *Spaces embeddable as regular closed subsets into acc spaces and (a)-spaces*, Topology Appl. **150** (2005), 19–31.
- [9] Song Y.-K., *Discretely star-Lindelöf spaces*, Tsukuba J. Math. **25** (2001), no. 2, 371–382.
- [10] Song Y.-K., *Remarks on star-Lindelöf spaces*, Questions Answer Gen. Topology **20** (2002), 49–51.
- [11] Song Y.-K., *Closed subsets of absolutely star-Lindelöf spaces II*, Comment. Math. Univ. Carolin. **44** (2003), no. 2, 329–334.
- [12] Song Y.-K., *Regular closed subsets of absolutely star-Lindelöf spaces*, Questions Answers Gen. Topology **22** (2004), 131–135.
- [13] Song Y.-K., *Some notes on star-Lindelöf spaces*, Questions Answers Gen. Topology **24** (2006), 11–15.
- [14] Vaughan J.E., *Absolutely countably compactness and property (a)*, Talk at 1996 Praha symposium on General Topology.
- [15] Yasui Y., Gao Z.M., *Spaces in countable web*, Houston J. Math. **25** (1999), 327–335.

INSTITUTE OF MATHEMATICS, SCHOOL OF MATHEMATICS AND COMPUTER SCIENCES, NANJING NORMAL UNIVERSITY, NANJING, 210097, P.R. CHINA

*E-mail*: songyankui@njnu.edu.cn

(Received August 21, 2006, revised January 25, 2007)