

František Štulajter

Some nonlinear statistical problems of a Poisson process

Kybernetika, Vol. 18 (1982), No. 5, 397--407

Persistent URL: <http://dml.cz/dmlcz/125869>

Terms of use:

© Institute of Information Theory and Automation AS CR, 1982

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these

Terms of use.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://project.dml.cz>

SOME NONLINEAR STATISTICAL PROBLEMS OF A POISSON PROCESS

FRANTIŠEK ŠTULAJTER

Some results of the theory of random vectors with values in linear spaces are used to study the structure of a space of random variables with finite dispersion generated by a Poisson process and the problem of estimation of nonlinear functionals of an intensity measure of a Poisson process.

1. INTRODUCTION

The aim of this paper is to study some nonlinear statistical problems of a Poisson random process. Similar problems are considered for example in [1] or in [3] for double stochastic Poisson processes. We shall study in more details the structure of the space $L^2(P(\lambda))$ of random variables with finite dispersion generated by a Poisson process with an intensity measure λ and the problem of estimation of nonlinear functionals of an unknown intensity measure λ of a Poisson process. It is shown that $L^2(P(\lambda))$ is equal to the orthogonal sum of $L_n^2(P(\lambda))$; $n \geq 0$, where $L_n^2(P(\lambda))$; $n \geq 0$ are (mutually orthogonal) subspaces of $L^2(P(\lambda))$. L_0^2 is the space of constants, L_1^2 is "the linear subspace" of $L^2(P(\lambda))$, generated by the centered Poisson process, L_2^2 is "the quadratic subspace" of $L^2(P(\lambda))$, and so on. Generating sets of $L_n^2(P(\lambda))$ for $n = 1, 2, 3$ and 4 are given. The same result is true for the space of random variables with a finite dispersion generated by a Gaussian process with zero mean value and a given covariance function as it is shown in [8]. But the rule according to which we form the generating sets of L_n^2 ; $n \geq 0$ for a Gaussian process is different from that derived here for the generating sets of a Poisson process.

In the Part 4 of this paper it is shown that every "polynomial" of a measure λ_f (given by $\lambda_f(A) = \int_A f d\lambda_0$) has an unbiased estimate. It is shown that a dispersion of the best unbiased estimate can be calculated by the same way as it is given in [9].

2. PRELIMINARIES REGARDING POISSON PROCESS

There are many possibilities to define a Poisson process. The best way, for our objective, is to define a Poisson process as a random point measure valued vector as it is done in [7], where the following statements can be found.

Let (T, \mathcal{F}) be a measurable space; denote by $\mathcal{M}(T, \mathcal{F})$ the vector space of finite measures defined on (T, \mathcal{F}) and by $\mathcal{L}_\infty(T, \mathcal{F})$ the space of bounded measurable functions defined on (T, \mathcal{F}) . Let $\mathcal{G}(\mathcal{M}, \mathcal{L}_\infty)$ be a σ -algebra of subsets of $\mathcal{M}(T, \mathcal{F})$, generated by linear transformations $\mu \rightarrow \mu(A)$; $A \in \mathcal{F}$. Then we have: for every fixed finite measure $\lambda \in \mathcal{M}(T, \mathcal{F})$ there exists a unique probability measure $P(\lambda)$ defined on $(\mathcal{M}(T, \mathcal{F}), \mathcal{G}(\mathcal{M}, \mathcal{L}_\infty))$ called the Poisson law with intensity λ on $\mathcal{M}(T, \mathcal{F})$. This measure is a distribution of a Poisson process X transforming a probability space $(\Omega, \mathcal{F}, P_\lambda)$ into $(\mathcal{M}, \mathcal{G})$. Realizations of the random process X have the form $X(\omega) = \sum_{j=1}^{n(\omega)} \delta_{t_j(\omega)}$, where $\{t_1(\omega), \dots, t_{n(\omega)}(\omega)\}$ is a finite set of points of T and δ is a Dirac measure. The random process X has the following properties: for every $A \in \mathcal{F}$ the random variable

$$\langle X(\omega), \chi_A \rangle = \int_A d\left(\sum_{j=1}^{n(\omega)} \delta_{t_j(\omega)}\right) = N_A(\omega) = \text{the number of points } t_j(\omega) \text{ in the set } A,$$

has a Poisson distribution with the parameter $\lambda(A)$. If f_1, \dots, f_n belong to $\mathcal{L}_\infty(T, \mathcal{F})$ and have disjoint supports, then $\langle X, f_1 \rangle, \dots, \langle X, f_n \rangle$ are independent random variables, where

$$\langle X, f_i \rangle(\omega) = \langle X(\omega), f_i \rangle = \int f_i d\left(\sum_{j=1}^{n(\omega)} \delta_{t_j(\omega)}\right); \quad i = 1, \dots, n.$$

The real Laplace transform of the probability space $(\mathcal{M}, \mathcal{G}, P(\lambda))$ is given by $L_{P(\lambda)}(f) = \int e^{\psi(f)} dP(\lambda)$, where ψ is an isomorphism between the vector space $L_0(T, \mathcal{F}, \lambda)$, consisting of classes of equivalence of real measurable functions defined on (T, \mathcal{F}) and the space $L(\mathcal{M}, \mathcal{G}, P(\lambda))$, given by

$$\psi(f) \left(\sum_{j=1}^{n(\omega)} \delta_{t_j(\omega)} \right) = \sum_{j=1}^{n(\omega)} f(t_j(\omega)).$$

We can write $L_{P(\lambda)}(f) = \exp \left\{ \int_T (e^f - 1) d\lambda \right\}$. The Laplace transform is finite, and so defined, for those functions $f \in L_0(T, \mathcal{F}, \lambda)$ for which a function $g = e^f$ belongs to $L(T, \mathcal{F}, \lambda)$; it is the set

$$D = \{f \in L_0 : f = \ln g; g \geq 0, g \in L(T, \mathcal{F}, \lambda)\}.$$

The function $f = 0 \text{ mod } \lambda$ is an inner point of the set D , from which we have that a transformation α_p defined by

$$[\alpha_p(Y)](f) = E_{P_\lambda}[Y \cdot e^{\psi(f)/2}]; \quad Y \in L^2(\mathcal{M}, \mathcal{G}, P_\lambda), \quad f \in D$$

is an isomorphism between the Hilbert spaces $L^2(\mathcal{M}, \mathcal{C}, P(\lambda))$ and a reproducing kernel Hilbert space $H(K_\lambda)$ with the kernel

$$K_\lambda(f, f') = L_{P(\lambda)}\left(\frac{f + f'}{2}\right); \quad f, f' \in D.$$

The problem of equivalence of two Poisson laws $P(\lambda)$ and $P(\lambda_0)$ is solved by the next assertion: let λ and λ_0 be two positive finite measures on (T, \mathcal{F}) . Then $P(\lambda)$ and $P(\lambda_0)$ are equivalent iff λ and λ_0 are equivalent. In the last case denote by $f_\lambda = d\lambda/d\lambda_0$. Then

$$\frac{dP(\lambda)}{dP(\lambda_0)} = \exp\left\{\psi(\ln f_\lambda) - \int_T (f_\lambda - 1) d\lambda_0\right\},$$

where ψ is the above mentioned isomorphism restricted to $L^1(T, \mathcal{F}, \lambda_0)$.

Now let $T = [0, T_0]$, $T_0 > 0$ be an interval on the real line. Then $N(t) = \langle X, \chi_{[0, t]} \rangle$; $0 \leq t \leq T_0$ is a Poisson process with an intensity measure λ , for which we have:

$$E_\lambda[N(t)] = \lambda([0, t]); \quad 0 \leq t \leq T_0$$

and

$$R_\lambda(s, t) = \text{Cov}_\lambda[N(s), N(t)] = \lambda([0, \min(s, t)]) = (\chi_{[0, s]}, \chi_{[0, t]})_{L^2(\lambda)}.$$

In a special case when λ is Lebesgue measure we get $R_\lambda(s, t) = \min(s, t)$, what is the covariance function of the Gaussian Wiener process, too. In the following section we show how these results can be used to solve some nonlinear statistical problems of a Poisson process. The results obtained, are similar to those valid for a Gaussian random process, described in [8] and [9].

3. THE STRUCTURE OF THE SPACE $L^2(\mathcal{M}, \mathcal{C}, P(\lambda))$

Let (T, \mathcal{F}) be a measurable space, λ a finite measure on it and $P(\lambda)$ a distribution of a Poisson process X with values in $(\mathcal{M}, \mathcal{C})$. To solve statistical problems of nonlinear estimation of random variables (for example problems of nonlinear filtration) based on a Poisson process it is necessary to know the structure of the space $L^2(\mathcal{M}, \mathcal{C}, P(\lambda)) = L^2(P(\lambda))$.

It was mentioned in the Section 1 that the Hilbert space $L^2(\mathcal{M}, \mathcal{C}, P(\lambda))$ is isomorphic with the reproducing kernel Hilbert space $H(K_\lambda)$ with a kernel

$$K_\lambda(f, g) = L_{P(\lambda)}\left(\frac{f + g}{2}\right) = \exp\left\{\int_T (e^{f/2} \cdot e^{g/2} - 1) d\lambda\right\},$$

where this kernel is defined on a set $E \times E$ with

$$E = \{f \in L_0 : e^{f/2} \in L^1(T, \mathcal{F}, \lambda)\} = \{f : f = \ln h; h \geq 0, h \in L^2(\lambda)\}.$$

According to this isomorphism, the system of random variables $\{\exp \psi(f); f \in E\}$

generates $L^2(\mathcal{M}, \mathcal{G}, P(\lambda))$. Since it is difficult to characterise the space $H(K_\lambda)$, we use the fact that the set of random variables $\{\exp\{\psi(f) - \int_T (e^f - 1) d\lambda\}; f \in E\}$ generates $L^2(P(\lambda))$ too, and according to Lemma 2 of [8] we have the following assertion: the Hilbert space $L^2(P(\lambda))$ is isomorphic with a reproducing kernel Hilbert space $H(M_\lambda)$ of functionals defined on E , with a kernel

$$\begin{aligned} M_\lambda(f, g) &= E_{P(\lambda)} \left[\exp \left\{ \psi(f) - \int_T (e^f - 1) d\lambda \right\} \exp \left\{ \psi(g) - \int_T (e^g - 1) d\lambda \right\} \right] = \\ &= \exp \left\{ \int_T (e^f - 1)(e^g - 1) d\lambda \right\}, \quad f, g \in E. \end{aligned}$$

Now let $H(N_\lambda)$ be a reproducing kernel Hilbert space with a kernel

$$N_\lambda(h, h') = \exp \left\{ \int_T h \cdot h' d\lambda \right\}; \quad h, h' \in F,$$

where

$$F = \{h \in L^2(T, \mathcal{F}, \lambda) : h \geq -1 \text{ mod } \lambda\}.$$

Define a transformation \mathcal{G} on a set of generating elements of $H(N_\lambda)$ onto a set of generating elements of $H(M_\lambda)$ by $\mathcal{G}(N_\lambda(\cdot, h)) = M_\lambda(\cdot, \ln(h + 1))$; $h \in F$. \mathcal{G} can be naturally extended to an isomorphism between $H(N_\lambda)$ and $H(M_\lambda)$, because we have:

$$\begin{aligned} \langle N_\lambda(\cdot, h), N_\lambda(\cdot, h') \rangle_{H(N_\lambda)} &= \langle \mathcal{G}(N_\lambda(\cdot, h)), \mathcal{G}(N_\lambda(\cdot, h')) \rangle_{H(M_\lambda)} = \\ &= \langle M_\lambda(\cdot, \ln(h + 1)), M_\lambda(\cdot, \ln(h' + 1)) \rangle_{H(M_\lambda)} = \exp \{(h, h')_{L^2(\lambda)}\}; \end{aligned}$$

$h, h' \in F$. Thus we have proved the following lemma:

Lemma 3.1. The Hilbert space $L^2(\mathcal{M}, \mathcal{G}, P(\lambda))$ is isomorphic with the reproducing kernel Hilbert space $H(N_\lambda)$ with the kernel $N_\lambda(h, h') = \exp \left\{ \int_T h h' d\lambda \right\}$; $h, h' \in F$.

Now we are able to give the following theorem.

Theorem 3.1. There exist an isomorphism say φ , between the Hilbert space $L^2(\mathcal{M}, \mathcal{G}, P(\lambda))$ and $\exp \odot L^2(T, \mathcal{F}, \lambda) = \bigoplus_{n \geq 0} L^2(T, \mathcal{F}, \lambda)^{\otimes n}$, where $L^2(\lambda)^{\otimes n}$ is the n -th symmetric tensor power of the space $L^2(\lambda)$.

Proof. It was proved in Lemma 3.1. that $L^2(P(\lambda))$ is isomorphic with $H(N_\lambda; F)$ where $N_\lambda(h, h') = \exp \{(h, h')_{L^2(\lambda)}\}$ is defined on $F \times F$, F being a subset of $L^2(\lambda)$. It is known from the properties of RKHS (see [5]) that $H(N_\lambda; F)$ is isomorphic with a subspace of RKHS $H(N_\lambda; L^2(\lambda))$ of functionals defined on $L^2(\lambda)$ generated by a set of functionals $\{N_\lambda(\cdot, h); h \in F\}$. Since $H(N_\lambda; L^2(\lambda))$ is isomorphic with $\exp \odot L^2(\lambda)$, it is enough to show that the set $\{N_\lambda(\cdot, h); h \in F\}$ generates $H(N_\lambda; L^2(\lambda))$. Let $f \in H(N_\lambda; L^2(\lambda))$ and let $\langle f, N_\lambda(\cdot, h) \rangle_{H(N_\lambda; L^2(\lambda))} = 0$ for all $h \in F$. We have to show that $f = 0$. In our case it holds that $f(g) = \sum_{n \geq 0} (f_n, g \otimes \dots \otimes g)_{L^2(\lambda)^{\otimes n}}$, where

$f_n \in L^2(\lambda)^{n\circ}$. Further we have: $N_\lambda(g, h) = \sum_{n \geq 0} ((1/n!) h \otimes \dots \otimes h, g \otimes \dots \otimes g)_{L^2(\lambda)^{n\circ}}$ and thus $0 = \langle f, N_\lambda(\cdot, h) \rangle_{H(N_\lambda)} = \sum_{n \geq 0} (g_n, h^{\otimes n} \dots \otimes h)_{L^2(\lambda)^{n\circ}}$ for all $h \in F$, where g_n is a projection of f_n onto the subspace $L^2(\lambda)^{n\circ}$ of $L^2(\lambda)^{n\circ}$. From the last equality we get that $\sum_{n \geq 0} t^n (g_n, h^{n\circ})_{L^2(\lambda)^{n\circ}} = 0$ for all $t \geq 0$ and for all $h \in L^2_+(\lambda) = \{h \in L^2(\lambda) : h \geq 0 \text{ mod } \lambda\}$, what is possible only in the case when $(g_n, h^{n\circ}) = 0$ for all $h \in L^2_+(\lambda)$. If we set $h = \sum_{j=1}^n c_j h_j$, where c_1, \dots, c_n are any nonnegative real numbers and $h_1, \dots, h_n \in L^2_+(\lambda)$, then we get that $(g_n, (\sum_{j=1}^n c_j h_j)^{n\circ})_{L^2(\lambda)^{n\circ}}$ – a polynomial in nonnegative variables c_1, \dots, c_n is identically equal to zero, from which we get that $(g_n, h_1 \circ \dots \circ h_n)_{L^2(\lambda)^{n\circ}}; h_1, \dots, h_n \in L^2_+(\lambda)$ – a coefficient of polynomial by a variable $c_1 \dots c_n$ is equal to zero. Since the set $L^2_+(\lambda)$ generates $L^2(\lambda)$, the set $\{h_1 \circ \dots \circ h_n; h_1, \dots, h_n \in L^2_+(\lambda)\}$ generates $L^2(\lambda)^{n\circ}$ for all $n \geq 0$, and thus g_n must be zero element for all $n \geq 0$. \square

Now we shall study in more details the special case when $T = [0, T_0]; T_0 > 0, \mathcal{T} = \mathcal{B}(T)$ and λ is a finite measure on (T, \mathcal{T}) . From Theorem 3.1. we have

Corollary 3.1. The Hilbert space $L^2(\mathcal{M}, \mathcal{G}, P(\lambda))$ is isomorphic with the Hilbert space $\exp \circ H(R_\lambda)$, where $R_\lambda(s, t); s, t \in T$ is the covariance function of a Poisson process $N(t) = \langle X, \chi_{[0, t]} \rangle; 0 \leq t \leq T_0$.

Proof. It was mentioned in Part 1 that $\langle R_\lambda(\cdot, s), R_\lambda(\cdot, t) \rangle_{H(R_\lambda)} = R_\lambda(s, t) = \langle \chi_{[0, s]}, \chi_{[0, t]} \rangle_{L^2(\lambda)}$. Since the system of functions $\{\chi_{[0, t]}; t \in T\}$ generates $L^2(\lambda)$ and the set $\{R_\lambda(\cdot, t); t \in T\}$ generates $H(R_\lambda)$, $L^2(\lambda)$ and $H(R_\lambda)$ are isomorphic. \square

It follows from the definition of $\exp \circ H(R_\lambda)$ as a direct sum of Hilbert spaces $H(R_\lambda)^{n\circ}; n \geq 0$ and from the isomorphism between $L^2(P(\lambda))$ and $\exp \circ H(R_\lambda)$, that the same partition to orthogonal components must hold for the space $L^2(P(\lambda))$, too. According to this we can write: $L^2(\mathcal{M}, \mathcal{G}, P(\lambda)) = \bigoplus_{n \geq 0} L_n^2(\mathcal{M}, \mathcal{G}, P(\lambda))$ where $L_n^2(P(\lambda))$ are orthogonal subspaces of $L^2(\lambda)$. For problems of nonlinear estimation of random variables the following theorems is useful.

Theorem 3.2. Let $T = [0, T_0], T_0 > 0$ and let λ be a finite positive measure on $(T, \mathcal{B}(T))$. Then for any random variable $U \in L^2(\mathcal{M}, \mathcal{G}, P(\lambda))$ we have $U = \bigoplus_{n \geq 0} U_n$, where

$$U_n = \varkappa[E_\lambda[U \cdot \varkappa(R_\lambda(\cdot, t_1) \circ \dots \circ R_\lambda(\cdot, t_n))]]; t_1, \dots, t_n \in T;$$

$n \geq 0$, and \varkappa is an isomorphism described in Corollary 3.1.

Proof. Since the set $\{R_\lambda(\cdot, t_1) \circ \dots \circ R_\lambda(\cdot, t_n); t_1, \dots, t_n \in T\}$ generates $H(R_\lambda)^{n\circ}$, the system of random variables $\{\varkappa(R_\lambda(\cdot, t_1) \circ \dots \circ R_\lambda(\cdot, t_n)); t_1, \dots, t_n \in T\}$ generates the Hilbert space $L_n^2(P(\lambda)); n \geq 0$. A symmetric function of n -variables

$(t_1, \dots, t_n) : E_i[U \cdot \varkappa(R_\lambda(\cdot, t_1) \odot \dots \odot R_\lambda(\cdot, t_n))]$ – an element of $H(R_\lambda^{\otimes n})$, we can identify with that element of the space $H(R_\lambda)^{\otimes n}$, the image of which by the isomorphism \varkappa is the random variable U_n – a projection of a random variable U on the subspace $L_n^2(P(\lambda))$. (For more details see [8]). \square

Now we shall try to clarify how the random variables $\varkappa(R_\lambda(\cdot, t_1) \odot \dots \odot R_\lambda(\cdot, t_n); t_1, \dots, t_n \in T)$ – generating elements of $L_n^2(P(\lambda))$ can be found for $n \geq 0$.

Let φ be the isomorphism from Theorem 3.1. Then we have

$$\varphi(\exp \odot (h - 1)) = \exp \left\{ \psi(\ln h) + \int_T (h - 1) d\lambda \right\}; \quad h \in L_+^2(\lambda),$$

where $\exp \odot h = \sum_{n \geq 0} 1/\sqrt{n!} h^{\otimes n}$ or

$$\varphi(\exp \odot (-f)) = \exp \left\{ \psi(\ln(1 - f)) + \int_T f d\lambda \right\},$$

where f is any function from $L^2(\lambda)$ such that $f \leq 1 \pmod{\lambda}$. If we set $f = \sum_{i=1}^n c_i \chi_{[0, t_i]}$, where $0 < t_1 \leq t_2 \leq \dots \leq t_n$ are any fixed points from the interval $[0, T_0]$, $n \geq 0$ and c_1, \dots, c_n are any suitable chosen real numbers such that $\sum_{i=1}^n c_i \chi_{[0, t_i]} \leq 1$, then we get

$$\begin{aligned} \varkappa(\exp \odot (-\sum_{i=1}^n c_i R_\lambda(\cdot, t_i))) &= \varphi(\exp \odot (-\sum_{i=1}^n c_i \chi_{[0, t_i]})) = \\ &= \exp \left\{ \psi(\ln(1 - \sum_{i=1}^n c_i \chi_{[0, t_i]})) + \sum_{i=1}^n c_i \int_T \chi_{[0, t_i]} d\lambda \right\}. \end{aligned}$$

From the equality

$$\exp \odot (-\sum_{i=1}^k c_i R_\lambda(\cdot, t_i)) = \sum_{n_1=0}^{\infty} \dots \sum_{n_k=0}^{\infty} \frac{c_1^{n_1}}{n_1!} \dots \frac{c_k^{n_k}}{n_k!} (-R_\lambda(\cdot, t_1))^{n_1 \odot} \odot \dots \odot (-R_\lambda(\cdot, t_k))^{n_k \odot}$$

we have that $(-1)^{\otimes n} R_\lambda(\cdot, t_1) \odot \dots \odot R_\lambda(\cdot, t_n)$ is a coefficient by a variable $c_1 \dots c_n$. Since $\{R_\lambda(\cdot, t_1) \odot \dots \odot R_\lambda(\cdot, t_n); t_1, \dots, t_n \in T\}$ generates $H(R_\lambda)^{\otimes n}$, to find $\varkappa(R_\lambda(\cdot, t_1) \odot \dots \odot R_\lambda(\cdot, t_n))$, it suffices to find a coefficient by $c_1 \dots c_n$ in an expansion of the random variable $\exp \left\{ \psi(\ln(1 - \sum_{i=1}^n c_i \chi_{[0, t_i]})) + \sum_{i=1}^n c_i \int_T \chi_{[0, t_i]} d\lambda \right\}$.

To do this, we can proceed as follows: using formally the expression $\ln(1 - x) = -\sum_{k=0}^{\infty} x^{k+1}/(k+1)$ we get

$$\exp \left\{ \psi(\ln(1 - \sum_{i=1}^n c_i \chi_{[0, t_i]})) + \sum_{i=1}^n c_i \int_T \chi_{[0, t_i]} d\lambda \right\} =$$

$$\begin{aligned}
&= \exp \left\{ - \int_T \sum_{k \geq 0} \frac{(\sum_{i=1}^n c_i \chi_{[0, t_i]})^{k+1}}{k+1} dN(t) + \sum_{i=1}^n c_i \int_T \chi_{[0, t_i]} d\lambda \right\} = \\
&= \exp \left\{ \sum_{i=1}^n c_i \left(\int_T \chi_{[0, t_i]} d\lambda - N(t_i) \right) - \int_T \sum_{k \geq 2} \frac{(\sum_{i=1}^n c_i \chi_{[0, t_i]})^k}{k} dN(t) \right\}.
\end{aligned}$$

Expanding the function exp into an infinite series we get the coefficient by $c_1 \dots c_n$ of this expansion. We are not able to derive an general expression for this coefficient for any $n \geq 0$. Here are the first four, derived by this method:

$$\kappa(R_\lambda(\cdot, t)) = N(t) - \int_T \chi_{[0, t]} d\lambda; \quad t \in T$$

Let us denote by $M(t) = N(t) - \int_T \chi_{[0, t]} d\lambda; \quad t \in T$. Then

$$\kappa(R_\lambda(\cdot, t_1) \odot R_\lambda(\cdot, t_2)) = M(t_1) M(t_2) - N(\min\{t_1, t_2\}); \quad t_1, t_2 \in T.$$

$$\begin{aligned}
\kappa(R_\lambda(\cdot, t_1) \odot R_\lambda(\cdot, t_2) \odot R_\lambda(\cdot, t_3)) &= \prod_{i=1}^3 M(t_i) - \sum_{i=1}^3 M(t_i) N(\min(T_3 - \{t_i\})) + \\
&+ 2N(\min T_3), \quad \text{where } T_3 = \{t_1, t_2, t_3\}; \quad t_1, t_2, t_3 \in [0, T_0] = T.
\end{aligned}$$

$$\begin{aligned}
\kappa(R_\lambda(\cdot, t_1) \odot \dots \odot R_\lambda(\cdot, t_4)) &= \prod_{i=1}^4 M(t_i) - \sum_{i < j} M(t_i) M(t_j) N(\min(T_4 - \{t_i, t_j\})) + \\
&+ 2! \sum_{i=1}^4 M(t_i) N(\min(T_4 - \{t_i\})) - 3! N(\min T_4) + \sum_{i=2}^4 N(\min\{t_1, t_i\}) \cdot \\
&\cdot N(\min(T_4 - \{t_1, t_i\})), \quad \text{where } T_4 = \{t_1, \dots, t_4\}; \quad t_1, \dots, t_4 \in T.
\end{aligned}$$

Remark. Setting $t_1 = \dots = t_n = T_0 = 1, n = 1, \dots, 4$ and $\lambda = l$ Lebesgue measure, where $l > 0$, we get the first four orthogonal polynomials of a complete orthogonal system of a Poisson distribution on integers with a parameter l :

$$\begin{aligned}
p_0(x) &= 1 \\
p_1(x) &= x - l \\
p_2(x) &= (x - l)^2 - x \\
p_3(x) &= (x - l)^3 - 3x(x - l) + 2x \\
p_4(x) &= (x - l)^4 - 6x(x - l)^2 + 8x(x - l) + 3x^2 - 6x,
\end{aligned}$$

where

$$\sum_{x \geq 0} p_i(x) p_j(x) \frac{l^x}{x!} e^{-l} = a_j \delta_{ij}; \quad 1, 2, \dots, 4.$$

4. ESTIMATION OF FUNCTIONALS OF AN UNKNOWN INTENSITY MEASURE OF A POISSON LAW

The basis for this part is a general theory of locally best unbiased estimates as given in [6] and used for example in [9]. Now we shall apply this theory to the special case of the estimation of functionals of an unknown intensity measure of a Poisson law.

As we mentioned in Part 2, for two Poisson laws with $\lambda \ll \lambda_0$ on (T, \mathcal{F}) , we have

$$\frac{dP(\lambda)}{dP(\lambda_0)} = \exp \left\{ \psi(\ln f_\lambda) - \int_T (f_\lambda - 1) d\lambda_0 \right\}, \quad \text{where } f_\lambda = \frac{d\lambda}{d\lambda_0}.$$

As we have shown in the preceding part, the system of random variables $\{\exp \{\psi(\ln f) - \int_T (f - 1) d\lambda_0\}; f \in L^2_+(\lambda_0)\}$ generates $L^2(\mathcal{M}, \mathcal{G}, P(\lambda_0))$. Every random variable of a type $\exp \{\psi(\ln f) - \int_T (f - 1) d\lambda_0\}; f \in L^2_+(\lambda_0)$ can be regarded as a Radon-Nikodym derivative $dP(\lambda_f)/dP(\lambda_0)$ of a measure $P(\lambda_f)$ with respect to the measure $P(\lambda_0)$, where λ_f is defined on (T, \mathcal{F}) by $\lambda_f(A) = \int_A f d\lambda_0; f \in L^2_+(\lambda_0), A \in \mathcal{F}$. Thus there exist a one-to-one correspondence between measures λ absolutely continuous with respect to λ_0 and functions (precisely equivalent classes of functions) from $L^2_+(\lambda_0)$.

From a general theory of locally unbiased estimates [6] we have that a functional $F(\cdot)$ defined on a set of measures, which are absolutely continuous with respect to λ_0 , or equivalently, on the set $L^2_+(\lambda_0)$, has an unbiased estimate with a finite dispersion at λ_0 , if and only if, $F(\cdot)$ belongs to a reproducing kernel Hilbert space $H(K_{\lambda_0})$ of functionals defined on $L^2_+(\lambda_0)$ with a kernel

$$K_{\lambda_0}(f, f') = E_{\lambda_0} \left[\frac{dP(\lambda_f)}{dP(\lambda_0)} \frac{dP(\lambda_{f'})}{dP(\lambda_0)} \right] = \exp \left\{ \int_T (f - 1) \cdot (f' - 1) d\lambda_0 \right\}; \quad f, f' \in L^2_+(\lambda_0).$$

It was shown in Theorem 3.1 that $H(K_{\lambda_0})$ and $\exp \odot L^2(\lambda_0)$ are isomorphic, from which we get the following characterization of the space $H(K_{\lambda_0})$, suitable for a case of estimation of functionals.

Theorem 4.1. The reproducing kernel Hilbert space $H(K_{\lambda_0})$ consists of functionals of a type $F_g(\cdot); g \in \exp \odot L^2(\lambda_0)$ defined on the space $L^2_+(\lambda_0)$ and such that

$$F_g(f) = \sum_{n \geq 0} (g_n, (f - 1)^{n \odot})_{L^2(\lambda_0)^{n \odot}}, \quad \text{where } g = \bigoplus_{n \geq 0} g_n \in \exp \odot L^2(\lambda_0)$$

and

$$\|F_g\|_{H(K_{\lambda_0})}^2 = \|g\|_{\exp \odot L^2(\lambda_0)}^2; \quad h^{n \odot} = \frac{1}{\sqrt{n!}} h^{n \otimes} \quad \text{for } h \in L^2(\lambda_0).$$

Proof. Setting $g_n = (g - 1)^{n \odot}; g \in L^2_+(\lambda_0)$ we get, that $F_g(\cdot) = K_{\lambda_0}(\cdot, g)$ is an element of $H(K_{\lambda_0})$. Using the definition of the norm for the class of functionals $F_g(\cdot)$ we get that

$$\langle F_g, K_{\lambda_0}(\cdot, f) \rangle_{H(K_{\lambda_0})} = \sum_{n \geq 0} (g_n, (f - 1)^{n \odot})_{L^2(\lambda_0)^{n \odot}} = F_g(f)$$

for every $g \in \exp \odot L^2(\lambda_0)$, $f \in L^2_+(\lambda_0)$ and the second property of reproducing kernel Hilbert space $H(K_{\lambda_0})$ is proved. \square

It was shown in Part 3 that in the case when $T = [0, T_0]$, $T_0 > 0$, the system $\{\varkappa(R(\cdot, t_1) \odot \dots \odot R(\cdot, t_n)); t_1, \dots, t_n \in T\}$ of random variables generates $L^2_+(P(\lambda_0))$ for every $n \geq 0$. From this we have

$$\begin{aligned} E_{\lambda_0}[\varkappa(R(\cdot, t_1) \odot \dots \odot R(\cdot, t_n))] &= E_{\lambda_0} \left[\varkappa(R[\cdot, t_1] \odot \dots \odot R(\cdot, t_n)) \frac{dP(\lambda_f)}{dP(\lambda_0)} \right] = \\ &= \int_{T^n} \chi_{[t_0, t_1]} \odot \dots \odot \chi_{[t_0, t_n]} (f-1)^{\otimes n} d\lambda_0^{\otimes n} = \\ &= \prod_{i=1}^n \int_T \chi_{[t_0, t_i]} (f-1) d\lambda_0 = \prod_{i=1}^n [\lambda_f([0, t_i]) - \lambda_0([0, t_i])] \end{aligned}$$

for any $f \in L^2_+(\lambda_0)$ and we see that a random variable $\varkappa(R(\cdot, t_1) \odot \dots \odot R(\cdot, t_n))$ is an unbiased estimate of a functional $F_g(f) = \prod_{i=1}^n [\lambda_f([0, t_i]) - \lambda_0([0, t_i])]$ depending on λ_0 .

We are interested in functionals independent of λ_0 . Analogically with results given in [9] we can show that any "polynomial" of a measure λ_f has an unbiased estimate. By a "polynomial of a degree p " we mean a functional $P_p(\cdot)$ given by

$$P_p(f) = \sum_{n=0}^p \int_{T^n} h_n \cdot f^{\otimes n} d\lambda_0^{\otimes n}; \quad f \in L^2_+(\lambda_0), h_n \in L^2(\lambda_0)^{\otimes n}.$$

According to the proof of Lemma 5.1 in [9] we have

$$\int_{T^n} h_n \cdot f^{\otimes n} d\lambda_0^{\otimes n} = \sum_{i=0}^n \binom{n}{i} \int_{T^i} \left(\int_{T^{n-i}} h_n d\lambda_0^{(n-i)\otimes} \right) (f-1)^{\otimes i} d\lambda_0^{\otimes i}$$

for any $n \geq 0$, from which we can derive that any polynomial has an unbiased estimate. For a dispersion of the best unbiased estimate \tilde{P}_p of a polynomial $P_p(f) = \sum_{n=0}^p \int_{T^n} h_n f^{\otimes n} d\lambda_0^{\otimes n}$, where $h_n \in L^2(\lambda_0)^{\otimes n}$ we have from Lemma 5.1. of [9]:

$$\text{Var}_{\lambda_0} [\tilde{P}_p] = \sum_{n=1}^p \sum_{m=1}^p \sum_{i=1}^{\min[m,n]} \binom{n}{i} \binom{m}{i} i! \int_{T^i} \left(\int_{T^{n-i}} h_n d\lambda_0^{(n-i)\otimes} \right) \left(\int_{T^{m-i}} h_m d\lambda_0^{(m-i)\otimes} \right) d\lambda_0^{\otimes i}.$$

Let us investigate a special case when

$$h_n = g_1 \odot \dots \odot g_n, \quad = \frac{1}{\sqrt{n!}} \sum_{\sigma} g_{\sigma_1} \otimes \dots \otimes g_{\sigma_n}.$$

Then we get:

$$\begin{aligned} P_n(f) &= \prod_{j=1}^n \int_T g_j f d\lambda_0 = \int_T h_n f^{\otimes n} d\lambda_0^{\otimes n} = \sum_{i=0}^n \binom{n}{i} \int_{T^i} \left(\int_{T^{n-i}} \frac{1}{n!} \sum_{\sigma} g_{\sigma_1} \otimes \dots \right. \\ &\quad \left. \dots \otimes g_{\sigma_{n-i}} d\lambda_0^{(n-i)\otimes} \right) g_{\sigma_{n-i+1}} \otimes \dots \otimes g_{\sigma_n} \cdot (f-1)^{\otimes i} d\lambda_0^{\otimes i} = \\ &= \sum_{i=0}^n \binom{n}{i} \frac{1}{n!} \sum_{\sigma} \left(\prod_{j=1}^{n-i} \int_T g_{\sigma_j} d\lambda_0 \right) \left(\prod_{j=n-i+1}^n \int_T g_{\sigma_j} (f-1) d\lambda_0 \right) \end{aligned}$$

and

$$\|P_n\|_{H(K\lambda_0)}^2 = E_{\lambda_0}[P_n^2] = \sum_{i=0}^n \binom{n}{i}^2 i! \left\| \frac{1}{n!} \sum_{\sigma} \prod_{j=1}^{n-i} \int_T g_{\sigma_j} d\lambda_0 \otimes_{j=n-i+1}^n g_{\sigma_j} \right\|_{L^2(\lambda_0), \otimes}^2$$

Example 4.1. Let $n = 2$. Then we get:

$$P_2(f) = \prod_{j=1}^2 \int_T g_j f d\lambda_0 = \int_T g_1 d\lambda_0 \int_T g_2 d\lambda_0 + \int_T g_1 d\lambda_0 \int_T g_2(f-1) d\lambda_0 + \int_T g_2 d\lambda_0 \int_T g_1(f-1) d\lambda_0 + \int_T g_1(f-1) d\lambda_0 + \int_T g_2(f-1) d\lambda_0.$$

The locally best unbiased estimate \tilde{P}_2 of P_2 is given

$$\tilde{P}_2 = \prod_{i=1}^2 \int g_i d\lambda_0 + \int_T g_1 d\lambda_0 \cdot \varphi(g_2) + \int_T g_2 d\lambda_0 \cdot \varphi(g_1) + \varphi(g_1 \odot g_2).$$

Setting $g_i = \chi_{[0, t_i]}$; $i = 1, 2$ we get that the random variable $\tilde{P}_2 = N(t_1) \cdot N(t_2) - N(\min\{t_1, t_2\})$ is the best unbiased estimate of the functional

$$P_2(f) = \lambda_f([0, t_1]) \cdot \lambda_f([0, t_2]); \quad t_1, t_2 \in T; \quad f \in L^2_i(\lambda_0)$$

with

$$\begin{aligned} \text{Var}_{\lambda_0}[\tilde{P}_2] &= \|P_2\|_{H(K\lambda_0)}^2 - P_2^2(1) = \int_T g_1^2 d\lambda_0 \int_T g_2^2 d\lambda_0 + \left(\int_T g_1 g_2 d\lambda_0 \right)^2 + \\ &+ \int_T \left[g_1 \int_T g_2 d\lambda_0 + g_2 \int_T g_1 d\lambda_0 \right]^2 d\lambda_0. \end{aligned}$$

If $g_i = \chi_{[0, t_i]}$; $i = 1, 2$, then

$$\begin{aligned} \text{Var}_{\lambda_0}[\tilde{P}_2] &= \lambda_0([0, t_1]) \cdot \lambda_0([0, t_2]) + \lambda_0^2([0, \min\{t_1, t_2\}]) + \\ &+ \lambda_0^2([0, t_2]) \lambda_0([0, t_1]) + 2\lambda_0^2([0, t_1]) \lambda_0([0, \min\{t_1, t_2\}]) + \\ &+ \lambda_0^2([0, t_1]) \lambda_0([0, t_2]). \end{aligned}$$

Setting $t_1 = t_2 = t$, we get

$$\text{Var}_{\lambda_0}[\tilde{P}_2] = 2\lambda_0([0, t]) + 4\lambda_0^3([0, t]) - \text{the classical result.}$$

Example 4.2. Let $P_3(f) = (\int g \cdot f d\lambda_0)^3$: Then

$$P_3(f) = \sum_{i=0}^3 \binom{3}{i} \left(\int g(f-1) d\lambda_0 \right)^i \left(\int_T g d\lambda_0 \right)^{3-i};$$

\tilde{P}_3 – the best unbiased estimate of P_3 is given by:

$$\tilde{P}_3 = \sum_{i=1}^3 \binom{3}{i} \left(\int g d\lambda_0 \right)^{3-i} \varphi(g^{i \odot});$$

$$\text{Var}_{\lambda_0}[\tilde{P}_3] = 6 \cdot \|g\|_{L^2(\lambda_0)}^6 + 18 \|g\|_{L^2(\lambda_0)}^4 \left(\int_T g d\lambda_0 \right)^2 + 9 \|g\|_{L^2(\lambda_0)}^2 \left(\int_T g d\lambda_0 \right)^4.$$

For $g = \chi_{[0, t]}$ we get $\bar{P}_3 = N(t)N(t) - 1(N(t) - 2)$ and

$$\text{Var}_{\lambda_0}[\bar{P}_3] = 6\lambda_0^3([0, t]) + 18\lambda_0^4([0, t]) + 9\lambda_0^5([0, t]),$$

what is again a classical result given in [2].

(Received July 15, 1981.)

REFERENCES

- [1] O. Aalen: Nonparametric inference for a family of counting processes. *Ann. Statist.* **4** (1978), 701–727.
- [2] C. L. Chiang: *Introduction to Stochastic Processes in Biostatistics*. Wiley, New York 1968.
- [3] J. Grandell: *Doubly Stochastic Poisson Processes*. Springer - Verlag, Berlin 1976.
- [4] J. Kerstan, K. Matthes, J. Mecke: *Unbegrenzt teilbare Punktprozesse*. Akademie - Verlag, Berlin 1974.
- [5] J. Neveu: *Processus Aléatoires Gaussiens*. Les Presses de l'Université de Montréal, Montréal 1968.
- [6] E. Parzen: *Time Series Analysis Papers*. Holden-Day, San Francisco 1967.
- [7] J. L. Soler: *Contribution à l'étude des structures statistiques infinidimensionnelles*. Thèse, Grenoble 1978.
- [8] F. Štulajter: RKHS Approach to nonlinear estimation of random variables. *Transaction of the Eighth Prague Conference*, Vol. B, Academia, Prague 1978, 239–246.
- [9] F. Štulajter: Nonlinear estimators of polynomials in mean values of a Gaussian stochastic process. *Kybernetika* **14** (1978), 3, 206–220.

RNDr. František Štulajter, CSc., Katedra teórie pravdepodobnosti a matematickej štatistiky MFF UK (Faculty of Mathematics and Physics, Department of Probability and Statistics – Comenius University), Mlynská dolina, 842 15 Bratislava, Czechoslovakia.