

Jiří Cerha

A method for determining constants in the linear combination of exponentials

*Mathematica Bohemica*, Vol. 121 (1996), No. 2, 121–122

Persistent URL: <http://dml.cz/dmlcz/126106>

## Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A METHOD FOR DETERMINING CONSTANTS IN THE LINEAR  
COMBINATION OF EXPONENTIALS

J. CERHA, Praha

(Received December 7, 1994)

*Summary.* Shifting a numerically given function  $b_1 \exp a_1 t + \dots + b_n \exp a_n t$  we obtain a fundamental matrix of the linear differential system  $\dot{y} = Ay$  with a constant matrix  $A$ . Using the fundamental matrix we calculate  $A$ , calculating the eigenvalues of  $A$  we obtain  $a_1, \dots, a_n$  and using the least square method we determine  $b_1, \dots, b_n$ .

*Keywords:* fundamental matrix, linear differential system, shifted exponentials, eigenvalues, the least square method

*AMS classification:* 65D15, 65L99, 34A30

Let  $n \geq 1$  denote an integer,  $a_1, \dots, a_n; b_1, \dots, b_n$  real numbers,  $a_i \neq a_j$  if  $i \neq j$ ,  $b_i \neq 0$  for  $i = 1, \dots, n$ ,

$$f(t) = b_1 \exp a_1 t + \dots + b_n \exp a_n t$$

for real  $t$ . Let  $h_1, \dots, h_n; k_1, \dots, k_n$  denote real numbers,  $h_1 = k_1 = 0$ ,  $h_i \neq h_j$  and  $k_i \neq k_j$  if  $i \neq j$ ;  $i, j = 1, \dots, n$ . Define the  $n \times n$ -matrix valued function

$$Y(t) = \begin{bmatrix} f(t - h_1 - k_1) & \dots & f(t - h_1 - k_n) \\ \dots & \dots & \dots \\ f(t - h_n - k_1) & \dots & f(t - h_n - k_n) \end{bmatrix} \quad \text{for real } t.$$

**Theorem.**  $Y$  is a fundamental matrix of the linear differential system  $\dot{y} = Ay$  with a constant  $n \times n$ -matrix  $A$ , and  $a_1, \dots, a_n$  are the eigenvalues of  $A$ .

*Proof.* Let us set  $y_i = \exp(-a_i)$ ;  $i = 1, \dots, n$ ,

$$E_1 = E(h_1, \dots, h_n) \equiv \begin{bmatrix} y_1^{h_1} & \dots & y_n^{h_1} \\ \dots & \dots & \dots \\ y_1^{h_n} & \dots & y_n^{h_n} \end{bmatrix}, \quad E_2 = E(k_1, \dots, k_n),$$

$$G = \text{diag}[a_1, \dots, a_n], \quad B = \text{diag}[b_1 \exp a_1 t, \dots, b_n \exp a_n t].$$

Using induction we shall prove that  $E_1$  is regular or, equivalently, the function

$$\varphi(y) = c_1 y^{h_1} + \dots + c_n y^{h_n}$$

has at most  $n - 1$  positive roots for arbitrary  $c_1, \dots, c_n$  excluding  $c_1 = \dots = c_n = 0$  and arbitrary  $h_1, \dots, h_n$  satisfying our assumptions. This is clear for  $n = 1$ . Let  $n > 1$ , let our assertion be true for  $n - 1$  and let us suppose  $\varphi$  has  $n$  positive roots. Hence, the derivative  $\varphi'$  has  $n - 1$  positive roots which, using  $h_1 = 0$ , contradicts the induction hypothesis. Similarly,  $E_2$  is regular. Using our notation we obtain  $Y = E_1 B E_2^T$ ,  $\dot{Y} = E_1 G B E_2^T$ . Hence  $Y$  is regular and  $A \equiv \dot{Y} Y^{-1} = E_1 G E_1^{-1}$  is constant, which proves our theorem.  $\square$

Let  $p \geq 2n$  be an integer,  $t_0, h > 0$  real numbers,  $f_i = f(t_0 - (i - 1)h)$  for  $i = 1, \dots, p$ . Let  $n, h, f_1, \dots, f_p$  be known, while  $a_1, \dots, a_n; b_1, \dots, b_n$  are to be determined. We put  $h_i = k_i = (i - 1)h$  for  $i = 1, \dots, n$ . (However, there exist many methods for choosing  $h_i, k_i$ .) Now, we may calculate  $Y(t)$  for

$$t \in M \equiv \{t_0 - (i - 1)h: i = 1, \dots, q\},$$

where  $q = p - 2n + 2$ ,  $q \geq 2$ . We will determine  $\dot{Y}(t)$  numerically for some fixed  $t \in M$  and put

$$A = \dot{Y}(t)Y(t)^{-1}.$$

Concerning numerical errors, we would probably obtain better results putting

$$A = \frac{1}{m} (\dot{Y}(t_1)Y(t_1)^{-1} + \dots + \dot{Y}(t_m)Y(t_m)^{-1}),$$

where  $m > 1$  is an integer,  $t_1, \dots, t_m \in M$ ,  $t_i \neq t_j$  if  $i \neq j$ . We obtain  $a_1, \dots, a_n$  calculating the eigenvalues of  $A$ .

Alternatively, the formula

$$A = \frac{1}{t_2 - t_1} \ln Y(t_2)Y(t_1)^{-1}$$

can be used. Let  $g_1, \dots, g_n$  denote the eigenvalues of the matrix  $Y(t_2)Y(t_1)^{-1}$  for some fixed  $t_1, t_2 \in M$ ,  $t_1 \neq t_2$ . Hence, the values  $a_1, \dots, a_n$  coincide with the values

$$\frac{1}{t_2 - t_1} \ln g_i; \quad i = 1, \dots, n.$$

Now,  $b_1, \dots, b_n$  may be determined using the least square method.

#### References

- [1] *P. Hartman*: Ordinary differential equations. John Wiley & Sons, New York, London, Sydney, 1964.

*Author's address: Jiří Cerha, Veltruská 533, 190 00 Praha 9, Czech Republic.*