Czechoslovak Mathematical Journal

Cihan Orhan; Ö. Çakar

Some inclusion theorems for absolute summability

Czechoslovak Mathematical Journal, Vol. 46 (1996), No. 4, 599–605

Persistent URL: http://dml.cz/dmlcz/127322

Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 46 (121) 1996, Praha

SOME INCLUSION THEOREMS FOR ABSOLUTE SUMMABILITY

C. ORHAN and Ö. ÇAKAR, Ankara[1]

(Received March 14, 1994)

1. INTRODUCTION

Let $\sum x_n$ be an infinite series with partial sums s_n, and let $A = (a_{nv})$ be a lower semi-matrix with nonzero diagonal entries. By (T_n) we denote the A-transform of the sequence $s = (s_n)$, i.e.,

$$T_n = \sum_{v=0}^{n} a_{nv} s_v \quad (n = 0, 1, 2, \ldots). \tag{1}$$

The series $\sum x_n$ is said to summable $|A|_k (k \geqslant 1)$, if

$$\sum_{n=1}^{\infty} n^{k-1} |T_n - T_{n-1}|^k < \infty \tag{2}$$

(see e.g. [4]).

In the special case of $A = (a_{nv})$ being a Riesz matrix, i.e., weighted mean matrix, we shall write $|R, p_n|_k$ for summability $|A|_k$. The case in which $k = 1$ reduces to the usual absolute weighted mean summability $|R, P_n|$. Recall that a weighted mean matrix is defined by

$$a_{nv} = p_v / P_n \quad \text{for } 0 \leqslant v \leqslant n$$

and

$$a_{nv} = 0 \quad \text{for } v > n$$

where (p_n) is a sequence of positive real numbers and

$$P_n = p_0 + p_1 + \ldots + p_n, \quad P_{-1} = 0.$$

[1] This paper was supported by the Scientific and Technical Research Council of Turkey (TBAG-ÇG 2).

Throughout the paper, we suppose that $P_n \to \infty$ as $n \to \infty$.

In this paper, using functional analytic techniques, we give necessary and sufficient conditions for the series $\sum x_n$ to be summable $|A|_k$ $(k \geqslant 1)$, whenever it is summable $|R, p_n|$, from which we deduce some knonw results.

2. The main result

Given a lower semi-matrix $A = (a_{nv})$, we introduce two lower semi-matrices $\bar{A} = (\bar{a}_{nv})$ and $\hat{A} = (\hat{a}_{nv})$ as follows:

$$\begin{aligned}
\bar{a}_{nv} &= \sum_{i=v}^{n} a_{ni};\ n, v = 0, 1, 2, \ldots, \\
\hat{a}_{nv} &= \bar{a}_{nv} - \bar{a}_{n-1,v};\ n = 1, 2, \ldots, \\
\hat{a}_{00} &= \bar{a}_{00} = a_{00}, \\
\hat{a}_{nv} &= \bar{a}_{nv} = 0 \text{ if } v \geqslant n.
\end{aligned}$$

Since A is a lower semi-matrix, so is $\hat{A}$.

We also note that

$$T_n = (As)_n = \sum_{v=0}^{n} a_{nv} s_v = \sum_{v=0}^{n} \sum_{i=v}^{n} = a_{ni} x_v = \sum_{v=0}^{n} \bar{a}_{nv} x_v$$

and

$$T_n - T_{n-1} = \sum_{v=0}^{n} (\bar{a}_{nv} - \bar{a}_{n-1,v}) x_v = \sum_{v=0}^{n} \Delta \hat{a}_{nv} x_v \tag{3}$$

where $s_v = x_0 + x_1 + \ldots + x_v$ and $\bar{a}_{n-1,n} = 0$.

Using this notation, we have

Theorem. *$|R, p_n|$ summability implies $|A|_k (k \geqslant 1)$ summability if and only if*

$$\text{(i)} \quad |\hat{a}_{vv}| \frac{P_v}{p_v} = O\left(v^{\frac{1}{k}-1}\right),$$

$$\text{(ii)} \quad \left(\sum_{n=v+1}^{\infty} n^{k-1} |\Delta \hat{a}_{nv}|^k \right)^{1/k} = O\left(\frac{p_v}{P_v}\right),$$

$$\text{(iii)} \quad \left(\sum_{n=v+1}^{\infty} n^{k-1} |\hat{a}_{n,v+1}|^k \right)^{1/k} = O(1)$$

where $\Delta \hat{a}_{nv} = \hat{a}_{nv} - \hat{a}_{n,v+1}$.

P r o o f. Necessity. Let t_n be the Riesz means of $\sum x_v$, i.e.

$$t_n = \frac{1}{P_n}\sum_{v=0}^{n} p_v s_v = \frac{1}{P_n}\sum_{v=0}^{n}(P_n - P_{v-1})x_v.$$

Now we have

$$c_n := t_n - t_{n-1} = \frac{p_n}{P_n P_{n-1}}\sum_{v=1}^{n} p_{v-1}x_v, \quad n \geqslant 1, \tag{4}$$
$$c_0 := x_0$$

and

$$C_n := T_n - T_{n-1} = \sum_{v=0}^{n} \hat{a}_{nv}x_v, \quad n \geqslant 1, \tag{5}$$
$$C_0 := x_0.$$

We are given that $|R, p_n| \Longrightarrow |A|_k$, $k \geqslant 1$. Hence

$$\sum_{n=1}^{\infty} n^{k-1}|C_n|^k < \infty \tag{6}$$

whenever

$$\sum |c_n| < \infty. \tag{7}$$

The spaces of sequences (x_v) satisfying (6) and (7) are BK-spaces (i.e., Banach spaces with continuous coordinates) if normed by

$$\|C\| = \left(|C_0|^k + \sum_{n=1}^{\infty} n^{k-1}|C_n|^k\right)^{1/k} \quad \text{and} \quad \|c\| = \sum_{n=1}^{\infty}|c_n|, \tag{8}$$

respectively.

Observe that (5) transforms the space of sequences satisfying (7) into the space of sequences satisfying (6). Applying the Banach-Steinhaus theorem, we find that there is a constant $M > 0$ such that

$$\|C\| \leqslant M\|c\| \tag{9}$$

for all sequences satisfying (7). Applying (4) and (5) to the sequence $x = e_v - e_{v+1}$, where e_v is the vth coordinate vector, we see that

$$c_n = \begin{cases} 0; & n < v \\ \frac{p_v}{P_v}; & n = v \\ \frac{-p_v p_n}{P_n P_{n-1}}; & n > v \end{cases} \quad \text{and} \quad C_n = \begin{cases} 0; & n < v \\ \hat{a}_{vv}; & n = v \\ \hat{a}_{nv} - \hat{a}_{n,v+1}; & n > v. \end{cases}$$

By (8), it follows that

$$\|c\| = \frac{2p_v}{P_v} \text{ and } \|C\| = \left(v^{k-1}|\hat{a}_{vv}|^k + \sum_{n=v+1}^{\infty} n^{k-1}|\Delta\hat{a}_{nv}|^k\right)^{1/k}.$$

By (9), we have

$$v^{k-1}|\hat{a}_{vv}|^k + \sum_{n=v+1}^{\infty} n^{k-1}|\Delta\hat{a}_{nv}|^k \leqslant (2M)^k\left(\frac{p_v}{P_v}\right)^k.$$

Since this holds for any $v \geqslant 1$, we get the necessity of (i) and (ii). To prove the necessity of (iii), we again apply (4) and (5) to the sequence $x = e_{v+1}$. Hence we get that

$$c_n = 0 \quad \text{if} \quad n < v+1$$

and

$$c_n = \frac{P_v p_n}{P_n P_{n-1}} \quad \text{if} \quad n \geqslant v+1,$$

and also

$$C_n = 0 \quad \text{if} \quad n < v+1$$

and

$$C_n = \hat{a}_{n,v+1} \quad \text{if} \quad n \geqslant v+1.$$

By (8) we have

$$\|c\| = 1 \text{ and } \|C\| = \left(\sum_{n=v+1}^{\infty} n^{k-1}|\hat{a}_{n,v+1}|^k\right)^{1/k}.$$

It follows from (9) that

$$\left(\sum_{n=v+1}^{\infty} n^{k-1}|\hat{a}_{n,v+1}|^k\right)^{1/k} = O(1),$$

which implies the necessity of (iii).

Sufficiency. By (4), we have

$$x_v = \frac{P_v}{p_v}c_v - \frac{P_{v-2}}{p_{v-1}}c_{v-1}; \quad P_{-1} = p_{-1} = 0. \tag{10}$$

Inserting (10) in to (5), we may write

$$\begin{aligned} C_n &= \sum_{v=0}^{n} \hat{a}_{nv} x_v = \hat{a}_{n0} c_0 + \sum_{v=1}^{n} \hat{a}_{nv} \Big(\frac{P_v}{p_v} c_v - \frac{P_{v-2}}{p_{v-1}} c_{v-1} \Big) \\ &= \hat{a}_{n0} c_0 + \hat{a}_{nn} \frac{P_n}{p_n} c_n + \sum_{v=1}^{n-1} (\hat{a}_{nv} P_v - \hat{a}_{n,v+1} P_{v-1}) \frac{c_v}{p_v} \\ &= \sum_{v=0}^{n-1} (\hat{a}_{nv} P_v - \hat{a}_{n,v+1} P_{v-1}) \frac{c_v}{p_v} + \hat{a}_{nn} \frac{P_n}{p_n} c_n. \end{aligned}$$

Since

$$\hat{a}_{nv} P_v - \hat{a}_{n,v+1} P_{v-1} = P_v \Delta \hat{a}_{nv} + p_v \hat{a}_{n,v+1},$$

we have

$$C_n = \sum_{v=0}^{n-1} \Big(\frac{P_v}{p_v} \Delta \hat{a}_{nv} + \hat{a}_{n,v+1} \Big) c_v + \hat{a}_{nn} \frac{P_n}{p_n} c_n.$$

Now set $H_n := n^{1-\frac{1}{k}} C_n$, $n \geqslant 1$. Then we get

$$H_n = \sum_{v=1}^{n} u_{nv} c_v$$

where

$$u_{nv} = \begin{cases} n^{(1-\frac{1}{k})} \cdot \Big(\frac{P_v}{p_v} \Delta \hat{a}_{nv} + \hat{a}_{n,v+1} \Big); & 1 \leqslant v \leqslant n-1, \\ n^{(1-\frac{1}{k})} \cdot \frac{P_n}{p_n} \hat{a}_{nn}; & v = n, \\ 0; & v > n. \end{cases}$$

Hence, $\sum x_v$ is summable $|A|_k$, $k \geqslant 1$, whenever $\sum x_v$ is summable $|R, p_n|$ if and only if

$$\sum |H_n|^k < \infty \quad \text{whenever} \quad \sum |c_n| < \infty$$

or equivalently, if and only if the matrix $U = (u_{nv})$ maps l_1 into l_k, $k \geqslant 1$, where

$$l_k = \Big\{ x = (x_v) \colon \sum_v |x_v|^k < \infty \Big\}.$$

Nonetheless, it is well-known that the matrix U maps l_1 into l_k, $k \geqslant 1$, if and only if

$$\sup_v \sum_{n=1}^{\infty} |u_{nv}|^k < \infty$$

(see e.g. [3], Theorem 5, p. 167). □

By the definition of $U = (u_{nv})$, we have

$$\sum_{n=v}^{\infty} |u_{nv}|^k = v^{k-1}\Big(\frac{P_n}{p_n}|\hat{a}_{nn}|\Big)^k + \sum_{n=v+1}^{\infty} n^{k-1}\Big|\frac{P_v}{p_v}\Delta\hat{a}_{nv} + \hat{a}_{n,n+1}\Big|^k.$$

Hence the conditions (i)–(iii) imply that $\sum\limits_{n=v}^{\infty} |u_{nv}|^k = O(1)$ as $v \to \infty$, whence the result.

Taking the matrix $A = (a_{nv})$ to be the weighted mean matrix (R, q_n) where $q_v > 0$ for each v and $Q_n = q_0 + q_1 + \ldots + q_n \to \infty$ as $n \to \infty$, we deduce some known results and list them below:

Corollary 1 ([5]). *$|R, p_n| \Rightarrow |R, q_n|_k$, $k \geqslant 1$ if and only if*

(i) $$\frac{q_v P_v}{Q_v p_v} = O\big(v^{\frac{1}{k}-1}\big),$$

(ii) $$q_v\Big(\sum_{n=v+1}^{\infty} n^{k-1}\Big(\frac{q_n}{Q_n Q_{n-1}}\Big)^k\Big)^{1/k} = O\Big(\frac{p_v}{P_v}\Big),$$

(iii) $$Q_v\Big(\sum_{n=v+1}^{\infty} n^{k-1}\Big(\frac{q_n}{Q_n Q_{n-1}}\Big)^k\Big)^{1/k} = O(1).$$

P r o o f. Apply Theorem with $A = (a_{nv})$ a weighted mean matrix (R, q_n). Observe that, in this case,

$$\hat{a}_{nv} = \frac{q_n Q_{v-1}}{Q_n Q_{n-1}} \quad \text{and} \quad \Delta\hat{a}_{nv} = \hat{a}_{nv} - \hat{a}_{n,v+1} = \frac{-q_n q_v}{Q_n Q_{n-1}}.$$

□

Corollary 2 ([2]). *$|R, p_n| \Rightarrow |R, q_n|$ if and only if*

(11) $$q_v P_v = O(Q_v p_v).$$

P r o o f. Apply Corollary 1 with $k = 1$.

Note that Corollary 2 has been obtained also by Sunouchi [6] in the sufficient form. When reviewing that paper Bosanquet has observed that condition (11) is not only sufficient but also necessary for $|R, p_n| \Rightarrow |R, q_n|$.

When $p_n = 1$ for all n, the $|R, p_n|$ summability is the same as $|C, 1|$ summability. Hence, using Theorem, one can write the necessary and sufficient conditions for $|C, 1| \Rightarrow |A|_k$, $k \geqslant 1$, immediately. So we omit the details. □

3. Concluding remarks

(a) Taking $A = (a_{nv})$ to be the weighted mean matrix (R, q_n) and defining

$$p_v = a^v \quad \text{and} \quad q_v = (v+1)^\alpha$$

where $a > 1$ and $\alpha > -1$, one can see that

$$\frac{P_v}{p_v} \sim \frac{a}{a-1} \quad \text{and} \quad Q_v \sim \frac{vq_v}{\alpha}.$$

Hence conditions (i)–(iii) of Theorem hold.

(b) If we take that matrix $A = (a_{nv})$ to be the weighted mean matrix (R, p_n), then by the condition (i) of Theorem, we must have

$$v^{1-\frac{1}{k}} = O(1),$$

which is impossible when $k > 1$. This means that there is a series $\sum x_n$ which is $|R, p_n|$ summable but not $|R, p_n|_k$, $k > 1$ summable. Actually, such a series is constructed in [5].

References

[1] *L.S. Bosanquet*:Mathematical Reviews *11* (1950), 654.

[2] *G. Das*: Tauberian Theorems for Absolute Nörlund Summability. Proc. London Math. Soc. *19* (1969), 357–384.

[3] *I.J. Maddox*: Elements of Functional Analysis. Cambridge University Press, 1970.

[4] *C. Orhan*: On Absolute Summability. Bull. Inst. Math. Acad. Sinica *15* (1987), 433–437.

[5] *C. Orhan and M.A. Sarigöl*: On Absolute Weighted Mean Summability. Rocky Mountain J. Math. *23* (1993), 1091–1097.

[6] *G. Sunouchi*: Notes on Fourier Analysis XVIII, Absolute Summability of a Series with Constant Terms. Tohoku Math. Journal *2* (1949), 57–65.

Author's address: Department of Mathematics, Faculty of Sciences, Ankara University, 06100 Ankara, Turkey.