Czechoslovak Mathematical Journal

Ján Jakubík

Convexities of normal valued lattice ordered groups

Czechoslovak Mathematical Journal, Vol. 50 (2000), No. 3, 641–649

Persistent URL: http://dml.cz/dmlcz/127599

Terms of use:

© Institute of Mathematics AS CR, 2000

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 50 (125) (2000), 641–649

CONVEXITIES OF NORMAL VALUED LATTICE ORDERED GROUPS

Ján Jakubík, Košice

(Received July 10, 1998)

Convexities of lattice ordered groups were investigated in [5]. Earlier, convexities of lattices and of d-groups had been dealt with in [3] or [4], respectively. Let us recall that the notation of convexity of lattices was introduced by Fried ([7], p. 225).

We denote by

$\mathscr{G}$—the class of all lattice ordered groups;

$\mathscr{A}$—the class of all abelian lattice ordered groups;

$\mathscr{N}$—the class of all normal valued lattice ordered groups;

X_0—the class of all one-element lattice ordered groups.

For $G \in \mathscr{G}$ we denote by $C(G)$ the convexity of lattice ordered groups which is generated by G. Let Z, Q and R be the additive group of all integers, rationals and reals, respectively, with the natural linear order.

If we consider a result on varieties of lattice ordered groups, torsion classes or radical classes, then we can ask whether a similar result holds for convexities.

The following result is well-known (cf., e.g., [1]):

(A) There exists a variety X_1 (namely, $X_1 = \mathscr{A}$) such that, whenever Y is a variety with $Y \neq X_0$, then $X_1 \subseteq Y$.

A result analogous to (A) holds neither for torsion classes nor for radical classes.

In the present paper we prove:

(B) There exists a convexity $Z_1 \neq X_0$ (namely, $Z_1 = C(R)$) such that, whenever Z is a convexity with $X_0 \neq Z \subseteq \mathscr{N}$, then $Z_1 \subseteq Z$.

Some further results are also proved.

Let us remark that the class $\mathscr{N}$ is large in the sense that whenever $\mathscr{V}$ is a variety with $\mathscr{V} \neq \mathscr{G}$, then $\mathscr{V} \subseteq \mathscr{N}$ (cf., e.g., [1]).

Supported by Grant 95/5305/471.

1. The partially ordered classes $\mathscr{C}$ and $\mathscr{C}_{nv}$

For $X \subseteq \mathscr{G}$ we denote by

HX—the class of all homomorphic images of elements of X;

CX—the class of all isomorphic images of convex ℓ-subgroups of elements of X;

PX—the class of all direct products of elements of X.

1.1. Definition. A nonempty subclass of X of $\mathscr{G}$ is called a convexity if $HX \subseteq X$, $CX \subseteq X$ and $PX \subseteq X$.

The class of all convexities of lattice ordered groups will be denoted by $\mathscr{C}$; it is partially ordered by the class-theoretical inclusion. Also, each nonempty subclass of $\mathscr{C}$ is partially ordered by the induced partial order.

1.2. Lemma. (Cf. [5].) *Let $\emptyset \neq X \subseteq \mathscr{G}$. Then*

(i) *$HCPX \in \mathscr{C}$;*

(ii) *for each $Y \in C$ with $X \subseteq Y$ the relation $HCPX \subseteq Y$ is valid.*

In view of 1.2, the convexity $HCPX$ will be said to be generated by X. If $X = \{G\}$ is a one-element set, then we put $HCPX = C(G)$.

For direct products of lattice ordered groups we apply the same notation and conventions as in [5], Section 1.

From 1.2 we immediately obtain

1.3. Lemma. *Let $\{X_i\}_{i\in I}$ be a nonempty subclass of $\mathscr{C}$. Then $\bigcap\limits_{i\in I} X_i \in \mathscr{C}$.*

Let X_0 be as above. It is obvious that X_0 is the least element of $\mathscr{C}$ and that $\mathscr{G}$ is the greatest element of $\mathscr{C}$. From this and from 1.3 we obtain

1.4. Lemma. *Let $\{X_i\}_{i\in I}$ be a nonempty subclass of $\mathscr{C}$. Then there exist Y_1 and Y_2 in $\mathscr{C}$ such that the relations $Y_1 = \inf\{X_i\}_{i\in I}$ and $Y_2 = \sup\{X_i\}_{i\in I}$ are valid in the partially ordered collection $\mathscr{C}$. Moreover, $Y_1 = \bigcap\limits_{i\in I} X_i$.*

In [5] it was proved that the collection $\mathscr{C}$ is large in the sense that there exists an injective mapping of the class of all infinite cardinals into $\mathscr{C}$.

Nevertheless, in view of 1.4 we can apply for $\mathscr{C}$ the usual lattice-theoretical terminology and notation. Thus, if Y_1 and Y_2 are in 1.4, then we write

$$Y_1 = \bigwedge_{i\in I} X_i, \quad Y_2 = \bigvee_{i\in I} X_i.$$

1.5. Lemma. *Let $\{X_i\}_{i\in I}$ be a nonempty subclass of $\mathscr{C}$. Then*

$$\bigvee_{i\in I} X_i = HCP\Big(\bigcup_{i\in I} X_i\Big).$$

P r o o f. This is a consequence of 1.2. □

1.6. Proposition. *Let $X_1, X_2 \in \mathscr{C}$. Next, let Y be the set of all $G \in \mathscr{G}$ such that there exist $G_1 \in X_1$ and $G_2 \in X_2$ with $G = G_1 \times G_2$. Then $X_1 \vee X_2 = Y$.*

P r o o f. Let $G \in Y$. Then (under the notation as above) we have $G_1, G_2 \in X_1 \vee X_2$, whence $G \in X_1 \vee X_2$.

Conversely, let $G \in X_1 \vee X_2$. In view of 1.5 there exists a set $\{G_j\}_{j\in J} \subseteq X_1 \cup X_2$ such that

$$G \in HCP\{G_j\}_{j\in J}.$$

Hence there are $A_1, A_2 \in \mathscr{G}$ with

$$A_1 \in P\{G_j\}_{j\in J}, \quad A_2 \in C\{A_1\}, \quad G \in H\{A_2\}.$$

Therefore

$$A_1 = \Big(\prod_{j\in J(1)} G_j\Big) \times \Big(\prod_{j\in J(2)} G_j\Big),$$

where

$$\{G_j\}_{j\in J(1)} \subseteq X_1, \quad \{G_j\}_{j\in J(2)} \subseteq X_2.$$

Put

$$\prod_{j\in J(1)} G_j = G_1^1, \qquad \prod_{j\in J(2)} G_j = G_2^1.$$

Hence $G_1^1 \in X_1$ and $G_2^1 \in X_2$.

From the relation $A_1 = G_1^1 \times G_2^1$ and from Lemma 1.2 in [5] we obtain

$$A_2 = (A_2 \cap G_1^1) \times (A_2 \cap G_2^1).$$

Since $A_2 \cap G_i^1$ is a convex ℓ-subgroup of G_i^1 we get $A_2 \cap G_i^1 \in X_i$ $(i = 1, 2)$. Now it suffices to apply Lemma 1.3 from [5] to verify that $G \in Y$. □

1.7. Theorem. *The lattice $\mathscr{C}$ is distributive.*

P r o o f. Let $X_1, X_2, X_3 \in \mathscr{C}$. We have to verify that the relation

$$X_1 \wedge (X_2 \vee X_3) = (X_1 \wedge X_2) \vee (X_1 \wedge X_3)$$

is valid. Clearly $(X_1 \wedge X_2) \vee (X_1 \wedge X_3) \subseteq X_1 \wedge (X_2 \vee X_3)$. Let $G \in X_1 \wedge (X_2 \vee X_3)$. Thus $G \in X_1$ and $G \in X_2 \vee X_3$. According to 1.6 there are $G_2 \in X_2$ and $G_3 \in X_3$ such that $G = G_2 \times G_3$. Hence $G_2, G_3 \in C\{G\}$ and therefore $G_2, G_3 \in X_1$. We get

$$G_i \in X_1 \wedge X_i \quad (i = 2, 3)$$

and hence $G \in (X_1 \wedge X_2) \vee (X_1 \wedge X_3)$, completing the proof. $\square$

A complete lattice is said to be infinitely distributive if it satisfies the identities

$$x \vee \left(\bigwedge_{i \in I} y_i \right) = \bigwedge_{i \in I} (x \vee y_i), \tag{1}$$

$$x \wedge \left(\bigvee_{i \in I} y_i \right) = \bigvee_{i \in I} (x \wedge y_i). \tag{2}$$

The collection of all radical classes of lattice ordered groups satisfies identically the relation (2). The question whether $\mathscr{C}$ satisfies (1) or (2) remains open.

Let $G \in \mathscr{G}$ and $0 \neq g \in G$. The convex ℓ-subgroup of G generated by g will be denoted by $[g]$. Next, let $C_1(g)$ be the set of all convex ℓ-subgroups of $[g]$ which do not contain the element g, and let $C_2(g)$ be the set of all maximal elements of $C_1(G)$.

A lattice ordered group G is said to be normal valued if, whenever $0 \neq g \in G$ and $G' \in C_2(g)$, then G' is a normal subgroup of $[g]$.

Let $\mathscr{N}$ be as above. From the fact that $\mathscr{N}$ is a variety (cf., e.g., [1]) we infer that $\mathscr{N}$ belongs to $\mathscr{C}$. Hence the class $\mathscr{C}_{nv}$ of all convexities X with $X \subseteq \mathscr{N}$ is the interval $[X_0, \mathscr{N}]$ of $\mathscr{C}$. Therefore 1.7 yields

1.8. Corollary. *$\mathscr{C}_{nv}$ is a distributive lattice.*

The following result is well-known.

1.9. Lemma. *Let $G \in \mathscr{N}$, $0 \neq g \in G$, $G' \in C_2(G)$. Then there exists an isomorphism φ of the lattice ordered group $[g]/G'$ into R.*

2. Convexities generated by ℓ-subgroups of R

We start by investigating the convexity $C(Z)$.

Let $\mathbb{N}$ be the set of all positive integers and for each $n \in \mathbb{N}$ let $G_n = Z$. Denote $G^1 = \prod_{n\in\mathbb{N}} G_n$. For $g \in G^1$ we denote by g_n the n-th component of g. If there exists a positive integer m such that $|g(n)| \leqslant m$ for each $n \in \mathbb{N}$, then g will be said to be bounded. (In an analogous sense we apply the notion of boundedness also when dealing with any direct product of ℓ-subgroups of R.) The set of all bounded elements of G^1 will be denoted by G^2.

The set G^2 is an ℓ-ideal of G^1, thus we can construct the factor lattice ordered group $\overline{G^1} = G^1/G^2$ and we have $\overline{G^1} \in C(Z)$. For $g \in G^1$ we put $\overline{g} = g + G^2$.

2.1. Lemma. *The lattice ordered group $\overline{G^1}$ is divisible.*

Proof. It suffices to verify that for each positive integer m and for each strictly positive element $\overline{g} = g + G^2$ of $\overline{G^1}$ there exists $g' \in G^1$ such that $m\overline{g'} = \overline{g}$.

If $\overline{g} > \overline{0}$, then without loss of generality we can suppose that $g_n \geqslant 0$ for each $m \in \mathbb{N}$, and that the sequence $(g_n)_{n\in\mathbb{N}}$ is not bounded.

Let $m, n \in \mathbb{N}$. There is a real x_n such that $mx_n = g_n$. Next, there is a real z_n such that $0 \leqslant z_n < 1$ and $x_n + z_n \in Z$. Put $x_n + z_n = y_n$.

There are $g', z' \in G'$ such that

$$g'_n = y_n, \quad z'_n = 1 \quad \text{for each } n \in \mathbb{N}.$$

Hence

$$g_n = mx_n \leqslant my_n = mx_n + mz_n < g_n + m.$$

Thus $g \leqslant mg' < g + z'$ and therefore

$$\overline{g} \leqslant m\overline{g'} \leqslant \overline{g} + m\overline{z'}.$$

Clearly $z' \in G^2$, whence $\overline{z'} = \overline{0}$. We conclude that $\overline{g} = m\overline{g'}$. □

2.2. Lemma. *There exists $\{0\} \neq G^3 \in C(Z)$ such that*

(i) *G^3 is an ℓ-subgroup of R;*

(ii) *G^3 is divisible.*

Proof. Let $\overline{G^1}$ be as in 2.1. Then $\overline{G^1}$ belongs to $C(Z)$. There exists $\overline{g} \in \overline{G^1}$ with $\overline{g} \neq \overline{0}$. Let $G' \in C_2(\overline{g})$ and denote

$$G^3_0 = [\overline{g}]/G'$$

(with respect to the lattice ordered group $\overline{G^1}$). Then G_0^3 is a nonzero lattice ordered group. We have $[\overline{g}] \in C(Z)$ and hence $G_0^3 \in C(Z)$. In view of 2.1, $[\overline{g}]$ is divisible and thus G_0^3 is divisible as well. Now it suffices to apply 1.9. $\square$

Let us consider the following condition for a convexity X:

$(*)$ There exists G^3 in X satisfying the conditions (i) and (ii) from 2.2.

Suppose that $(*)$ is valid. Put $H_n = G^3$ for each $n \in \mathbb{N}$ and

$$G^4 = \prod_{n \in \mathbb{N}} H_n.$$

For $g \in G^4$ let g_n be the component of g in H_n. Next, let K be the set of all bounded elements of G^4. We investigate the factor lattice ordered group G^4/K; we denote $\overline{g} = g + K$. Since G^3 is divisible, so are G^4 and G^4/K. Hence for each $q \in Q$ we can construct $qg \in G^4$ and $q\overline{g} \in G^4/K$.

Let $0 < r \in R$. Put

$$Q_1 = \{g \in Q : 0 < q < r\}, \quad Q_2 = \{q \in Q : r < q\}.$$

There exists a sequence $\{q_{(n)}\}_{n \in \mathbb{N}}$ of elements of Q_1 such that $q_{(n)} < q_{(n+1)}$ for each $n \in \mathbb{N}$, and $\sup\{q_{(n)}\}_{n \in \mathbb{N}} = r$.

Under the above notation we have

2.3. Lemma. *Let $0 < g \in G^4$ with $\overline{g} > \overline{0}$ and let $0 < r \in R$. There exists $g' \in G^4$ such that $q_1\overline{g} \leqslant \overline{g'} \leqslant q_2\overline{g}$ for each $q_1 \in Q_1$ and each $q_2 \in Q_2$.*

Proof. There is $g' \in G^4$ such that

$$g'_n = q_{(n)} g_n$$

for each $n \in \mathbb{N}$. Thus $g'_n < q_2 g_n$ for each $n \in \mathbb{N}$ and each $q_2 \in Q_2$. Hence

$$\overline{g'} \leqslant q_2\overline{g} \quad \text{for each } q_2 \in Q_2. \tag{1}$$

Let $q_1 \in Q_1$. There exists $m \in \mathbb{N}$ such that $q_1 < q_{(m)}$. Further, there are elements $g_{(1,m)}$ and $g_{(2,m)}$ in G^4 such that

$(g_{(1,m)})_n = g_n$ if $n < m$, and $(g_{(1,m)})_n = 0$ otherwise;

$(g_{(2,m)})_n = g_n$ if $n \geqslant m$, and $(g_{(2,m)})_n = 0$ otherwise.

Similarly we construct $g'_{(1,m)}$ and $g'_{(2,m)}$ (with g replaced by g'). Thus

$$g = g_{(1,m)} + g_{(2,m)}, \tag{2}$$

$$g' = g'_{(1,m)} + g'_{(2,m)}. \tag{3}$$

Clearly $g_{(1,m)}, g'_{(1,m)} \in K$, whence

$$\overline{g} = \overline{g_{(2,m)}}, \quad \overline{g'} = \overline{g'_{(2,m)}}. \tag{4}$$

We have also

$$q_1 g_{(2,m)} < g'_{(2,m)},$$

whence

$$q_1 \overline{g_{(2,m)}} \leqslant \overline{g'_{(2,m)}}. \tag{5}$$

The relations (2)–(5) yield that $q_1\overline{g} \leqslant \overline{g'}$ for each $q_1 \in Q_1$. Hence, by virtue of (1), the proof is complete. □

2.4. Lemma. *Let X be a convexity satisfying the condition $(*)$. Then $R \in X$.*

Proof. Let G^4, g and r be as in 2.3. Then $G^4 \in X$. Also, G^4/K belongs to X. Let $G' \in C_2(\overline{g})$. We construct the lattice ordered group $[\overline{g}]/G'$ (with respect to G^4/K). For $\overline{x} \in [\overline{g}]$ we denote $\overline{\overline{x}} = \overline{x} + G'$.

According to 1.9 there exists an isomorphism φ of $[\overline{g}]/G'$ into R. Denote $\varphi(\overline{\overline{g}}) = r_0$. Then $r_0 > 0$. From this we infer that there exists an isomorphism φ_1 of $[\overline{g}]/G'$ into R such that $\varphi_1(\overline{\overline{g}}) = 1$.

Let $\overline{g'}$ be as in 2.3. Put $\varphi_1(\overline{\overline{g'}}) = r'$. In view of 2.3 we have

$$q_1\varphi_1(\overline{\overline{g}}) \leqslant \varphi_1(\overline{\overline{g'}}) \leqslant q_2\varphi_1(\overline{\overline{g}}),$$

hence $q_1 \leqslant r' \leqslant q_2$ whenever $q_1 < r < q_2$. Thus $r = r'$ and therefore $r \in \varphi_1([g]/G')$. Then, clearly, φ_1 is an epimorphism. Since $[\overline{g}]/G'$ belongs to X we get that R belongs to X as well. □

2.5. Lemma. $R \in C(Z)$.

Proof. This is a consequence of 2.1–2.4. □

Now let G be an ℓ-subgroup of R such that $G \neq \{0\}$ and G fails to be isomorphic to Z. By an elementary argument we obtain

2.6. Lemma. *Let $0 < x \in R$. Then there exists $g_0 \in G$ such that $0 < g_0 < x$.*

For each $n \in \mathbb{N}$ let $G_n = G$. Put $G_0^1 = \prod_{n\in\mathbb{N}} G_n$. Next, let G^2 be the ℓ-subgroup of G_0^1 consisting of all bounded elements of G_0^1. We denote $\overline{G_0^1} = G_0^1/G^2$, $\overline{g} = g + G^2$, where $g \in G_0^1$.

2.7. Lemma. *The lattice ordered group $\overline{G_0^1}$ is divisible.*

P r o o f. We proceed in the same way as in the proof of 2.1 with the distinction that instead of

$$y_n = x_n + z_n \in Z$$

we now consider the relation

$$y_n = x_n + z_n \in G$$

and apply 2.6. □

2.8. Lemma. $R \in C(G)$.

P r o o f. This is a consequence of 2.7, 2.3 and 2.4. □

2.9. Theorem. *Let X be a convexity of normal valued lattice ordered groups and let $X \neq X_0$. Then $C(R) \subseteq X$.*

P r o o f. There is $G_1 \in X$ with $G_1 \neq \{0\}$. Hence in view of 1.9 there exists $G \in X$ such that $G \neq \{0\}$ and G is an ℓ-subgroup of R. According to 2.5 and 2.8, R belongs to X. Thus $C(R) \subseteq X$. □

In other words, we have proved that the interval $[X_0, \mathscr{N}]$ of $\mathscr{C}$ has a unique atom.

2.10. Corollary. $C(R) \subset C(Z)$.

P r o o f. In view of 1.2, $C(R)$ is divisible. Hence $Z \notin C(R)$ and thus, according to 2.9, $C(R) \subset C(Z)$. □

We denote by $X_{v\ell}$ the class of all lattice ordered groups G which satisfy the following condition:

(vℓ) We can define a multiplication of elements of G by reals such that G turns out to be a vector lattice.

It is obvious that

(i) $X_{v\ell}$ is closed with respect to H, C and P; hence $X_{v\ell}$ is a convexity;
(ii) if G_1 is an ℓ-subgroup of R with $\{0\} \neq G_1 \in X_{v\ell}$, then $G_1 = R$.

The following result generalizes 2.10.

2.11. Proposition. *Let G_1 be an ℓ-subgroup of R such that $\{0\} \neq G_1 \neq R$. Then $C(R) \subset C(G_1)$.*

P r o o f. In view of 2.9 we have $C(R) \subseteq C(G_1)$. By way of contradiction, suppose that $C(R) = C(G_1)$. Hence $G_1 \in C(R)$. But $X \in X_{v\ell}$ and thus according to (i), G_1 belongs to $X_{v\ell}$ as well. This contradicts (ii). □

In particular, $C(R) \subset C(Q)$. An open question: what are the relations between $C(Q)$ and $C(Z)$?

References

[1] *A. M. W. Glass, W. C. Holland, S. M. McCleary*: The structure of ℓ-group varieties. Algebra Universalis *10* (1980), 1–20.

[2] *J. Jakubík*: Radical classes and radical mappings of lattice ordered groups. Sympos. Math. *31* (1977), 451–477. Academic Press, NewYork-London.

[3] *J. Jakubík*: On convexities of lattices. Czechoslovak Math. J. *42* (1992), 325–330.

[4] *J. Jakubík*: On convexities of d-groups. Czechoslovak Math. J. *44* (1994), 305–314.

[5] *J. Jakubík*: Convexities of lattice ordered groups. Math. Bohem. *121* (1996), 59–67.

[6] *J. Martinez*: Torsion theory of lattice ordered groups. Czechoslovak Math. J. *25* (1975), 284–299.

[7] *R. Mlitz*: General Algebra 1988. Proc. Internat. Conf. Krems 1988. North Holland, Amsterdam 1990.

Author's address: Matematický ústav SAV, Grešákova 6, 040 01 Košice, Slovakia.