Czechoslovak Mathematical Journal

Carlos Bosch; Jan Kučera

Sequential completeness and regularityof inductive limits of webbed spaces

Czechoslovak Mathematical Journal, Vol. 52 (2002), No. 2, 329–332

Persistent URL: http://dml.cz/dmlcz/127720

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 52 (127) (2002), 329–332

SEQUENTIAL COMPLETENESS AND REGULARITY OF INDUCTIVE LIMITS OF WEBBED SPACES

Carlos Bosch, San Angel, and Jan Kučera, Pullman

(Received March 22, 1999)

Abstract. Any inductive limit of bornivorously webbed spaces is sequentially complete iff it is regular.

Keywords: localy convex space, webbed space, sequential completeness, regularity of an inductive limit

MSC 2000: 46A13, 46A04

A web in a vector space F is a countable family of balanced subsets of F, arranged in "layers". The first layer of the web consists of a sequence $(A_p\colon\ p = 1, 2, \ldots)$ whose union absorbs each point of F. For each set A_p of the first layer there is a sequence $(A_{pq}\colon\ q = 1, 2, \ldots)$ of sets, called the sequence determined by A_p such that

$$A_{pq} + A_{pq} \subset A_p \text{ for each } q, \text{ and}$$
$$\bigcup\{A_{pq}\colon\ q = 1, 2, \ldots\} \text{ absorbs each point of } A_p.$$

Further layers are made up in a corresponding way so that each set of the k-th layer is indexed by a row of k integers and at each step the two conditions above are satisfied. Suppose that one chooses a set A_p then A_{pq} from the sequence determined by A_p, and so on. The resulting sequence $S = (A_p, A_{pq}, A_{pqr}, \ldots)$ is called a strand. Whenever we are dealing with only one strand we can simplify the notation by writing $W_1 = A_p$, $W_2 = A_{pq}$, etc.; thus $S = (W_k)$ is a strand where for each k, W_k is a set in the k-th layer.

A web in a locally convex space is called strict if each of its sets is absolutely convex and for any strand (W_k) there exists a sequence $a_1 > a_2 > \ldots > 0$, called

Partially supported by Fulbright grant No. 22729.

completing, such that for any choice of $b_k \in [0, a_k]$, $x_k \in W_k$, $k \in \mathbb{N}$, and any $n \in \mathbb{N}$, the sequence $\sum\limits_{k=n+1}^{\infty} b_k x_k$ converges to some element in W_n. A locally convex space F is strictly webbed if it admits a strict web. It is bornivorously webbed if it is strictly webbed and for every set B bounded in F there exist a strand (W_k) of a strict web and a sequence (λ_k) such that $B \subset \lambda_k W_k$ for every $k \in \mathbb{N}$ ([1], [2], [3]).

Lemma. *Let (F, α) be a strictly webbed space with a locally convex topology α. Take a strand (W_k) of a strict webb in F and denote by γ the topology generated by the subbasis $\{W_k\colon\ k \in \mathbb{N}\}$. Then:*

(a) *The topology γ is stronger than α.*

(b) *The space (F, γ) is Fréchet.*

P r o o f. (a) Let (a_k) be a completing sequence for the strand (W_k). Assume that the topology γ is not stronger than α. Then there exists a balanced 0-neighbourhood $U \in \alpha$ such that $a_k W_k \setminus U \neq \emptyset$ for any $k \in \mathbb{N}$. Take a 0-neighbourhood V in α such that $V - V \subset U$, choose $y_k \in a_k W_k \setminus U$, and put $x_k = a_k^{-1} y_k$, $k \in \mathbb{N}$. Since the space (F, α) is strictly webbed, the series $\sum\limits_{k=1}^{\infty} a_k x_k$ converges in (F, α) and there exists $k \in \mathbb{N}$ such that for any $m \geqslant k$, we have $\sum\limits_{n=m}^{\infty} a_k x_k \in V$. This implies $y_k = a_k x_k = \sum\limits_{n=k}^{\infty} a_n x_n - \sum\limits_{n=k+1}^{\infty} a_n x_n \in V - V \subset U$, a contradiction.

(b) Since the topology γ has a countable subbasis, it is metrizable. It remains to show that (F, γ) is complete. Take a Cauchy sequence $\{y_k;\ k \in \mathbb{N}\}$ in (F, γ) and a completing sequence (a_k) for the strand (W_k). Without a loss of generality, we may assume that $y_{k+1} - y_k \in a_k W_k$, $k \in \mathbb{N}$. Then $x_k = a_k^{-1}(y_{k+1} - y_k) \in W_k$ for any $k \in \mathbb{N}$ and $\sum\limits_{k=1}^{\infty} a_k x_k = \sum\limits_{k=1}^{\infty} (y_{k+1} - y_k)$ converges to an element $y \in F$. Moreover $\sum\limits_{k=n+1}^{\infty} (y_{k+1} - y_k) = \sum\limits_{k=n+1}^{\infty} a_k x_k \in W_n$, $n \in \mathbb{N}$. This implies that, in the topology γ, the series $\sum\limits_{k=1}^{\infty} (y_{k+1} - y_k)$ converges to an element $y \in F$ and $y_{k+1} = y_1 + \sum\limits_{k=1}^{n} (y_{k+1} - y_k) \longrightarrow y_1 + y \in F$. $\square$

Let $E_1 \subset E_2 \subset \ldots$ be a sequence of locally convex spaces with respective topologies τ_n, $n \in \mathbb{N}$, such that the identity map $\mathrm{id}\colon (E_n, \tau_n) \to (E_{n+1}, \tau_{n+1})$ is continuous for every $n \in \mathbb{N}$. Its locally convex inductive limit is denoted by $\operatorname{ind} E_n$.

We call $\operatorname{ind} E_n$ regular if for each set B, bounded in $\operatorname{ind} E_n$, there exists $n \in \mathbb{N}$ such that B is bounded in (E_n, τ_n).

Theorem. *If each space E_n, $n \in \mathbb{N}$, is bornivorously webbed then* $\operatorname{ind} E_n$ *is sequentially complete if and only if it is regular.*

P r o o f. For brevity, we will write just E instead of $\operatorname{ind} E_n$.

Suppose that the space E is sequentially complete, then it is fast complete. By Theorem 1 in [4], fast complete inductive limit of webbed spaces is regular.

Let E be regular, $\{x_n;\ n \in \mathbb{N}\}$ a Cauchy sequence in E, and $B_n = \operatorname{cl}_E \operatorname{co} \bigcup \{x_m;\ m \geqslant n\}$, $n \in \mathbb{N}$. The set B_1 is bounded in E and, by the regularity of E, it is also bounded in some space E_m. Without a loss of generality, we may assume $m = 1$.

The space (E_1, τ_1) is bornivorously webbed, hence there exists a strand (W_k) in E_1 and a sequence $\{\alpha_k;\ k \in \mathbb{N}\}$ such that $B_1 \subset \alpha_k W_k$ for each $k \in \mathbb{N}$. Denote by γ the topology on E_1 generated by the subbasis $\{W_k;\ k \in \mathbb{N}\}$ and, for brevity, by F the space (E_1, γ).

The set $B_1 \subset E_1$ is closed in E. Hence it is closed in (E_1, τ_1), and by Lemma, it is closed in the locally convex space $F = (E_1, \gamma)$. Since B_1 is convex, it is also weakly closed in F.

By Lemma, F is a Fréchet space. Hence the canonical inbedding $F \to F''$, where F'' is the second dual of F equipped with the strong topology, is a topological isomorphism into F''. Since F is complete, it is closed in F'' and each functional from the strong dual F' of F can be continuously extended to F''. Thus the $\sigma(F, F')$-closed set B_1 is also $\sigma(F'', F')$-closed in F''.

Further, since B_1 is bounded in F'', it is equicontinuous in F'. Hence, by Alaoglu Theorem, the set B_1 is relatively $\sigma(F'', F')$-compact. This, together with the $\sigma(F'', F')$-closedness, implies that B_1 is $\sigma(F'', F')$-compact in F''.

Similarly, all sets B_n, $n \in \mathbb{N}$, are $\sigma(F'', F')$-compact. Every finite intersection $\bigcap \{B_n;\ 1 \leqslant n \leqslant m\} = B_m$, $m \in \mathbb{N}$, is non-empty. Hence there exists $x_0 \in \bigcap \{B_n;\ n \in \mathbb{N}\} \subset B_1 \subset E_1$. This implies the existence of an upper-triangular matrix $\Lambda = (\lambda_{nm})$ with all entries $\lambda_{nm} \geqslant 0$, only finite number of non-zeros in each row, and the sum of all entries in each row equal to 1, such that the sequence $\{y_n = \sum\limits_{m=n}^{\infty} \lambda_{nm} x_m;\ n \in \mathbb{N}\}$ converges to x_0 in the topology γ. Then the continuity of the identity maps: $(E_1, \gamma) \to (E_1, \tau_1) \to \operatorname{ind} E_n$ implies the convergence $y_k \to x_0$ in $\operatorname{ind}\ E_n$.

Take a balanced, convex, 0-neighbourhood V in E. Then there exist $p, q \in \mathbb{N}$ such that $y_n - x_0 \in V$ for $n \geqslant p$ and $x_m - x_n \in V$ for $m \geqslant n \geqslant q$. Then for $n \geqslant \max(p, q)$, we have $x_0 - x_n = (x_0 - y_n) + (y_n - x_n) = (x_0 - y_n) + \sum\limits_{m=n}^{\infty} \lambda_{nm}(x_m - x_n) \in V + V$. This implies $x_n \to x_0$ in the space E. $\square$

Corollaries.

(a) *Since any LF-space is bornivorously webbed, it is sequentially complete iff it is regular.*

(b) *Theorem* 1 *in* [4] *reads: If all locally convex spaces* (E_n, τ_n), $n \in \mathbb{N}$, *are webbed and their* $\operatorname{ind} E_n$ *is fast complete, then* $\operatorname{ind} E_n$ *is regular. This result, combined with our Theorem, implies: In a regular LF-space, the sequential and fast completenesses are equivalent.*

References

[1] *M. De Wilde*: Closed Graph Theorem and Webbed Spaces. Pitman, 1978.

[2] *G. Köthe*: Topological Vector Spaces II. Springer-Verlag, 1979.

[3] *W. Robertson*: On the closed graph theorem with webs. Proc. London Math. Soc. *24* (1972), 692–738.

[4] *J. Kučera, C. Bosch*: Bounded sets in fast complete inductive limits. Internat J. Math. *7* (1984), 615–617.

[5] *J. Kučera*: Sequential completeness of LF-spaces. Czechoslovak Math. J *51* (2001), 181–183.

Authors' addresses: C. Bosch, Departamento de Matemáticas ITAM, Rio Hondo No. 1, Col. Tizapan San Angel, 01000 Mexico, e-mail: `bosch@itam.mx`, visiting at Dept. Math. WSU; J. Kučera, Department of Mathematics WSU, Pullman, WA 99164-3113, U.S.A., e-mail: `kucera@math.wsu.edu`.