

Luděk Zajíček

On sets of non-differentiability of Lipschitz and convex functions

*Mathematica Bohemica*, Vol. 132 (2007), No. 1, 75–85

Persistent URL: <http://dml.cz/dmlcz/133997>

## Terms of use:

© Institute of Mathematics AS CR, 2007

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON SETS OF NON-DIFFERENTIABILITY OF  
LIPSCHITZ AND CONVEX FUNCTIONS

LUDĚK ZAJÍČEK, Praha

(Received November 4, 2005)

*Abstract.* We observe that each set from the system  $\tilde{\mathcal{A}}$  (or even  $\tilde{\mathcal{C}}$ ) is  $\Gamma$ -null; consequently, the version of Rademacher's theorem (on Gâteaux differentiability of Lipschitz functions on separable Banach spaces) proved by D. Preiss and the author is stronger than that proved by D. Preiss and J. Lindenstrauss. Further, we show that the set of non-differentiability points of a convex function on  $\mathbb{R}^n$  is  $\sigma$ -strongly lower porous. A discussion concerning sets of Fréchet non-differentiability points of continuous convex functions on a separable Hilbert space is also presented.

*Keywords:* Lipschitz function, convex function, Gâteaux differentiability, Fréchet differentiability,  $\Gamma$ -null sets, ball small sets,  $\delta$ -convex surfaces, strong porosity

*MSC 2000:* 46G05, 26B05

## 1. INTRODUCTION

There exist several infinite-dimensional versions of Rademacher's theorem (on Gâteaux differentiability of Lipschitz functions on separable Banach spaces). They assert that a Lipschitz mapping  $f$  of a separable Banach space to a Banach space with the Radon-Nikodým property is Gâteaux differentiable "almost everywhere". The version due to Aronszajn [1] (and Phelps [11]) says that the set  $NG(f)$  of Gâteaux non-differentiability points of  $f$  is null in Aronszajn's (or, equivalently, in Gaussian) sense; see [3]. Further versions of Rademacher's theorem were proved in [16] ( $NG(f)$  belongs to the class  $\tilde{\mathcal{A}}$ ) and in [8] ( $NG(f)$  is  $\Gamma$ -null). Since  $\tilde{\mathcal{A}}$  is a proper subsystem of the system  $\mathcal{A}$  of Aronszajn's (Gaussian) null sets, the version of [16] is stronger than that of [1]. As shown in [8], the version of [8] is incomparable with that of [1]. In Section 2 we observe (Theorem 2.4) that the lemma ([8, Lemma 2.2]) which was used in [8] for proving that  $NG(f)$  is  $\Gamma$ -null easily implies that each set from  $\tilde{\mathcal{A}}$  (or

from the possibly larger class  $\tilde{\mathcal{C}}$  is  $\Gamma$ -null. Since it is well-known that there exists a  $\Gamma$ -null set which is not in  $\mathcal{A}$  (see Remark 2.5), the version of Rademacher's theorem from [16] is stronger than that of [8].

The above mentioned result that  $\mathcal{A} \setminus \tilde{\mathcal{A}} \neq \emptyset$  was proved (in any separable space  $X$ ) in [16] by a direct construction. We observe (see Corollary 2.6) that Theorem 2.4 together with a deep result of [8] give (if  $X$  is superreflexive) an alternative proof. Namely, we show that the set  $A \in \mathcal{A}$  constructed in [9] is not in  $\tilde{\mathcal{A}}$ .

If  $f$  is a continuous convex function on a separable Banach space  $X$ , then the set  $NG(f)$  of Gâteaux non-differentiability points of  $f$  can be covered by countably many of d.c. (that is, delta-convex) hypersurfaces (see [19, Theorem 2] or [3, Theorem 4.20]); moreover, this result is the optimal one, if we are interested in the smallness of sets  $NG(f)$  (for a continuous convex  $f$ ) only. Note that a full characterization of these sets is given in [10] for  $X = \mathbb{R}^n$ , but even the case of a Hilbert space  $X$  is open. In Section 3 we present some consequences of the above mentioned result of [19] and of P.Hartman's theorem on superposition of delta-convex mappings. In particular, we show that the set  $NG(f)$  is  $\sigma$ -strongly lower porous for each convex function on  $\mathbb{R}^n$ . The infinite-dimensional case is open; see Problem 1 below. If  $X^*$  is separable and  $f$  is a continuous convex function on  $X$ , then ([2]) the set  $NF(f)$  of all Fréchet non-differentiability points is a first category set. This result was strengthened in [14], [15] ( $NF(f)$  is  $\sigma$ -porous; even "angle small"). Moreover, in [8] it is proved that  $NF(f)$  is  $\Gamma$ -null. As written in [7], p. 32, it is difficult to conjecture a precise result on the nature of the sets  $NF(f)$ . In Section 3, we briefly discuss this problem and formulate a related natural Problem 2.

## 2. EACH SET FROM $\tilde{\mathcal{C}}$ IS $\Gamma$ -NULL

First we recall definitions of the systems  $\tilde{\mathcal{A}}, \tilde{\mathcal{C}}$  from [16].

(The symbol  $B(x, r)$  denotes an open ball.)

**Definition 2.1.** Let  $X$  be a separable Banach space. Then we define the following systems:

- (i) If  $0 \neq v \in X$  and  $\varepsilon > 0$ , then  $\tilde{\mathcal{A}}(v, \varepsilon)$  is the system of all Borel sets  $B \subset X$  such that  $\{t: \varphi(t) \in B\}$  is Lebesgue null whenever  $\varphi: \mathbb{R} \rightarrow X$  is such that the function  $x \mapsto \varphi(x) - xv$  has Lipschitz constant at most  $\varepsilon$ .
- (ii)  $\tilde{\mathcal{C}}$  is the system of all Borel sets  $B$  which are of the form  $B = \bigcup_{n=1}^{\infty} B_n$ , where  $B_n \in \tilde{\mathcal{A}}(v_n, \varepsilon_n)$  for some  $v_n \neq 0$  and  $\varepsilon_n > 0$ .

- (iii)  $\tilde{\mathcal{A}}$  is the system of all Borel sets  $B \subset X$  such that, if  $v_i \neq 0$  are vectors from  $X$  with  $\overline{\text{span}}(v_i)_{i=1}^\infty = X$ , then  $B$  can be represented as a union  $\bigcup_{i=1}^\infty \bigcup_{k=1}^\infty B(i, k)$ , where  $B(i, k) \in \tilde{\mathcal{A}}(v_i, 1/k)$  for all  $i$  and  $k$ .

**Remark 2.2.**

- (i) Clearly  $\tilde{\mathcal{A}} \subset \tilde{\mathcal{C}}$  but it is not known whether this inclusion is proper. If  $\mathcal{A}$  denotes the system of all Aronszajn's (equivalently, Gaussian) null sets, then trivially  $\tilde{\mathcal{A}} \subset \mathcal{A}$  and it is not difficult to show that  $\tilde{\mathcal{C}} \subset \mathcal{A}$  (see the beginning of the proof of [16, Proposition 13]).
- (ii) The property of  $\varphi$  from Definition 2.1 (i) is clearly satisfied (see [16, Remark 5]) if  $\varphi$  is Lipschitz and  $\|\varphi'(x) - v\| \leq \varepsilon$  for almost all  $x \in \mathbb{R}$ . Further, a simple extension argument shows that we obtain the same notion of  $\tilde{\mathcal{A}}(v, \varepsilon)$  if we consider  $\varphi$  defined on an arbitrary closed interval (instead on the whole  $\mathbb{R}$ ); see [16], p. 16.

Now we introduce the definition of  $\Gamma$ -null sets from [8], which is based on a sophisticated (but very useful) combination of the notions of a first category set and of a Lebesgue null set. Let  $T := [0, 1]^\mathbb{N}$  be endowed with the product topology and the product Lebesgue measure  $\mu$ . Let  $X$  be a Banach space and let  $\Gamma(X)$  be the space of all continuous mappings  $\gamma: T \rightarrow X$  having continuous partial derivatives  $D_j\gamma$  (with one-sided derivatives at points where the  $j$ -th coordinate is 0 or 1). The set  $\Gamma(X)$  is equipped with the topology generated by the seminorms  $\|\gamma\|_0 = \sup_{t \in T} \|\gamma(t)\|$  and  $\|\gamma\|_k = \sup_{t \in T} \|D_k\gamma(t)\|$ . (This topology is metrizable by a complete separable metric.)

A Borel set  $N \subset X$  is called  $\Gamma$ -null if  $\mu\{t \in T: \gamma(t) \in N\} = 0$  for residually many  $\gamma \in \Gamma(X)$  (i.e., for all  $\gamma$  except a first category set). A possibly non Borel subset of  $X$  is called  $\Gamma$ -null if it is contained in a Borel  $\Gamma$ -null set.

Our observation (Theorem 2.4) is an easy consequence of the following fact which is a special case of [8, Lemma 2.2].

**Lemma 2.3.** *Let  $v \in X$  and  $\varepsilon > 0$ . Then, for residually many  $\gamma \in \Gamma(X)$ , there exist  $k \in \mathbb{N}$  and  $c > 0$  such that  $\sup_{t \in T} \|cD_k\gamma(t) - v\| < \varepsilon$ .*

**Theorem 2.4.** *If  $X$  is separable and  $N \in \tilde{\mathcal{C}}$ , then  $N$  is  $\Gamma$ -null.*

**Proof.** Let  $v \in X$  and  $\varepsilon > 0$  be given. By the definition of  $\tilde{\mathcal{C}}$ , it is sufficient to prove that each  $B \in \tilde{\mathcal{A}}(v, \varepsilon)$  is  $\Gamma$ -null. By Lemma 2.3 there exists a residual set  $S \subset \Gamma(X)$  such that, for each  $\gamma \in S$ , there exist  $k \in \mathbb{N}$  and  $c > 0$  such that

$$(1) \quad \sup_{t \in T} \|cD_k\gamma(t) - v\| < \varepsilon.$$

Choose an arbitrary  $\gamma \in S$  and find  $k \in \mathbb{N}$  and  $c > 0$  such that (1) holds. For each  $s = (s_1, s_2, \dots) \in T$ , set  $p_s(y) := \gamma(s_1, \dots, s_{k-1}, y, s_{k+1}, \dots)$  for  $y \in [0, 1]$  and define  $\varphi_s(x) := p_s(cx)$  for  $x \in [0, 1/c]$ . By (1) we have  $\|\varphi'_s(x) - v\| < \varepsilon$  for each  $x \in [0, 1/c]$ . Consequently, using Remark 2.2 (ii), for each  $B \in \tilde{\mathcal{A}}(v, \varepsilon)$  we obtain that  $(\varphi_s)^{-1}(B)$  is Lebesgue null; consequently also  $(p_s)^{-1}(B) = c \cdot (\varphi_s)^{-1}(B)$  is Lebesgue null. Since  $\gamma$  is continuous and  $B$  is Borel, we can use the Fubini theorem for  $\mu$  and  $\gamma^{-1}(B)$  and obtain  $\mu(\gamma^{-1}(B)) = 0$ , which completes the proof.  $\square$

**Remark 2.5.** As observed in [8], the decomposition result of [9] easily implies that, in the case of an infinite-dimensional superreflexive  $X$ , there exists a  $\Gamma$ -null set which is not in  $\mathcal{A}$  and thus it is not in  $\tilde{\mathcal{A}}$  (not even in  $\tilde{\mathcal{C}}$ ). (Note that clearly  $\tilde{\mathcal{A}} \subset \mathcal{A}$  and also  $\tilde{\mathcal{C}} \subset \mathcal{A}$ ; see Remark 2.2). Consequently, the version of Rademacher's theorem from [16] is stronger than that of [8].

Theorem 2.4 together with a theorem of [8] and the above-mentioned decomposition result of [9] give a new proof of the following known result. (Recall that a set in  $\mathcal{A} \setminus \tilde{\mathcal{A}}$  is constructed in any separable  $X$  in [16]; see Introduction.)

**Corollary 2.6.** *In each separable infinite-dimensional superreflexive space  $X$ , there exists a set in  $\mathcal{A}$  (i.e., a Gaussian null set) which is not in  $\tilde{\mathcal{A}}$  (not even in  $\tilde{\mathcal{C}}$ ).*

**Proof.** By [9],  $X$  can be decomposed into a union of a set  $A \in \mathcal{A}$  and a set  $N$  for which there exists a continuous convex function  $f$  on  $X$  such that  $f$  is Fréchet differentiable at no point of  $N$ . We will show that  $A \notin \tilde{\mathcal{C}}$ . Indeed, suppose that  $A \in \tilde{\mathcal{C}}$ . Then  $A$  is  $\Gamma$ -null by Theorem 2.4. Since  $N$  is also  $\Gamma$ -null by [8, Corollary 3.11], we obtain a contradiction.  $\square$

### 3. EVERY CONVEX FUNCTION ON $\mathbb{R}^n$ IS DIFFERENTIABLE OUTSIDE A $\sigma$ -STRONGLY LOWER POROUS SET

We start with recalling the notion of  $\sigma$ -porosity (for a recent survey about applications of  $\sigma$ -porosity in Banach spaces see [20]).

**Definition 3.1.** Let  $X$  be a Banach space and let  $M \subset X$ ,  $x \in X$ ,  $R > 0$  be given. Then we define  $\gamma(x, R, M)$  as the supremum of all  $r \geq 0$  for which there exists  $z \in X$  such that  $B(z, r) \subset B(x, R) \setminus M$ . Further, define the upper and lower porosity of  $M$  at  $x$  as

$$\bar{p}(M, x) := 2 \limsup_{R \rightarrow 0^+} \frac{\gamma(x, R, M)}{R} \quad \text{and} \quad \underline{p}(M, x) := 2 \liminf_{R \rightarrow 0^+} \frac{\gamma(x, R, M)}{R}.$$

We say that  $M$  is upper (lower, strongly upper, strongly lower) porous at  $x$  if  $\bar{p}(M, x) > 0$  ( $\underline{p}(M, x) > 0$ ,  $\bar{p}(M, x) = 1$ ,  $\underline{p}(M, x) = 1$ ).

We say that  $M$  is upper (lower, strongly upper, strongly lower) porous if  $M$  is upper (lower, strongly upper, strongly lower) porous at each point  $y \in M$ . We say that  $M$  is  $\sigma$ -upper ( $\sigma$ -lower,  $\sigma$ -strongly upper,  $\sigma$ -strongly lower) porous if it is a countable union of upper (lower, strongly upper, strongly lower) porous sets.

Clearly the  $\sigma$ -ideal of all  $\sigma$ -strongly lower porous sets is the smallest from the four just defined  $\sigma$ -ideals.

Another definition we need is that of the notion (see [19], [20]) of d.c. (that is, delta-convex) surfaces of finite codimension. First recall the definition (see [18]) of d.c. mappings.

**Definition 3.2.** Let  $X, Y$  be Banach spaces,  $C \subset X$  an open convex set, and  $F: C \rightarrow Y$  a continuous mapping. We say that  $F$  is d.c. if there exists a continuous convex function  $f: C \rightarrow \mathbb{R}$  such that  $y^* \circ F + f$  is convex whenever  $y^* \in Y^*$ ,  $\|y^*\| \leq 1$ .

**Definition 3.3.** Let  $X$  be a Banach space and  $n \in \mathbb{N}$ ,  $1 \leq n < \dim X$ . We say that  $A \subset X$  is a *d.c. surface* of codimension  $n$  if there exist an  $n$ -dimensional linear space  $F \subset X$ , its topological complement  $E$  and a d.c. (that is, delta-convex) mapping  $\varphi: E \rightarrow F$  such that  $A = \{x + \varphi(x): x \in E\}$ . A d.c. surface of codimension 1 will be called a d.c. hypersurface.

By the smooth part  $A_s$  of the surface  $A$  we mean the set  $A_s = \{x + \varphi(x): x \in S\}$ , where  $S$  is the set of Gâteaux differentiability points of  $\varphi$ .

For the purpose of this paper, we define d.c. surfaces of codimension  $n$  (and their smooth parts) also in the singular case  $n = \dim X < \infty$  as singletons.

**Remark 3.4.**

- (i) Note that, since  $F$  is finite dimensional, it is clear that (cf. [18, Corollary 1.8])  $\varphi$  is a d.c. (delta-convex) mapping if and only if  $y^* \circ \varphi$  is a d.c. function (i.e. the difference of two continuous convex functions) for each  $y^* \in F^*$  (or equivalently: for each  $y^*$  from a fixed basis of  $F^*$ ).
- (ii) It is easy to show (since  $F$  is finite-dimensional and  $\varphi$  is locally Lipschitz) that  $x \in A_s$  if and only if the (Bouligand) tangent cone of  $A$  at  $x$  is a closed linear subspace of  $X$  of codimension  $n$ . Thus  $A_s$  does not depend on the choice of  $E$ ,  $F$  and  $\varphi$ .
- (iii) An easy standard argument shows that each (smooth part of a) d.c. surface of codimension  $n$  ( $n \geq 2$ ) is a subset of a (smooth part of a) d.c. surface of codimension  $n - 1$ .
- (iv) If  $\dim X < \infty$  then, in the definition of  $A_s$ , we can replace Gâteaux differentiability by Fréchet differentiability. Indeed, each d.c. function is locally Lipschitz

and a Lipschitz function on a finite-dimensional space is Fréchet differentiable if and only if it is Gâteaux differentiable. Therefore it is easy to see that each d.c. surface  $A$  of codimension 1 in  $\mathbb{R}^n$  is strongly lower porous at each point of  $A_s$  (and thus  $A_s$  is a strongly lower porous set).

**Lemma 3.5.** *Let  $X$  be a separable Banach space and  $A \subset X$  a d.c. surface of codimension  $k$  ( $1 \leq k < \dim X$ ). Then  $A \setminus A_s$  can be covered by countably many d.c. surfaces of codimension  $k + 1$ .*

*Proof.* Let  $F, E, \varphi: E \rightarrow F$  be as in Definition 3.3. By [19, Theorem 2] (or [3, Theorem 4.20]) there exist sets  $B_i \subset E$  ( $i \in \mathbb{N}$ ) such that each set  $B_i$  is a d.c. surface of codimension 1 in  $E$  and  $\varphi$  is Gâteaux differentiable at each point of  $E \setminus \bigcup B_i$ . Clearly  $A \setminus A_s \subset \bigcup_{i \in \mathbb{N}} T_i$ , where  $T_i := \{e + \varphi(e) : e \in B_i\}$ .

If  $k = \dim X - 1$ , then the assertion of the lemma holds, since all  $B_i$  and  $T_i$  are singletons.

If  $k < \dim X - 1$ , consider an arbitrary  $i \in \mathbb{N}$  and choose a closed linear subspace  $E_1$  of  $E$  of codimension 1,  $F_1 = \text{span}\{f_1\}$  with  $E = E_1 \oplus F_1$  and a d.c. mapping  $\varphi_1: E_1 \rightarrow F_1$  such that  $B_i = \{u + \varphi_1(u) : u \in E_1\}$ . It is easy to see that  $\varphi_2(u) := u + \varphi_1(u)$ ,  $u \in E_1$ , is a locally d.c. mapping from  $E_1$  to  $E$  (cf. [18, Lemma 1.7]). Consequently,  $\varphi_3(u) := \varphi(u + \varphi_1(u))$ ,  $u \in E_1$ , is a locally d.c. mapping from  $E_1$  to  $F$  by [18, Theorem 4.2]. Therefore  $\varphi_4(u) := \varphi_1(u) + \varphi(u + \varphi_1(u))$ ,  $u \in E_1$ , is a locally d.c. mapping from  $E_1$  to  $F_1 \oplus F$ . Using separability of  $E_1$ , Remark 3.4 (i) and a standard extension procedure for continuous convex functions, it is easy to obtain the assertion of the lemma.  $\square$

We will apply the preceding lemma in the case of a finite dimensional  $X$  only; in this case we can use Hartman's superposition theorem [5] for d.c. mappings instead of its generalization [18, Theorem 4.2]. However, the lemma is perhaps of some interest also in the infinite dimensional case.

**Proposition 3.6.** *Let  $X$  be a finite dimensional Banach space and  $A \subset X$  a d.c. surface of codimension  $k$  ( $1 \leq k < \dim X$ ). Then*

- (i)  $A \setminus A_s$  can be covered by countably many smooth parts of d.c. surfaces of codimension  $k + 1$ ,
- (ii)  $A$  can be covered by countably many smooth parts of d.c. surfaces of codimension  $k$ , and
- (iii)  $A$  is  $\sigma$ -strongly lower porous.

*Proof.* Clearly (i) implies (ii) and (ii) implies (iii) (by Remark 3.4 (iii), (iv)).

If  $k = \dim X - 1$  then Lemma 3.5 implies that  $A \setminus A_s$  is countable; so (i) trivially holds.

Now suppose that the proposition holds for some  $1 < k < \dim X$  and  $A$  is a d.c. surface of codimension  $k - 1$ . Then, by Lemma 3.5,  $A \setminus A_s$  can be covered by countably many d.c. surfaces of codimension  $k$  and thus the induction hypothesis (condition (ii)) gives that  $A \setminus A_s$  can be covered by countably many smooth parts of d.c. surfaces of codimension  $k$ . Thus we obtain that the proposition holds for  $k - 1$ . By induction, the proposition follows.  $\square$

Proposition 3.6 and the result of [19] mentioned in Introduction immediately imply the main result of this section (in which, of course, we can write  $NF(f)$  instead of  $NG(f)$ ).

**Corollary 3.7.** *Let  $X$  be a finite dimensional Banach space and  $f$  a continuous convex function on  $X$ . Then the set  $NG(f)$  of all Gâteaux non-differentiability points is  $\sigma$ -strongly lower porous.*

It is not probable that the generalization of Corollary 3.7 holds in an infinite-dimensional space. On the other hand, using [19, Theorem 2] and the fact that every d.c. function is locally Lipschitz, it is easy to see that  $NG(f)$  is  $\sigma$ -lower porous whenever  $f$  is a continuous convex function on a separable Banach space. However, it seems to be probable that the following problem has a negative answer.

**Problem 1.** Let  $f$  be a continuous convex function on  $\ell_2$ . Is it true that then  $NG(f)$  is  $\sigma$ -strongly upper porous?

#### 4. ON THE SET OF POINTS WHERE A CONTINUOUS CONVEX FUNCTION IS NOT FRÉCHET DIFFERENTIABLE

Suppose that  $X$  is an infinite-dimensional Banach space such that  $X^*$  is separable (i.e.,  $X$  is a separable Asplund space) and  $f$  is a continuous convex function on  $X$ . Denote by  $NF(f)$  the set of points at which  $f$  is not Fréchet differentiable. Then  $NF(f)$  is not only of the first category, but it is also an angle small set ([15], see also [12]) and a  $\Gamma$ -null set [8].

Problem 1 of [15] asks whether  $NF(f)$  can be covered by countably many closed convex sets with empty interior and by countably many d.c. hypersurfaces.

This problem was answered in negative by S. V. Konyagin [6] in the case when  $X$  is Hilbert. E. Matoušková [9] observed that the negative answer follows from the main decomposition result of [9] in the case when  $X$  is superreflexive.

The main aim of the present section is to show that now the most natural related problem is probably the following one.

**Problem 2.** Let  $f$  be a continuous convex function on  $\ell_2$ . Is it true that then  $NF(f)$  can be covered by a ball small set and by countably many d.c. hypersurfaces?

Recall the definition of ball small sets:

**Definition 4.1.** Let  $X$  be a Banach space and  $r > 0$ . We say that  $A \subset X$  is  $r$ -ball porous if for each  $x \in A$  and  $\varepsilon \in (0, r)$  there exists  $y \in X$  such that  $\|x - y\| = r$  and  $B(y, r - \varepsilon) \cap A = \emptyset$ . We say that  $A \subset X$  is ball small if it can be written in the form  $A = \bigcup_{n=1}^{\infty} A_n$ , where each  $A_n$  is  $r_n$ -ball porous for some  $r_n > 0$ .

Let us note the following facts:

a) The Hahn-Banach theorem easily implies that, in a Hilbert space, each closed convex set with empty interior is ball small.

b) If  $X$  is Hilbert and  $M \subset X$  is ball small, then there exists (see [15, Theorem 2]) a continuous convex function on  $X$  such that  $M \subset NF(f)$ .

c) Examples in both [6] and [9] give a ball small set which cannot be covered by countably many closed convex sets with empty interior and by countably many d.c. hypersurfaces.

These facts suggest that Problem 2 is very natural. Moreover, we will show that, in this problem, it is not possible to avoid d.c. hypersurfaces. Indeed, in any Hilbert space  $X$  with  $\dim X \geq 2$  there exists a continuous convex  $f$  such that  $NF(f)$  is not ball small. This fact, which is mentioned in [15] (without a proof) for  $X = \mathbb{R}^n$ , follows immediately from Proposition 4.3 below, since for each d.c. hypersurface  $C \subset X$  there exists (by [19, Theorem 2]) a continuous convex function on  $X$  such that  $C \subset NG(f)$ .

Before proving Proposition 4.3 we need to recall some (essentially well-known) facts.

**Definition 4.2.** Let  $X$  be a Banach space,  $M \subset X$ ,  $a \in M$  and  $r > 0$ . We say that  $M$  is  $r$ -ball supported at  $a$  if there exists  $z \in X$  such that  $\|z - a\| \geq r$  and  $B(z, \|z - a\|) \cap M = \emptyset$ . We say that  $M$  is ball supported at  $a$  if it is  $r$ -ball supported at  $a$  for some  $r > 0$ .

Recall (see e.g. [4]) that  $f: (a, b) \rightarrow \mathbb{R}$  is called semiconcave (with a linear modulus) if it is locally of the form  $f(x) = c(x) + Kx^2$ , where  $c(x)$  is concave and  $K > 0$ . A function is called semiconvex, if  $-f$  is semiconcave. Each semiconvex function is a d.c. function (since it is locally d.c., see [5]), but a very special one. For example, the second distributional derivative  $D^2f$  of  $f$  is clearly a signed Radon measure whose negative part is absolutely continuous with respect to the Lebesgue measure.

We can clearly choose a d.c. function  $\varphi: \mathbb{R} \rightarrow \mathbb{R}$  such that, for each open interval  $I$ , neither the positive nor the negative part of  $D^2(\varphi|_I)$  is absolutely continuous with

respect to the Lebesgue measure. Thus, for each open interval  $I$ , the function  $\varphi|_I$  is neither semiconvex nor semiconcave. Then we have that

(\*) there exists a dense subset  $D$  of the graph  $G$  of  $\varphi$  such that  $G$  is ball supported at no point of  $D$ .

Indeed, suppose on the contrary that there exists an open interval  $I$  such that  $G$  is ball supported at each point of  $G \cap (I \times \mathbb{R})$ . The Baire Category Theorem easily implies that there exists an open interval  $J \subset I$ , a dense subset  $S$  of  $G \cap (J \times \mathbb{R})$  and  $r > 0$  such that either

- (a)  $G$  is  $r$ -supported from above at each point  $a \in S$ , or
- (b)  $G$  is  $r$ -supported from below at each point  $a \in S$ .

Here “ $r$ -supported from above” means that the point  $z$  from Definition 4.2 can be chosen to be above  $G$ .

If (a) holds, then an easy compactness argument shows that  $G$  is  $r$ -ball porous from above even at each point of  $G \cap (J \times \mathbb{R})$ . But this implies (see e.g. [17]) that  $\varphi|_J$  is semiconcave, which is a contradiction. Similarly we obtain a contradiction if (b) holds.

**Proposition 4.3.** *In every Hilbert space  $X$  with  $\dim X \geq 2$  there exists a d.c. hypersurface  $C$  which is not ball small.*

*Proof.* Choose a d.c. function  $\varphi: \mathbb{R} \rightarrow \mathbb{R}$  such that (\*) holds.

Further, fix unit orthogonal vectors  $v, w \in X$ , put  $Y := \text{span}\{v, w\}^\perp$  and then define  $C := \{tv + y + \varphi(t)w : t \in \mathbb{R}, y \in Y\}$ . It is clear that  $C$  is a d.c. hypersurface.

To prove that  $C$  is not ball small, suppose on the contrary that  $C = \bigcup_{n=1}^{\infty} P_n$ , where  $P_n$  is  $r_n$ -ball porous. Since each d.c. hypersurface is clearly closed, by the Baire Category Theorem we can choose  $k \in \mathbb{N}$ ,  $c \in C$  and  $\varepsilon > 0$  such that

$$(2) \quad P_k \text{ is dense in } C \cap B(c, \varepsilon).$$

Let  $c = t_0v + y_0 + \varphi(t_0)w$ . Choose  $\delta > 0$  so small that  $c(t, y) := tv + y + \varphi(t)w \in B(c, \varepsilon/4)$  whenever  $t \in I := (t_0 - \delta, t_0 + \delta)$  and  $y \in B(y_0, \delta)$ .

Now consider an arbitrary open interval  $J \subset I$ . Using (2), we can choose  $t^* \in J$  and  $y^* \in B(y_0, \delta)$  such that  $c(t^*, y^*) \in P_k \cap B(c, \varepsilon/4)$ . Set  $r := \min(r_k, \varepsilon/4)$ ; clearly  $P_k$  is  $r$ -ball porous. Therefore, for each  $j \in \mathbb{N}$  with  $1/j < r$ , there exists a point  $z_j = t_jv + y_j + \tau_jw$  such that  $\|z_j - c(t^*, y^*)\| = r$  and  $B(z_j, r - 1/j) \cap P_k = \emptyset$ . Since clearly  $B(z_j, r - 1/j) \subset B(c, \varepsilon)$ , (2) implies  $B(z_j, r - 1/j) \cap C = \emptyset$ .

Since  $(z_j - c(t^*, y_j)) \perp Y$ , we have  $\|z_j - c(t^*, y_j)\| \leq \|(z_j - c(t^*, y_j)) + (y_j - y^*)\| = \|z_j - c(t^*, y^*)\| = r$ .

Putting  $\tilde{z}_j := z_j - y_j$ , we have  $\|\tilde{z}_j - c(t^*, 0)\| = \|z_j - c(t^*, y_j)\| \leq r$  and  $B(\tilde{z}_j, r - 1/j) \cap C = \emptyset$ , since  $C = C - y_j$ .

Identifying  $tv + \tau w$  with  $(t, \tau)$ , we have a sequence  $\tilde{z}_j = (t_j, \tau_j) \in \mathbb{R}^2$  which satisfies  $\|\tilde{z}_j - (t^*, \varphi(t^*))\| \leq r$  and  $B(\tilde{z}_j, r - 1/j) \cap G = \emptyset$  (recall  $G$  is the graph of  $\varphi$ , cf. (\*)). Using a standard compactness argument, we obtain that  $G$  is  $r$ -ball supported at  $(t^*, \varphi(t^*))$ . Since  $J \subset I$  was arbitrary,  $G$  is  $r$ -ball supported at each point of a dense subset of  $G \cap (I \times \mathbb{R})$ . Thus, another simple compactness argument implies that  $G$  is  $r$ -ball supported at each point of  $G \cap (I \times \mathbb{R})$ , which contradicts the choice of  $\varphi$ .  $\square$

**A c k n o w l e d g e m e n t s .** The research was supported by the institutional grant MSM 0021620839 and by the grant GAČR 201/03/0931. The author thanks the Erwin Schrödinger Institute for Mathematical Physics in Vienna for its support.

### References

- [1] *N. Aronszajn*: Differentiability of Lipschitzian mappings between Banach spaces. *Studia Math.* 57 (1976), 147–190. [Zbl 0342.46034](#)
- [2] *E. Asplund*: Fréchet differentiability of convex functions. *Acta Math.* 121 (1968), 31–47. [Zbl 0162.17501](#)
- [3] *Y. Benyamini, J. Lindenstrauss*: Geometric Nonlinear Functional Analysis Vol. I. Colloquium publications (American Mathematical Society) no. 48, Providence, 2000. [Zbl 0946.46002](#)
- [4] *P. Cannarsa, C. Sinestrari*: Semiconcave Functions, Hamilton-Jacobi Equations, and Optimal Control. Birkhäuser, Boston, 2004. [Zbl pre02129788](#)
- [5] *P. Hartman*: On functions representable as a difference of convex functions. *Pacific J. Math.* 9 (1959), 707–713. [Zbl 0093.06401](#)
- [6] *S. V. Konyagin*: On points of existence and nonuniqueness of elements of best approximation. *Theory of Functions and Its Applications*, P.L. Ul'yanov ed., Izdat. Moskov. Univ. 1986, pp. 34–43. (In Russian.)
- [7] *J. Lindenstrauss, D. Preiss*: Fréchet Differentiability of Lipschitz Functions (a survey). *Recent Progress in Functional Analysis*, Elsevier, 2001. [Zbl 1037.46043](#)
- [8] *J. Lindenstrauss, D. Preiss*: On Fréchet differentiability of Lipschitz maps between Banach spaces. *Annals Math.* 157 (2003), 257–288. [Zbl pre01933158](#)
- [9] *E. Matoušková*: An almost nowhere Fréchet smooth norm on superreflexive spaces. *Studia Math.* 133 (1999), 93–99. [Zbl 0923.46016](#)
- [10] *D. Pavlica*: On the points of non-differentiability of convex functions. *Comment. Math. Univ. Carolin.* 45 (2004), 727–734.
- [11] *R. R. Phelps*: Gaussian null sets and differentiability of Lipschitz maps on Banach spaces. *Pacific J. Math.* 18 (1978), 523–531. [Zbl 0396.46041](#)
- [12] *R. R. Phelps*: Convex Functions, Monotone Operators and Differentiability. Second edition, *Lecture Notes in Math.* 1364, Springer, Berlin, 1993. [Zbl 0921.46039](#)
- [13] *D. Preiss*: Differentiability of Lipschitz functions on Banach spaces. *J. Funct. Anal.* 91 (1990), 312–345. [Zbl 0711.46036](#)
- [14] *D. Preiss, L. Zajíček*: Fréchet differentiation of convex functions in a Banach space with a separable dual. *Proc. Amer. Math. Soc.* 91 (1984), 202–204. [Zbl 0521.46034](#)
- [15] *D. Preiss, L. Zajíček*: Stronger estimates of smallness of sets of Fréchet nondifferentiability of convex functions. *Rend. Circ. Mat. Palermo (2), Suppl. no. 3 (1984)*, 219–223. [Zbl 0547.46026](#)
- [16] *D. Preiss, L. Zajíček*: Directional derivatives of Lipschitz functions. *Israel J. Math.* 125 (2001), 1–27. [Zbl 1022.46026](#)

- [17] *Yu. G. Reshetnyak*: On a generalization of convex surfaces. *Mat. Sbornik* 40 (1956), 381–398. (In Russian.) [Zbl 0072.17601](#)
- [18] *L. Veselý, L. Zajíček*: Delta-convex mappings between Banach spaces and applications. *Dissertationes Math. (Rozprawy Mat.)* 289 (1989). [Zbl 0685.46027](#)
- [19] *L. Zajíček*: On the differentiation of convex functions in finite and infinite dimensional spaces. *Czechoslovak Math. J.* 29 (1979), 340–348. [Zbl 0429.46007](#)
- [20] *L. Zajíček*: On  $\sigma$ -porous sets in abstract spaces. *Abstr. Appl. Anal.* 2005 (2005), 509–534. [Zbl pre05038297](#)

*Author's address:* *Luděk Zajíček*, Charles University, Faculty of Mathematics and Physics, Sokolovská 83, 186 75 Praha 8, Czech Republic, e-mail: [zajicek@karlin.mff.cuni.cz](mailto:zajicek@karlin.mff.cuni.cz).