

Ladislav Bican

Non-singular covers over ordered monoid rings

Mathematica Bohemica, Vol. 131 (2006), No. 1, 95–104

Persistent URL: <http://dml.cz/dmlcz/134079>

Terms of use:

© Institute of Mathematics AS CR, 2006

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

NON-SINGULAR COVERS OVER ORDERED MONOID RINGS

LADISLAV BICAN, Praha

(Received November 7, 2005)

Abstract. Let G be a multiplicative monoid. If RG is a non-singular ring such that the class of all non-singular RG -modules is a cover class, then the class of all non-singular R -modules is a cover class. These two conditions are equivalent whenever G is a well-ordered cancellative monoid such that for all elements $g, h \in G$ with $g < h$ there is $l \in G$ such that $lg = h$. For a totally ordered cancellative monoid the equalities $Z(RG) = Z(R)G$ and $\sigma(RG) = \sigma(R)G$ hold, σ being Goldie's torsion theory.

Keywords: hereditary torsion theory, torsion theory of finite type, Goldie's torsion theory, non-singular module, non-singular ring, monoid ring, precover class, cover class

MSC 2000: 16S90, 18E40, 16D80

In what follows, R stands for an associative ring with the identity element and $R\text{-mod}$ denotes the category of all unitary left R -modules. If G is a multiplicative monoid with the unit e , then RG will denote the monoid ring over R consisting of all elements of the form $\sum_{i=1}^n r_i g_i$ with $r_i \in R$, $g_i \in G$, $i = 1, \dots, n$, where the addition is given naturally and the multiplication is given by $\left(\sum_{i=1}^n r_i g_i\right) \left(\sum_{j=1}^m s_j h_j\right) = \sum_{i=1}^n \sum_{j=1}^m r_i s_j g_i h_j$. Recall, that a monoid G is called *left cancellative* if for any three elements $h, g_1, g_2 \in G$ the equality $hg_1 = hg_2$ implies that $g_1 = g_2$. The *right cancellative* monoid is defined similarly and G is called *cancellative* if it is both left and right cancellative. The basic properties of rings and modules can be found in [1].

A class \mathcal{G} of modules is called *abstract*, if it is closed under isomorphic copies. Recall that a *hereditary torsion theory* $\tau_R = (\mathcal{T}_\tau, \mathcal{F}_\tau)$, or simply $\tau = (\mathcal{T}, \mathcal{F})$, for

The work is a part of the research project MSM 0021620839 financed by MSMT and partly supported by the Grant Agency of the Charles University, grant #GAUK 448/2004/B-MAT (301-10-203117).

the category $R\text{-mod}$ consists of two abstract classes \mathcal{T} and \mathcal{F} , the τ -torsion class and the τ -torsionfree class, respectively, such that $\text{Hom}(T, F) = 0$ whenever $T \in \mathcal{T}$ and $F \in \mathcal{F}$, the class \mathcal{T} is closed under submodules, factor-modules, extensions and arbitrary direct sums, the class \mathcal{F} is closed under submodules, extensions and arbitrary direct products and for each module M there exists a short exact sequence $0 \rightarrow T \rightarrow M \rightarrow F \rightarrow 0$ such that $T \in \mathcal{T}$ and $F \in \mathcal{F}$. It is easy to see that every module M contains the unique largest τ -torsion submodule (isomorphic to T), which is called the τ -torsion part of the module M and it is usually denoted by $\tau(M)$. A submodule K of a module M is τ -dense in M if the factor-module M/K is τ -torsion. Associated to each hereditary torsion theory τ is the Gabriel filter \mathcal{L}_τ (or simply \mathcal{L}) of left ideals of R consisting of all the left ideals $I \leq R$ such that $R/I \in \mathcal{T}$. Recall that τ is said to be of *finite type*, if \mathcal{L} contains a cofinal subset of finitely generated left ideals, i.e. if every element of \mathcal{L} contains a finitely generated left ideal of R lying in \mathcal{L} .

For a module M , a submodule K is called *essential* in M , $K \leq' M$ in short, if $K \cap L \neq 0$ for each non-zero submodule L of M and the *singular submodule* $Z(M)$ consists of all elements $a \in M$, the *annihilator left ideal* $(0 : a)_R = \{r \in R; ra = 0\}$, or simply $(0 : a)$, of which is essential in R . Goldie's torsion theory for the category $R\text{-mod}$ is the hereditary torsion theory $\sigma = (\mathcal{T}, \mathcal{F})$, where $\mathcal{T} = \{M \in R\text{-mod}; Z(M/Z(M)) = M/Z(M)\}$ and $\mathcal{F} = \{M \in R\text{-mod}; Z(M) = 0\}$. Note, that throughout this paper the letter σ will always denote Goldie's torsion theory and that the modules from the class \mathcal{F}_σ are usually called *non-singular modules*. A ring R is said to be (*left*) *non-singular* if it is non-singular as a left R -module. For more details on torsion theories we refer to [10] or [9].

If \mathcal{G} is an abstract class of modules, then a homomorphism $\varphi: G \rightarrow M$ with $G \in \mathcal{G}$ is called a \mathcal{G} -precover of the module M , if for each homomorphism $f: F \rightarrow M$ with $F \in \mathcal{G}$ there exists a homomorphism $g: F \rightarrow G$ such that $\varphi g = f$. A \mathcal{G} -precover φ of M is said to be a \mathcal{G} -cover, if every endomorphism f of G with $\varphi f = \varphi$ is an automorphism of the module G . An abstract class \mathcal{G} of modules is called a *precover (cover) class*, if every module has a \mathcal{G} -precover (\mathcal{G} -cover). A more detailed study of precovers and covers can be found in [14].

Recently, in [4; Corollary 3], it has been proved that for each hereditary torsion theory τ with $\tau \geq \sigma$ in the usual sense that $\mathcal{T}_\sigma \subseteq \mathcal{T}_\tau$ the class of all τ -torsionfree modules is a precover class if and only if it is a cover class and these conditions are satisfied exactly when the torsion theory τ is of finite type. Moreover, one of the main results in [5] states that these conditions are equivalent for Goldie's torsion theory for all members of the countable set $\mathfrak{M} = \{R, R/\sigma(R), R[x_1, \dots, x_n], R[x_1, \dots, x_n]/\sigma(R[x_1, \dots, x_n]), n < \omega\}$ of rings whenever they are equivalent for an arbitrary member of this set.

The purpose of this note is to study some relations between the class of non-singular modules in the category $R\text{-mod}$ and that in the category $RG\text{-mod}$, G being a multiplicative monoid. Especially, in Corollary 15 we shall obtain a direct generalization of [5; Theorem 16] dealing with the polynomial rings. The main result may be concentrate in the following Theorem.

Theorem. *Let G be a monoid and let R be an arbitrary ring. Then*

- (i) *if RG is a non-singular ring and the class of all non-singular RG -modules is a cover class, then the class of all non-singular R -modules is a cover class;*
- (ii) *if G is a totally ordered and cancellative monoid, then the equalities $Z(RG) = Z(R)G$ and $\sigma(RG) = \sigma(R)G$ hold;*
- (iii) *if G is a well-ordered cancellative monoid such that for all elements $g, h \in G$ with $g < h$ there is $l \in G$ such that $lg = h$, then the class of all non-singular R -modules is a cover class if and only if the class of all non-singular RG -modules is a cover class.*

Proof. With respect to [4; Corollary 3] it suffices to apply the following Theorems 8, 11 and 14, respectively. □

Now we are ready to start our investigations.

Lemma 1. *If every essential left ideal of the ring R contains a σ -dense finitely generated left ideal, then every left ideal of R contains a σ -dense finitely generated left ideal.*

Proof. Let $0 \neq I \leq R$ be an arbitrary non-essential left ideal of the ring R and let $J \leq R$ be a left ideal of R maximal with respect to $I \cap J = 0$. Then $I \oplus J$ is essential in R and consequently the hypothesis yields the existence of a finitely generated left ideal $K = \sum_{i=1}^n Ra_i$ which is σ -dense in $I \oplus J$ and hence in R . Now $a_i = b_i + c_i$, $b_i \in I$, $c_i \in J$, $i = 1, \dots, n$, and it remains to show that the left ideal $L = \sum_{i=1}^n Rb_i$ is σ -dense in I . If $s \in I$ and $r \in (K : s)$ are arbitrary elements then $rs = \sum_{i=1}^n r_i b_i + \sum_{i=1}^n r_i c_i$ for suitable elements $r_1, \dots, r_n, r \in R$ and consequently $rs = \sum_{i=1}^n r_i b_i \in L$. Thus $(K : s) \subseteq (L : s)$ and so $(L : s) \in \mathcal{L}$, showing that L is σ -dense in I . □

Lemma 2. *The following conditions are equivalent for Goldie's torsion theory σ for the category $R\text{-mod}$:*

- (i) *σ is of finite type;*

- (ii) every left ideal of R contains a σ -dense finitely generated left ideal;
- (iii) every essential left ideal of R contains a σ -dense finitely generated left ideal;
- (iv) every non-singular left ideal of R essentially contains a finitely generated left ideal.

Proof. Obviously, (iii) follows from (ii) trivially, while the converse follows from Lemma 1. Further, if (i) holds then especially (iii) holds and (i) follows from (ii) trivially.

(ii) implies (iv). Let $I \leq R$ be a non-singular left ideal of R . By the hypothesis there is a finitely generated left ideal K of R which is σ -dense in I . Let $J \leq I$ be a left ideal maximal with respect to $K \cap J = 0$. Then $J \cong (J \oplus K)/K \in \mathcal{T} \cap \mathcal{F} = 0$ and so K is essential in I .

(iv) implies (iii). Let I be an essential left ideal of R and let $J \leq I$ be maximal with respect to $(I \cap \sigma(R)) \cap J = 0$. For $J = 0$ we see that $I \cap \sigma(R)$ and hence $\sigma(R)$ is essential in R and the assertion is trivial. In the opposite case J is a non-singular left ideal of R and consequently there is a finitely generated left ideal K of R which is essential in J . Summarizing we have $K \subseteq J \subseteq J \oplus (I \cap \sigma(R)) \subseteq I$ where all the inclusions are obviously σ -dense and we are through. \square

Lemma 3. *Let G be a monoid and let $0 \neq a \in R$ be an arbitrary element. Then $(0 : a)_{RG} = RG(0 : a)_R = (0 : a)_{RG}$.*

Proof. For the sake of simplicity we shall denote by I the left annihilator ideal $(0 : a)_R$ of a in R and by J the left annihilator ideal $(0 : a)_{RG}$ of a in RG . For any element $u = \sum_{i=1}^n r_i g_i \in RG$ and any $r \in I$ we have $ra = 0$, hence $0 = ura = \sum_{i=1}^n (r_i r a) g_i$, which proves the inclusion $RGI \subseteq J$. Conversely, let $u = \sum_{i=1}^n r_i g_i \in J$ be an arbitrary element. Then $0 = ua = \sum_{i=1}^n (r_i a) g_i$ yields $r_i a = 0$ and consequently $r_i \in I$ for each $i = 1, \dots, n$. But this means that $u = \sum_{i=1}^n r_i g_i \in RGI$ and the proof is complete, the rest being obvious. \square

Lemma 4. *Let G be a monoid and let $I \leq R$ be an essential left ideal of the ring R . Then $J = IG = RGI$ is an essential left ideal of the ring RG . Especially, if the left annihilator ideal $(0 : a)_R$ of an element $0 \neq a \in R$ is essential in R , then the left annihilator ideal $(0 : a)_{RG}$ of a is essential in RG .*

Proof. Let $u = \sum_{i=1}^n r_i g_i$ be an arbitrary element of the ring RG with $r_i \neq 0$, $i = 1, \dots, n$, which does not belong to J . If $r_1 \in I$ then we put $s_1 = 1$, while in

the opposite case there is an element $s_1 \in R$ such that $0 \neq s_1 r_1 \in I$. Continuing by the induction let us assume that the elements $s_1, \dots, s_m \in R$, $1 \leq m < n$, such that $s_m \dots s_1 r_i \in I$ for all $i = 1, \dots, m$, and such that at least one of these elements is non-zero, have been already constructed. If $s_m \dots s_1 r_{m+1} \in I$ then we put $s_{m+1} = 1$ and we shall find $s_{m+1} \in R$ such that $0 \neq s_{m+1} s_m \dots s_1 r_{m+1} \in I$ in the opposite case. It is clear now, that after n steps we obtain a non-zero multiple ru which lies in J . The special statement now immediately follows from Lemma 3. \square

Lemma 5. *Let G be a monoid. If I is a left ideal of the ring R such that the left ideal $J = RGI$ is essential in RG , then I is essential in R .*

Proof. Let $0 \neq r \in R$ be an arbitrary element. Then $r = re \in RG$ and consequently there is an element $u = \sum_{i=1}^n r_i g_i \in RG$ such that $0 \neq ur \in J$. Thus there is a non-zero coefficient $r_i r$ of ur , which obviously lies in I and the proof is complete. \square

Proposition 6. *If G is a monoid, then the inclusions $Z(R)G \subseteq Z(RG)$ and $\sigma(R)G \subseteq \sigma(RG)$ hold. Especially, if the ring RG is non-singular, then so is R .*

Proof. If $u = \sum_{i=1}^n r_i g_i$ is an element of $Z(R)G$, then $(0 : r_i)$ is essential in R for each $i = 1, \dots, n$ and consequently the intersection $I = \bigcap_{i=1}^n (0 : r_i)$ is essential in R . By Lemma 4 the left ideal IG is essential in RG and the obvious inclusion $IG \subseteq (0 : u)$ yields that $u \in Z(RG)$, as we wished to show. So, let $u = \sum_{i=1}^n r_i g_i \in \sigma(R)G$ be arbitrary. Then $(Z(R) : r_i) \leq' R$ for each $i = 1, \dots, n$ and consequently $I = \bigcap_{i=1}^n (Z(R) : r_i)$ is essential in R . By Lemma 4 the left ideal IG is essential in RG . For an arbitrary element $v = \sum_{j=1}^m s_j h_j \in IG$ we have $s_j \in I$ and consequently $s_j r_i \in Z(R)$ for all relevant indices i and j . Thus $vu \in Z(R)G$ and so $v \in (Z(R)G : u)$. This means that $IG \subseteq (Z(R)G : u) \subseteq (Z(RG) : u)$, consequently $u \in \sigma(RG)$ and the inclusion $\sigma(R)G \subseteq \sigma(RG)$ is verified. The rest is now clear. \square

Lemma 7. *Let G be a monoid. If I is a left ideal of the ring R such that the left ideal $J = RGI$ essentially contains a finitely generated left ideal of RG , then I essentially contains a finitely generated left ideal of the ring R .*

Proof. By the hypothesis the left ideal J contains a finitely generated left ideal $K = \sum_{i=1}^n RG u_i$, which is essential in J . So, we can write $u_i = \sum_{j=1}^m r_{ij} g_j$ for each

$i = 1, \dots, n$, where some coefficients may be zero. Now we put $L = \sum_{i=1}^n \sum_{j=1}^m Rr_{ij}$ and we are going to verify that L is essential in I . If $r \in I \setminus L$ is an arbitrary element, then especially $r \in J = RGI$ and consequently $0 \neq ur \in K$ for some element $u \in RG$. Thus $ur = \sum_{i=1}^n v_i u_i$ for suitable elements $v_i \in RG$, $i = 1, \dots, n$. Now it is clear that any non-zero coefficient of ur is a left multiple of r and it lies in L . \square

Theorem 8. *Let G be a monoid and let RG be a non-singular ring. If Goldie's torsion theory for the category $RG\text{-mod}$ is of finite type, then Goldie's torsion theory for the category $R\text{-mod}$ is of finite type, too.*

Proof. Since R is non-singular by Proposition 6, the Gabriel filter of Goldie's torsion theory for the category $R\text{-mod}$ consists of essential left ideals, only. So, let I be an arbitrary essential left ideal of the ring R . Then the left ideal $J = RGI$ is essential in RG by Lemma 4 and consequently it essentially contains a finitely generated left ideal of the ring RG by the hypothesis. An application of Lemma 7 yields the existence of a finitely generated left ideal of R , which is essential in I and the proof is therefore complete. \square

Notation. Let G be a totally ordered monoid. If J is a left ideal of the ring RG then we denote by $J[g]$ the set of all coefficients at the element $g \in G$ of all elements $u \in J$ of the form $u = \sum_{i=1}^n r_i g_i$, where $g = g_1 > \dots > g_n$. Note, that this notation is the same as that in [5] which works with "leading" coefficients of the polynomials.

Lemma 9. *Let G be a totally ordered monoid, let J be a left ideal of the ring RG and let $g, h \in G$ be arbitrary elements. Then $J[g]$ is a left ideal of R and if G satisfies the left cancellation law, then $J[g] \subseteq J[hg]$.*

Proof. If $a, b \in J[g]$ and $r \in R$ are arbitrary elements, then there are elements $u = ag + \sum_{i=1}^n a_i g_i$ and $v = bg + \sum_{j=1}^m b_j h_j$ from J such that $g > g_1 > \dots > g_n$ and $g > h_1 > \dots > h_m$. So, $u - v = (a - b)g + \sum_{i=1}^n a_i g_i - \sum_{j=1}^m b_j h_j \in J$, $ru = rag + \sum_{i=1}^n ra_i g_i \in J$ and consequently $a - b, ra \in J[g]$, showing that $J[g]$ is a left ideal of R . Further, $hu = ahg + \sum_{i=1}^n a_i hg_i$, which yields that $a \in J[hg]$ in view of the fact that $hg > hg_1 > \dots > hg_n$ by the left cancellation law for the monoid G . \square

Lemma 10. *Let G be a totally ordered cancellative monoid and let $u = \sum_{k=1}^n r_k g_k$ be a non-zero element of the ring RG such that $g_1 > \dots > g_n$ and $r_k \neq 0$ for each $k = 1, \dots, n$. If K is a left ideal of the ring R such that the left ideal $J = (RGK : u)$ is essential in RG , then the left ideal $I = (K : r_1)$ is essential in R .*

Proof. Proving indirectly let us suppose that there exists a non-zero left ideal L of R such that $L \cap I = 0$. Now RGL is a non-zero left ideal of RG and we are going to show that $RGL \cap J = 0$. Assume, on the contrary, that $v = \sum_{l=1}^m s_l h_l$ is a non-zero element of $RGL \cap J$ such that $h_1 > \dots > h_m$ and $s_l \neq 0$ for each $l = 1, \dots, m$. Note that $h_j g_1 > h_j g_i$ by the left cancellation law and $h_1 g_i > h_j g_i$ by the right cancellation law. Thus $h_1 g_1 > h_j g_i$ for all $j = 1, \dots, m$ and $i = 1, \dots, n$, where at least one of the indices i, j is different from 1. Now $v \in J$ yields $vu = \sum_{k=1}^n \sum_{l=1}^m s_l r_k h_l g_k \in RGK$ and consequently $s_1 r_1 \in K$. On the other hand, $0 \neq s_1 \in L$ means that $s_1 \notin I$, hence $s_1 r_1 \notin K$, which is a contradiction finishing the proof. \square

Theorem 11. *If G is a totally ordered cancellative monoid, then the equalities $Z(RG) = Z(R)G$ and $\sigma(RG) = \sigma(R)G$ hold. Especially, a ring R is non-singular if and only if the ring RG is so.*

Proof. We start with the equality $Z(RG) = Z(R)G$. The inclusion $Z(R)G \subseteq Z(RG)$ holds by Proposition 6. In order to prove the converse let $u = \sum_{k=1}^n r_k g_k \in Z(RG)$ be an arbitrary non-zero element such that $g_1 > \dots > g_n$ and $r_k \neq 0$ for each $k = 1, \dots, n$. Then $(0 : u) \leq' RG$ and so $(0 : r_1) \leq' R$ by Lemma 10. Hence $r_1 \in Z(R)$ yields that $r_1 g_1 \in Z(R)G \subseteq Z(RG)$. Thus $u - r_1 g_1 \in Z(RG)$ and continuing by the induction we finally obtain that $u = \sum_{k=1}^n r_k g_k \in Z(R)G$, as we wished to show.

Now we are going to prove the second equality in the similar way. By Proposition 6 we know that $\sigma(R)G \subseteq \sigma(RG)$. Thus, let $0 \neq u = \sum_{k=1}^n r_k g_k$ be an arbitrary element of $\sigma(RG)$ such that $g_1 > \dots > g_n$ and $r_k \neq 0$ for each $k = 1, \dots, n$. Then $(Z(RG) : u)$ is essential in RG and so the left annihilator ideal $(Z(R) : r_1)$ is essential in R by Lemma 10 in view of the equality $Z(RG) = Z(R)G$ proved in the first part of the proof. Thus $r_1 \in \sigma(R)$ gives that $r_1 g_1 \in \sigma(R)G \subseteq \sigma(RG)$. From this we infer that $u - r_1 g_1 \in \sigma(RG)$ and we can proceed by the induction. Finally we obtain that $u = \sum_{k=1}^n r_k g_k \in \sigma(R)G$, as required. The rest is now obvious. \square

Corollary 12. *Let G be a totally ordered cancellative monoid and let $u = \sum_{k=1}^n r_k g_k$ be a non-zero element of the ring RG . If the left annihilator ideal $(0 : u)$ is essential in RG then the intersection $\bigcap_{k=1}^n (0 : r_k)$ is essential in R .*

Proof. Without loss of generality we may assume that $g_1 > \dots > g_n$. In the proof of Theorem 11 we have shown that $(0 : r_1) \leq' R$ and that $u - r_1 g_1 \in Z(RG)$. Continuing by the induction we shall obtain that $(0 : r_k) \leq' R$ for each $k = 1, \dots, n$, from which the assertion follows immediately. \square

Lemma 13. *Let G be a well-ordered cancellative monoid such that for all $g, h \in G$ with $g < h$ there is $l \in G$ such that $lg = h$. If every essential left ideal of a non-singular ring R essentially contains a finitely generated left ideal, then every essential left ideal of the ring RG essentially contains a finitely generated left ideal.*

Proof. Let J be an essential left ideal of the ring RG . It follows immediately from Lemma 9 that the set $\{J[g]; g \in G\}$ is ordered by the inclusion. Note, that G has not the largest element. Clearly, if $h \in G$ is such that $g \leq h$ for each $g \in G$, then for $g < h$ there is $l \in G$ with $lg = h$. But then the left cancellation law gives $h = lg < lh \leq h$, which is impossible. Let $g_0 \in G$ be the smallest element in the well-order on G . If $J[g_0]$ is not essential in $J[h]$ for each $h \in G$, then let $g_1 \in G$ be the first element of G such that $J[g_0]$ is not essential in $J[g_1]$. Continuing by the induction, after a finite number of steps we shall come to $J[g_k]$ which is essential in $J[h]$ for each $h \geq g_k$. Clearly, in the opposite case we shall construct an infinite sequence $g_0 < g_1 < \dots$ of elements of G such that $J[g_i]$ is not essential in $J[g_{i+1}]$ for each $i < \omega$. Thus there is a non-zero left ideal $L_i \leq J[g_{i+1}]$ such that $J[g_i] \cap L_i = 0$ and so we obtain an infinite direct sum $\bigoplus_{i < \omega} L_i$ of σ -torsionfree left ideals of the ring R . Since Goldie's torsion theory σ for the category $R\text{-mod}$ is of finite type by the hypothesis and Lemma 2, we shall come to a contradiction with [13; Theorem 2.1] stating that σ is of finite type if and only if R contains no infinite direct sum of σ -torsionfree left ideals. We have proved the existence of an element $g \in G$ such that the left ideal $J[g]$ is essential in R and consequently it essentially contains a finitely generated left ideal $K = \sum_{i=1}^r Ra_i$. Now for each a_i there is an element $u_i = a_i g + \sum_{j=1}^s b_{ij} g_j \in J$ such that $g > g_1 > \dots > g_s$ and some coefficients may be zero. Now we put $L = \sum_{i=1}^r RGu_i$ and we are going to show that L is essential in J . So, let $u = \sum_{i=1}^s b_i h_i$ be an element of J such that $h_1 < \dots < h_s$. If $h_s < g$ and if $l' \in G$ is such that $g = l'h_s$ then we can take $l'u$ instead of u and so we may

assume that $h_s \geq g$. Then there is $l \in G$ such that $h_s = lg$ and taking an element $t \in (K : b_s) \setminus (0 : b_s)$ we have $0 \neq tb_s = \sum_{i=1}^r t_i a_i$ and consequently the coefficient of the element $tu - l \sum_{i=1}^r t_i u_i$ at h_s is equal to $tb_s - \sum_{i=1}^r t_i a_i = 0$. Thus we have shown that $tu + u_s = \sum_{j=1}^m c_j k_j$, where $k_1 < \dots < k_m < h_s$, $u_s \in L$ and tu is non-zero. Now we can proceed by the induction. Since G is well-ordered, after a finite number of steps we have to come to zero, from which it immediately follows the existence of a non-zero multiple of u belonging to L and the proof is complete. \square

Theorem 14. *Let G be a well-ordered cancellative monoid such that for all elements $g, h \in G$ with $g < h$ there is $l \in G$ such that $lg = h$. Then Goldie's torsion theory for the category $R\text{-mod}$ is of finite type if and only if Goldie's torsion theory for the category $RG\text{-mod}$ is of finite type.*

Proof. By [5; Theorem 5] Goldie's torsion theory σ for the category $R\text{-mod}$ is of finite type if and only if Goldie's torsion theory for the category $R/\sigma(R)\text{-mod}$ is of finite type. By Theorem 11 we have $\sigma(RG) = \sigma(R)G$ from which we easily obtain the ring isomorphism $(R/\sigma(R))G \cong RG/\sigma(RG)$. So, if Goldie's torsion theory for the category $RG\text{-mod}$ is of finite type, then so is that for the category $RG/\sigma(RG)\text{-mod}$ and consequently that for the category $R/\sigma(R)\text{-mod}$ by Theorem 8. Conversely, if Goldie's torsion theory for the category $R\text{-mod}$ is of finite type then so is that for the category $R/\sigma(R)\text{-mod}$ by [5; Theorem 5] and so that for the category $RG/\sigma(RG)\text{-mod}$ by Lemma 13. Now it remains to use [5; Theorem 5] again. \square

Corollary 15. *If G is an infinite cyclic monoid then Goldie's torsion theory for the category $R\text{-mod}$ is of finite type if and only if Goldie's torsion theory for the category $RG\text{-mod}$ is of finite type.*

Proof. Obvious. \square

Note. It is clear that Theorem 14 holds once we replace the well-order on G into an inverse well-order on G , i.e. into a total order satisfying the maximum condition. Moreover, Corollary 15 says that Goldie's torsion theory for the category $R\text{-mod}$ is of finite type if and only if Goldie's torsion theory for the category $R[x]\text{-mod}$ is of finite type and consequently Theorem 14 generalizes [5; Theorem 16].

References

- [1] *F. W. Anderson, K. R. Fuller*: Rings and Categories of Modules. Graduate Texts in Mathematics, vol. 13, Springer, 1974. [Zbl 0765.16001](#)
- [2] *L. Bican*: Torsionfree precovers. Proceedings of the Klagenfurt Conference 2003 (66. AAA). Verlag Johannes Heyn, Klagenfurt, 2004, pp. 1–6. [Zbl 1074.16002](#)
- [3] *L. Bican*: Precovers and Goldie’s torsion theory. *Math. Bohem.* 128 (2003), 395–400. [Zbl 1057.16027](#)
- [4] *L. Bican*: On torsionfree classes which are not precover classes. To appear.
- [5] *L. Bican*: Non-singular precovers over polynomial rings. To appear.
- [6] *L. Bican, R. El Bashir, E. Enochs*: All modules have flat covers. *Proc. London Math. Society* 33 (2001), 649–652. [Zbl 1029.16002](#)
- [7] *L. Bican, B. Torrecillas*: Precovers. *Czechoslovak Math. J.* 53 (2003), 191–203. [Zbl 1016.16003](#)
- [8] *L. Bican, B. Torrecillas*: On covers. *J. Algebra* 236 (2001), 645–650. [Zbl 0973.16002](#)
- [9] *L. Bican, T. Kepka, P. Němec*: Rings, Modules, and Preradicals. Marcel Dekker, New York, 1982. [Zbl 0483.16026](#)
- [10] *J. Golan*: Torsion Theories. Pitman Monographs and Surveys in Pure and Applied Mathematics, 29, Longman Scientific and Technical, 1986. [Zbl 0657.16017](#)
- [11] *S. H. Rim, M. L. Teply*: On coverings of modules. *Tsukuba J. Math.* 24 (2000), 15–20. [Zbl 0985.16017](#)
- [12] *M. L. Teply*: Torsion-free covers II. *Israel J. Math.* 23 (1976), 132–136. [Zbl 0321.16014](#)
- [13] *M. L. Teply*: Some aspects of Goldie’s torsion theory. *Pacif. J. Math.* 29 (1969), 447–459. [Zbl 0174.06803](#)
- [14] *J. Xu*: Flat Covers of Modules. Lecture Notes in Mathematics 1634, Springer, Berlin, 1996. [Zbl 0860.16002](#)

Author’s address: Ladislav Bican, KA MFF UK, Sokolovská 83, 186 00 Praha 8-Karlín, Czech Republik, e-mail: bican@karlin.mff.cuni.cz.