

Riadh Khennoufa; Olivier Togni

A note on radio antipodal colourings of paths

Mathematica Bohemica, Vol. 130 (2005), No. 3, 277–282

Persistent URL: <http://dml.cz/dmlcz/134100>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON RADIO ANTIPODAL COLOURINGS OF PATHS

RIADH KHENNOUFA, OLIVIER TOGNI, Dijon

(Received November 12, 2004)

Abstract. The radio antipodal number of a graph G is the smallest integer c such that there exists an assignment $f: V(G) \rightarrow \{1, 2, \dots, c\}$ satisfying $|f(u) - f(v)| \geq D - d(u, v)$ for every two distinct vertices u and v of G , where D is the diameter of G . In this note we determine the exact value of the antipodal number of the path, thus answering the conjecture given in [G. Chartrand, D. Erwin and P. Zhang, *Math. Bohem.* 127 (2002), 57–69]. We also show the connections between this colouring and radio labelings.

Keywords: radio antipodal colouring, radio number, distance labeling

MSC 2000: 05C78, 05C12, 05C15

1. INTRODUCTION

Let G be a connected graph and let k be an integer, $k \geq 1$. The distance between two vertices u and v of G is denoted by $d(u, v)$ and the diameter of G by $D(G)$ or simply D . A *radio k -colouring* f of G is an assignment of positive integers to the vertices of G such that

$$|f(u) - f(v)| \geq 1 + k - d(u, v)$$

for every two distinct vertices u and v of G .

Following the notation of [1], [3], we define the *radio k -colouring number* $rc_k(f)$ of a radio k -colouring f of G to be the maximum colour assigned to a vertex of G and the *radio k -chromatic number* $rc_k(G)$ to be $\min\{rc_k(f)\}$ taken over all radio k -colourings f of G .

Radio k -colourings generalize many graph colourings. For $k = 1$, $rc_1(G) = \chi(G)$, the chromatic number of G . For $k = 2$, the radio 2-colouring problem corresponds to the well studied $L(2, 1)$ -colouring problem and $rc_2(G) = \lambda(G)$ (see [5] and references therein). For $k = D(G) - 1$, the radio $(D - 1)$ -colouring is referred to as the *radio*

antipodal colouring, because only antipodal vertices can have the same colour. In that case, $rc_k(G)$ is called the *radio antipodal number*, also denoted by $ac(G)$. Finally, for the case $k = D(G)$, $rc_k(G)$ is called the *radio number* and is studied in [1], [6].

In [2] the antipodal number for cycles was discussed and bounds were given. In [3], Chartrand et al. gave general bounds for the antipodal number of a graph. The authors proved the following result for the radio antipodal number of the path:

Theorem 1 ([3]). *For every positive integer n ,*

$$ac(P_n) \leq \binom{n-1}{2} + 1.$$

Moreover, they conjectured that the above upper bound is the value of the antipodal number of the path. In [4], the authors found a sharper bound for the antipodal number of an odd path (thus showing that the conjecture was false):

Theorem 2 ([4]). *For the path P_n of odd order $n \geq 7$,*

$$ac(P_n) \leq \binom{n-1}{2} - \frac{n-1}{2} + 4.$$

In this note we completely determine the antipodal number of the path:

Theorem 3. *For any $n \geq 5$,*

$$ac(P_n) = \begin{cases} 2p^2 - 2p + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 4p + 5 & \text{if } n = 2p. \end{cases}$$

Notice that for $n = 2p + 1$ we have $\binom{n-1}{2} - \frac{n-1}{2} + 4 = p(2p-1) - p + 4 = 2p^2 - 2p + 4$, thus the bound of Theorem 2 is one from the optimal.

Examples of minimal antipodal colourings of P_7 and P_8 are given in Figure 1.




Figure 1. Antipodal colouring of P_7 and P_8 .

In order to prove Theorem 3, we shall use a result of Liu and Zhu [6] about the radio number of the path. Notice that Liu and Zhu allow 0 to be used as a colour but we do not. Then, when presenting their result, we will make the necessary adjustment (adding “one”) to be consistent with the rest of the paper.

Theorem 4 ([6]). For any $n \geq 3$

$$\text{rc}_{n-1}(P_n) = \begin{cases} 2p^2 + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 2p + 2 & \text{if } n = 2p. \end{cases}$$

2. RADIO k -COLOURINGS

Lemma 1. Let G be a graph of order n and let k be an integer. If f is a radio k -colouring of G then, for any integer $k' > k$, there exists a radio k' -colouring f' of G with $\text{rc}_{k'}(f') \leq \text{rc}_k(f) + (n-1)(k' - k)$.

Proof. We construct a radio k' -colouring f' of G with $\text{rc}_{k'}(f') = c + (n-1)(k' - k)$ from a radio k -colouring f with $\text{rc}_k(f) = c$ in the following way: Let x_1, x_2, \dots, x_n be an ordering of the vertices of G such that $f(x_i) \leq f(x_{i+1})$, $1 \leq i \leq n-1$, and set

$$f'(x_i) = f(x_i) + (i-1)(k' - k).$$

For any two integers i and j , $1 \leq i < j \leq n$, we have $|f'(x_j) - f'(x_i)| = |f(x_j) - f(x_i)| + (j-i)(k' - k)$.

As $|f(x_j) - f(x_i)| \geq 1 + k - d(x_j, x_i)$ and $j - i \geq 1$, we obtain $|f'(x_j) - f'(x_i)| \geq 1 + k + (j-i)(k' - k) - d(x_j, x_i) \geq 1 + k' - d(x_j, x_i)$. Thus f' is a radio k' -colouring of G and $\text{rc}_{k'}(f') = c + (n-1)(k' - k)$. \square

The above result can be strengthened a little in some cases:

Lemma 2. Let G be a graph of order n and let k, k' be integers, $k' > k$. Given a radio k -colouring f of G , let x_1, x_2, \dots, x_n be an ordering of the vertices of G such that $f(x_i) \leq f(x_{i+1})$, $1 \leq i \leq n-1$ and let $\varepsilon_i = |f(x_i) - f(x_{i-1})| - (1 + k - d(x_i, x_{i-1}))$, $2 \leq i \leq n$. Consider a set $I = \{i_1, i_2, \dots, i_s\} \subset \{2, \dots, n\}$, where $1 \leq s \leq n-1$, such that $i_{j+1} > i_j + 1$ for all j , $1 \leq j \leq s-1$. Then there exists a radio k' -colouring f' of G with $\text{rc}_{k'}(f') \leq \text{rc}_k(f) + (n-1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i)$.

Proof. A radio k' -colouring f' of G is obtained simply by setting for all j with $1 \leq j \leq n-1$:

$$f'(x_j) = f(x_j) + (j-1)(k' - k) - \sum_{i \in I, i \leq j} \min(k' - k, \varepsilon_i).$$

The vertex x_n has the maximum colour: $f'(x_n) = f(x_n) + (n-1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i) = \text{rc}_k(f) + (n-1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i)$.

Then, for any two integers j_1 and j_2 , $1 \leq j_1 < j_2 \leq n$, let us show that the condition

$$|f'(x_{j_2}) - f'(x_{j_1})| \geq 1 + k' - d(x_{j_2}, x_{j_1})$$

is verified, i.e. that

$$|f(x_{j_2}) - f(x_{j_1})| + (j_2 - j_1)(k' - k) - \sum_{i \in I, j_1 < i \leq j_2} \min(k' - k, \varepsilon_i) \geq 1 + k' - d(x_{j_2}, x_{j_1}).$$

If $j_2 = j_1 + 1$, then $|f(x_{j_2}) - f(x_{j_1})| = 1 + k - d(x_{j_2}, x_{j_1}) + \varepsilon_{j_2}$. Thus $|f'(x_{j_2}) - f'(x_{j_1})| \geq 1 + k - d(x_{j_2}, x_{j_1}) + \varepsilon_{j_2} + (k' - k) - \min(k' - k, \varepsilon_{j_2}) \geq 1 + k' - d(x_{j_2}, x_{j_1})$.

If $j_2 > j_1 + 1$, then $\sum_{i \in I, j_1 < i \leq j_2} \min(k' - k, \varepsilon_i) \leq (j_2 - j_1 - 1)(k' - k)$ since by the hypothesis there are no two consecutive integers in the set I . Thus $|f'(x_{j_2}) - f'(x_{j_1})| \geq 1 + k - d(x_{j_2}, x_{j_1}) + (j_2 - j_1)(k' - k) - (j_2 - j_1 - 1)(k' - k) = 1 + k' - d(x_{j_2}, x_{j_1})$.

Therefore, f' is a radio k' -colouring of G and $\text{rc}_{k'}(f') = \text{rc}_k(f) + (n - 1)(k' - k) - \sum_{i \in I} \min(k' - k, \varepsilon_i)$. \square

3. ANTIPODAL COLOURINGS OF PATHS

Theorem 3 derives from the next two theorems.

Theorem 5. For any $n \geq 5$,

$$\text{ac}(P_n) \leq \begin{cases} 2p^2 - 2p + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 4p + 5 & \text{if } n = 2p. \end{cases}$$

Proof. The fact that $\text{ac}(P_5) = 7$ is easily checked (see [3]). Thus take $n \geq 6$ and let $P_n = (u_1, u_2, \dots, u_n)$. We consider two cases depending on whether n is even or odd.

Case 1. $n = 2p + 1$ is odd for an integer $p \geq 3$. Define a colouring f of P_{2p+1} by

$$\begin{cases} f(u_1) = 3p + 2, \\ f(u_2) = p + 1, \\ f(u_i) = i(2p - 1) - p + 3, & 3 \leq i \leq p, \\ f(u_{p+1}) = 2p + 2, \\ f(u_{p+2}) = 1, \\ f(u_{p+i}) = i(2p - 1) - 2p + 3, & 3 \leq i \leq p, \\ f(u_{2p+1}) = p + 2. \end{cases}$$

Then the vertex u_p has the maximum colour: $f(u_p) = p(2p-1)-p+3 = 2p^2-2p+3$. We only have to show that the distance condition is verified for two vertices u_i and u_{p+j} , $3 \leq i, j \leq p$ (the other cases can be easily checked). We want

$$\begin{aligned} |f(u_{p+j}) - f(u_i)| &\geq 1 + (D - 1) - d(u_{p+j}, u_i) \Leftrightarrow \\ |j(2p - 1) - 2p + 3 - (i(2p - 1) - p + 3)| &\geq 2p - (p + j - i) \Leftrightarrow \\ |(j - i)(2p - 1) - p| &\geq p - j + i. \end{aligned}$$

If $j - i \geq 1$ then $|(j - i)(2p - 1) - p| = (j - i)(2p - 1) - p \geq 2p - 1 - p = p - 1 \geq p - j + i$.
If $j - i < 1$ then $|(j - i)(2p - 1) - p| = -(j - i)(2p - 1) + p = (i - j)(2p - 1) + p \geq p - j + i$ for $p \geq 1$.

Case 2. $n = 2p$ is even for an integer $p \geq 3$. Define a colouring f of P_{2p} by

$$\begin{cases} f(u_1) = p, \\ f(u_i) = (p - i)(2p - 1) + 2, & 2 \leq i \leq p - 1, \\ f(u_p) = 2p^2 - 4p + 5, \\ f(u_{p+i}) = 2p^2 - 4p + 6 - f(u_{p-i+1}), & 1 \leq i \leq p. \end{cases}$$

Then the vertex u_p has the maximum colour: $f(u_p) = 2p^2 - 4p + 5$. We only have to show that the distance condition is verified for two vertices u_i and u_{p+j} , $2 \leq i \leq p - 1, 1 \leq j \leq p$ (the other cases can be easily checked). We want

$$\begin{aligned} |f(u_{p+j}) - f(u_i)| &\geq 1 + (D - 1) - d(u_{p+j}, u_i) \Leftrightarrow \\ |(p - j)(2p - 1) + 3 - ((p - i)(2p - 1) - p + 2)| &\geq 2p - 1 - (p + j - i) \Leftrightarrow \\ |(i - j)(2p - 1) + p + 1| &\geq p - j + i - 1. \end{aligned}$$

If $i - j \geq 0$ then $|(i - j)(2p - 1) + p + 1| = (i - j)(2p - 1) + p + 1 \geq p - j + i - 1$ since $(i - j)(2p - 2) \geq -1$ for $p \geq 1$.

If $i - j < 0$, i.e. if $j - i \geq 1$ then $|(i - j)(2p - 1) + p + 1| = (j - i)(2p - 1) - p - 1 \geq p - j + i - 1$ since $2p(j - i) \geq 2p$. \square

Theorem 6. For any $n \geq 5$,

$$\text{ac}(P_n) \geq \begin{cases} 2p^2 - 2p + 3 & \text{if } n = 2p + 1, \\ 2p^2 - 4p + 5 & \text{if } n = 2p. \end{cases}$$

Proof. For $n = 2p + 1$, by Lemma 1 we have $\text{rc}_{n-1}(P_n) \leq \text{ac}(P_n) + (n - 1)$. This together with Theorem 4 gives $\text{ac}(P_{2p+1}) \geq 2p^2 + 3 - 2p$.

For $n = 2p$, let $D = D(P_{2p}) = 2p - 1$. We will use Lemma 2 with the radio $(D - 1)$ -colouring f of P_{2p} described in the proof of Theorem 5 and with $k = D - 1 = 2p - 1$ and $k' = D = 2p$. Keeping the notation of Lemma 2, one can see that f is such that $x_1 = u_{p+1}$, $x_2 = u_1$, $x_3 = u_{2p-1}$, $x_4 = u_{p-1}$, \dots , $x_{2j+1} = u_{2p-j+1}$, $x_{2j} = u_{p-j+1}$, \dots , $x_{2p-1} = u_{2p}$, $x_{2p} = u_p$. Thus ε_3 verifies

$$\begin{aligned} \varepsilon_3 &= |f(x_3) - f(x_2)| - (1 + k - d(x_3, x_2)) \\ &= |f(u_{2p-1}) - f(u_1)| - (1 + 2p - 2 - (2p - 2)) \\ &= |2p^2 - 4p + 6 - f(u_2) - f(u_1)| - 1 \\ &= |2p^2 - 4p + 6 - (p - 2)(2p - 1) - 2 - p| - 1 = 1. \end{aligned}$$

A similar calculus gives $\varepsilon_{2p-1} = 1$ and $\varepsilon_i = 0$ for all other indices.

Thus, as $k' - k = 1$ and $p \geq 3$, applying Lemma 2 with $I = \{3, 2p - 1\}$ gives

$$\text{rc}_{2p-1}(P_{2p}) \leq \text{ac}(P_{2p}) + (2p - 1) - \varepsilon_3 - \varepsilon_{2p-1},$$

that is

$$\text{ac}(P_{2p}) \geq \text{rc}_{2p-1}(P_{2p}) - (2p - 1) + \varepsilon_3 + \varepsilon_{2p-1}.$$

By virtue of Theorem 4 we obtain $\text{ac}(P_{2p}) \geq 2p^2 - 2p + 2 - (2p - 1) + 1 + 1 = 2p^2 - 4p + 5$. □

References

- [1] *G. Chartrand, D. Erwin, F. Harary, P. Zhang*: Radio labelings of graphs. *Bull. Inst. Combin. Appl.* 33 (2001), 77–85.
- [2] *G. Chartrand, D. Erwin, P. Zhang*: Radio antipodal colorings of cycles. *Congr. Numerantium* 144 (2000), 129–141.
- [3] *G. Chartrand, D. Erwin, P. Zhang*: Radio antipodal colorings of graphs. *Math. Bohem.* 127 (2002), 57–69.
- [4] *G. Chartrand, L. Nebeský, P. Zhang*: Radio k -colorings of paths. *Discuss. Math. Graph Theory* 24 (2004), 5–21.
- [5] *D. Kuo, J.-H. Yan*: On $L(2, 1)$ -labelings of Cartesian products of paths and cycles. *Discrete Math.* 283 (2004), 137–144.
- [6] *D. Liu, X. Zhu*: Multi-level distance labelings for paths and cycles. To appear in *SIAM J. Discrete Math.*

Author's address: Riadh Khennoufa, Olivier Togni, LE2I, UMR 5158 CNRS, Université de Bourgogne, BP 47870, 21078 Dijon cedex, France, e-mail: olivier.togni@u-bourgogne.fr.