

I. Gelbukh

On the structure of a Morse form foliation

*Czechoslovak Mathematical Journal*, Vol. 59 (2009), No. 1, 207–220

Persistent URL: <http://dml.cz/dmlcz/140474>

## Terms of use:

© Institute of Mathematics AS CR, 2009

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## ON THE STRUCTURE OF A MORSE FORM FOLIATION

I. GELBUKH, Mexico City

(Received May 6, 2007)

*Abstract.* The foliation of a Morse form  $\omega$  on a closed manifold  $M$  is considered. Its maximal components (cylinders formed by compact leaves) form the foliation graph; the cycle rank of this graph is calculated. The number of minimal and maximal components is estimated in terms of characteristics of  $M$  and  $\omega$ . Conditions for the presence of minimal components and homologically non-trivial compact leaves are given in terms of  $\text{rk } \omega$  and  $\text{Sing } \omega$ . The set of the ranks of all forms defining a given foliation without minimal components is described. It is shown that if  $\omega$  has more centers than conic singularities then  $b_1(M) = 0$  and thus the foliation has no minimal components and homologically non-trivial compact leaves, its foliation graph being a tree.

*Keywords:* number of minimal components, number of maximal components, compact leaves, foliation graph, rank of a form

*MSC 2010:* 57R30, 58K65

## 1. INTRODUCTION AND ANNOUNCEMENT OF THE RESULTS

Consider a connected closed oriented manifold  $M$  with a Morse form  $\omega$ , i.e., a closed 1-form with Morse singularities—locally the differential of a Morse function. The set of its singularities  $\text{Sing } \omega$  is finite. This form defines a foliation  $\mathcal{F}_\omega$  on  $M \setminus \text{Sing } \omega$ . Its leaves  $\gamma$  can be classified into compact, *compactifiable* ( $\gamma \cup \text{Sing } \omega$  is compact), and non-compactifiable.

Such foliations have remarkably regular structure. A connected component  $\mathcal{C}_i^{\max}$  of the union of compact leaves—which we call *maximal component*—is an open cylinder over any its leaf, whose levels are leaves. In particular, all leaves in a maximal component are diffeomorphic. A connected component  $\mathcal{C}_i^{\min}$  of the union of non-compactifiable leaves is called *minimal component*. Its topology can be arbitrarily complex—say, such a component can cover the whole  $M \setminus \text{Sing } \omega$  [1]—but it cannot be too simple: a minimal component contains at least two cycles with non-

commensurable integrals [10]. Each non-compactifiable leaf is dense in its minimal component [7]. The boundary of a maximal or minimal component consists of a finite number of non-compact compactifiable leaves  $\gamma_i^0$  and singularities. This gives a decomposition of  $M$  into a finite number of non-intersecting pieces

$$M = \left( \bigcup \mathcal{C}_i^{\max} \right) \cup \left( \bigcup \mathcal{C}_i^{\min} \right) \cup \left( \bigcup \gamma_i^0 \right) \cup \text{Sing } \omega,$$

interconnected in a way resembling the structure of a simplicial complex.

The way the pieces are put together is described by the *foliation graph*  $\Gamma$ —a connected graph (allowing loops and multiple edges) whose edges are  $\mathcal{C}_i^{\max}$  and vertices are connected components of  $M \setminus \bigcup \mathcal{C}_i^{\max}$ . This notion has been used for studying foliation structure [2], [6]. Note that unlike [10], we define the graph on the whole  $M$ , including  $\text{Sing } \omega$ .

In this paper, we show (Theorem 2.1) that

$$m(\Gamma) = c(\omega),$$

where  $m(\Gamma)$  is the cycle rank of the foliation graph  $\Gamma$  and  $c(\omega)$  is the number of homologically independent compact leaves of  $\mathcal{F}_\omega$ .

The number  $m(\omega)$  of minimal components is bounded by the first Betti number:  $2m(\omega) \leq b_1(M)$  [1], [8]. We obtain a stronger estimate (Theorem 3.1):

$$2m(\omega) + c(\omega) \leq b_1(M)$$

and an independent estimate (our main theorem, Theorem 3.2):

$$m(\omega) + c(\omega) \leq h(M),$$

where  $h(M)$  is the maximum rank of a subgroup in  $H^1(M, \mathbb{Z})$  with trivial cup-product [11]. There are practical methods of calculating  $h(M)$  (Remark 3.1).

We also estimate (Theorem 3.3) the total number of components for a singular form:

$$M(\omega) + m(\omega) \leq h(M) + |\text{Sing } \omega| - 1,$$

where  $M(\omega)$  is the number of maximal components (obviously, for a non-singular form  $M(\omega) + m(\omega) = 1$  instead). In addition, for a singular form

$$M(\omega) + 2m(\omega) \leq b_1(M) + |\text{Sing } \omega| - 1.$$

In addition to the bound for  $m(\omega) + c(\omega)$  above, we present some conditions for  $m(\omega) = 0$  and for  $c(\omega) = 0$ . A foliation having no minimal components is called

*compactifiable*. In [3] we have presented some conditions for compactifiability of the foliation in terms of the structure of  $\ker[\omega] \subseteq H_1(M)$ , where  $[\omega]$  is the integration map. Here we consider other characteristics of the form: the form's  $\text{rk } \omega \stackrel{\text{def}}{=} \text{rk im}[\omega]$  and the number of singularities of different indices.

If  $\omega$  is rational ( $\text{rk } \omega \leq 1$ ) then  $\mathcal{F}_\omega$  is compactifiable [13]. We show that the converse is true only in the sense that any compactifiable foliation *can* be defined by a rational Morse form. Namely, a compactifiable foliation  $\mathcal{F}_\omega$  is defined by forms  $\omega'$  with

$$(0 \text{ or } 1) \leq \text{rk } \omega' \leq c(\omega)$$

(Theorem 4.1); whether 0 is included depends on the structure of the directed foliation graph [4]. In particular, if  $\text{rk } \omega > h(M)$  then  $\mathcal{F}_\omega$  is not compactifiable. For instance, if  $\text{rk } \omega = b_1(M)$  (i.e. maximal) and the cup-product on  $H^1(M, \mathbb{Z})$  is non-trivial then  $\mathcal{F}_\omega$  is non-compactifiable (Corollary 4.3). If in addition  $H_1(M)$  has no torsion and the cup-product is non-degenerate then all compact leaves of  $\mathcal{F}_\omega$  are homologically trivial (Proposition 4.1).

No necessary condition for compactifiability of  $\mathcal{F}_\omega$  can be obtained in terms of  $\text{Sing } \omega$ : for any foliation there exists a rational Morse form with the same set of singularities of each index [14]. However,  $\text{Sing } \omega$  can give useful information on  $M$ . If  $\text{Sing } \omega = \emptyset$  then  $M$  is a bundle over  $S^1$  [15], so the topology of  $\mathcal{F}_\omega$  is defined by  $\text{rk } \omega$ : if  $\text{rk } \omega = 1$  then  $\mathcal{F}_\omega$  is compact (all leaves are compact), otherwise it is minimal. If all singularities are centers then  $M = S^n$  and  $\mathcal{F}_\omega$  is obviously compact. (By centers, called also spherical singularities, we mean those of index 0 or  $\dim M$ .)

We generalize the latter fact: If there are more centers than conic singularities then (Theorem 4.2)

$$b_1(M) = 0;$$

in particular, in this case  $m(\omega) + c(\omega) = 0$ , i.e.,  $\mathcal{F}_\omega$  is compactifiable, all its leaves are homologically trivial, and its foliation graph is a tree (Theorem 4.3).

The paper is organized as follows. In Section 2 we recall some facts about qualitative structure of a Morse form foliation and give the necessary definitions. We introduce minimal and maximal foliation components and describe their properties (Section 2.1). Then we define the foliation graph and calculate its cycle rank (Section 2.2). In Section 3 we give the upper bounds on the number of minimal components (Section 3.1) and on the total number of (minimal and maximal) components of the foliation (Section 3.2). Finally, in Section 4 we give some conditions on the presence of minimal components in terms of the rank of the form (Section 4.1) and the indices of its singularities (Section 4.2).

## 2. GENERAL STRUCTURE

The general structure of a Morse form foliation has been studied in [1], [3], [7], [10].

**Definition 2.1** [3]. A leaf  $\gamma \in \mathcal{F}_\omega$  is called compactifiable if  $\gamma \cup \text{Sing } \omega$  is compact; otherwise it is called non-compactifiable.

Note that compact leaves are compactifiable.

The set covered by all non-compactifiable leaves is open [7]. A compact leaf  $\gamma$  has an open neighborhood consisting solely of compact leaves [1], [2]: indeed, integrating  $\omega$  gives  $f$  with  $df = \omega$  near  $\gamma$ ; hence the set covered by all compact leaves is also open. The number of non-compact compactifiable leaves  $\gamma_i^0$  is finite since each singularity belongs to the closure of no more than four non-compact compactifiable leaves (see Figure 1); thus such leaves are isolated.




Figure 1. A conic singularity has locally four (a) or two (b), (c) adjacent (non-compact) leaves; centers have no such leaves, and other singularities have one (d). Here, (d) and (c) visualize non-compact leaves for  $\dim M = 4$  (the lowest dimension where singularities other than conic or centers exist) as connected components of the set  $\{\mathbb{R}^4 \setminus p \mid x^2 + y^2 - z^2 = \pm t^2\}$ , respectively, foliated by the “time”  $t$ .

**2.1. Minimal and maximal foliation components.** We call *foliation components* the connected components  $\mathcal{C}_i$  of the set  $M \setminus (\bigcup \gamma_i^0 \cup \text{Sing } \omega)$ , i.e. the union of all compact and non-compactifiable leaves. Each component  $\mathcal{C}_i$  is open, and  $\partial \mathcal{C}_i \subseteq \bigcup \gamma_j^0 \cup \text{Sing } \omega$ . For any  $\gamma_j^0$  there exist either one or two components  $\mathcal{C}_i$  such that  $\gamma_j^0 \cap \partial \mathcal{C}_i \neq \emptyset$ ; see Figure 2(a). Thus the number of components is finite.

A component consists entirely of leaves of one type: either non-compactifiable or compact. Components of the former kind are *minimal neighborhoods* consisting of non-compactifiable leaves; each such leaf is dense in its minimal component [1], [7]:

**Definition 2.2** ([1]). A connected component of the union of all non-compactifiable leaves is called a *minimal component* of the foliation.




Figure 2. (a) The leaf  $\gamma_2^0$  adjoins only one component, while  $\gamma_1^0$  connects two components. (b) The corresponding foliation graph has a loop.

Components of the latter kind are *maximal* connected sets consisting of compact leaves:

**Definition 2.3** ([3]). A connected component of the union of all compact leaves is called a *maximal component* of the foliation.

A maximal component  $\mathcal{C}$  of a singular Morse form foliation is cylindrical:  $\mathcal{C} = \gamma \times (0, 1)$ ,  $\gamma \times x \in \mathcal{F}_\omega$ , where  $\gamma$  is any (compact) leaf in  $\mathcal{C}$ . Note that for a non-singular form a maximal component—which exists iff  $\omega$  is rational—is a bundle over  $S^1$  with fiber  $\gamma \in \mathcal{F}_\omega$  and covers the whole manifold [15].

While a maximal component is topologically simple, the topology of a minimal component can be arbitrarily complex (e.g., it can cover the whole  $M \setminus \text{Sing } \omega$  [1]). Moreover, it cannot be too simple: a minimal component contains at least two (homologically independent in  $M$ ) 1-cycles with incommensurable periods [10].

Thus  $M$  can be decomposed into a finite number of non-intersecting pieces interconnected in a way resembling the structure of a simplicial complex:

$$(2.1) \quad M = \left( \bigcup \mathcal{C}_i^{\max} \right) \cup \left( \bigcup \mathcal{C}_j^{\min} \right) \cup \left( \bigcup \gamma_k^0 \right) \cup \text{Sing } \omega,$$

with maximal components  $\mathcal{C}_i^{\max}$  being cylindrical and minimal components  $\mathcal{C}_j^{\min}$  having at least two independent 1-cycles. For examples of such a decomposition see Figures 2 (a), 3 (a, c), and 4 (b).

**2.2. Foliation graph.** The configuration formed by the maximal components in the decomposition (2.1) is described by the *foliation graph* [3]. Rewrite (2.1) as

$$M = \left( \bigcup \mathcal{C}_i^{\max} \right) \cup \left( \bigcup P_j \right),$$

where  $P_j$  are connected components of the union  $P$  of all non-compact leaves and singularities. Since a maximal component is a cylinder,  $\partial \mathcal{C}_i^{\max} \subseteq P$  consists of one or




Figure 3. (a), (c) Examples of the decomposition. (b) The vertices of  $\Gamma$  can include singularities, non-compact compactifiable leaves and (d) whole minimal components.

two connected components; thus each  $\mathcal{C}_i^{\max}$  adjoins one or two of the  $P_j$ . This allows representing  $M$  as a connected graph  $\Gamma$  (admitting multiple edges and loops) with edges  $\mathcal{C}_i^{\max}$  and vertices  $P_j$ : an edge  $\mathcal{C}_i^{\max}$  is incident to a vertex  $P_j$  if  $\partial\mathcal{C}_i^{\max} \cap P_j \neq \emptyset$ ; see Figures 2 and 3.

Let  $c(\omega) = \text{rk } H_\omega$ , where  $H_\omega \subseteq H_{n-1}(M)$  is generated by all compact leaves, i.e., the number of homologically independent compact leaves. Denote by  $m(\Gamma)$  the cycle rank of the graph  $\Gamma$ .

**Theorem 2.1.** *It holds*

$$(2.2) \quad c(\omega) = m(\Gamma).$$

Moreover, there exist  $\gamma_1, \dots, \gamma_{c(\omega)}$  such that for any compact leaf  $\gamma$  it holds

$$[\gamma] = \sum_{i \in I_\gamma} \pm [\gamma_i].$$

**Proof.** In  $H_\omega$  there exists a basis  $e$  consisting of homology classes of leaves  $[\gamma_1], \dots, [\gamma_c]$ ,  $c = c(\omega)$  [3]. Each  $\gamma_i$ ,  $i = 1, \dots, c$ , defines a maximal component  $\mathcal{C}_i^{\max} = \gamma_i \times (0, 1)$ , which is an edge  $h_i$  in the foliation graph  $\Gamma$ ; see Figure 4(a).

Consider the subgraph  $\Gamma^0 = \Gamma \setminus \bigcup_{i=1}^c h_i$ ; see Figure 4(b). It has the same set of vertices. Independence of  $e$  implies that  $\Gamma^0$  is connected.

Suppose that  $\Gamma^0$  contains a cycle  $z$ ; associate with it a curve  $\alpha \subset M$ . Obviously,  $\alpha \cap \gamma_i = \emptyset$  for all  $i$ . Let  $h \in z$  be an edge, and  $\gamma \subset M$  a corresponding leaf. Then  $[\alpha] \cdot [\gamma] \neq 0$  and  $[\alpha] \cdot [\gamma_i] = 0$ , which contradicts the maximality of  $e$ .




Figure 4. Proof of Theorem 2.1.

Thus  $\Gamma^0$  is a spanning tree. Since the cycle rank of a graph equals to the number of chords of its spanning tree [5], we have  $m(\Gamma) = c(\omega)$ . What is more, since removing any edge splits a tree, for any compact leaf  $\gamma$  we have  $[\gamma] = \sum_{i \in I_\gamma} \pm [\gamma_i]$ ; see Figure 4(c).  $\square$

**Corollary 2.1.** *If all compact leaves of  $\mathcal{F}_\omega$  are homologically trivial then  $\Gamma$  is a tree. If in addition  $\mathcal{F}_\omega$  has no minimal components then  $\omega = df$ .*

The second statement follows from Corollary 4.1 below.

### 3. NUMBER OF COMPONENTS

Given a specific  $M$ , we can estimate the number of components of Morse form foliations on  $M$ . This may give useful information on their structure.

**3.1. Number of minimal components.** Denote by  $m(\omega)$  the number of minimal components of  $\mathcal{F}_\omega$ . It is known that  $2m(\omega) \leq b_1(M)$  [1], [8]. We generalize this estimate:

**Theorem 3.1.**  $2m(\omega) + c(\omega) \leq b_1(M)$ .

*Proof.* Let  $z_1, \dots, z_{c(\omega)}$  be 1-cycles dual to homologically independent compact leaves  $\gamma_1, \dots, \gamma_{c(\omega)}$ , i.e.,  $z_i \cdot [\gamma_j] = \delta_{ij}$ . Each minimal component  $C_i^{\min}$  contains two cycles  $z'_i, z''_i$  with incommensurable periods [10]. Suppose that

$$(3.1) \quad \sum_{i=1}^{m(\omega)} (n'_i z'_i + n''_i z''_i) + \sum_{j=1}^{c(\omega)} m_j z_j = 0.$$

Intersecting (2) with  $[\gamma_k]$  gives  $m_k = 0$  for all  $k$ . Thus for any  $p$  we can rewrite (3.1) as

$$(3.2) \quad n'_p z'_p + n''_p z''_p = - \sum_{i \neq p} (n'_i z'_i + n''_i z''_i) = z.$$

On the one hand,  $z$  is induced from  $\mathcal{C}_p^{\min}$  and on the other hand, from  $M \setminus \mathcal{C}_p^{\min}$ , thus by the Mayer-Vietoris sequence it is induced from  $\partial\mathcal{C}_p^{\min}$ . Since  $\partial\mathcal{C}_p^{\min}$  consists of leaves and singularities,  $\int_z \omega = 0$ . Thus  $n'_p = n''_p = 0$ , since the corresponding periods are incommensurable.  $\square$

**Example 3.1.** For a connected sum  $M = (S^2 \times S^1) \# (S^2 \times S^1)$ , Theorem 3.1 gives  $2m(\omega) + c(\omega) \leq 2$ . This characterizes the topology of  $\mathcal{F}_\omega$ . Indeed, if  $\mathcal{F}_\omega$  is compactifiable, then it has at most two homologically independent compact leaves. Otherwise, it has exactly one (uniquely ergodic [1]) minimal component, and all its compact leaves are homologically trivial.

Denote by  $h(M)$  the maximum rank of a subgroup in  $H^1(M, \mathbb{Z})$  with trivial cup-product [11].

**Remark 3.1.** Some methods of calculating  $h(M)$  can be found in [12]:

- (i)  $h(M_1 \times M_2) = \max\{h(M_1), h(M_2)\}$ .
- (ii)  $h(M_1 \# M_2) = h(M_1) + h(M_2)$  for  $\dim M_i \geq 2$ .
- (iii) Let  $b_i = b_i(M)$  be the Betti numbers and  $r = \text{rk ker } \smile$ , where  $\smile$  is the cup-product on  $H^1(M, \mathbb{Z})$ . Then

$$\frac{b_1 + b_2 r}{b_2 + 1} \leq h(M) \leq \frac{b_1 b_2 + r}{b_2 + 1}.$$

In particular, if  $b_2 = 1$  then  $h(M) = \frac{1}{2}(b_1 + r)$ ; if  $r = b_1$  then  $h(M) = b_1$ .

- (iv) If  $\smile$  is surjective then  $h(M) \leq r + \frac{1}{2} + \sqrt{(b_1 - r - \frac{1}{2})^2 - 2b_2}$ .

**Example 3.2.**  $h(T^n) = 1$  (torus);  $h(M_g^2) = g$ .

**Theorem 3.2.**  $m(\omega) + c(\omega) \leq h(M)$ .

*Proof.* (i) Let us show that for a minimal component  $\mathcal{C}^{\min}$  there exist  $u \in H_1(\mathcal{C}^{\min})$  and  $z \in H_{n-1}(\mathcal{C}^{\min})$  such that their intersection  $u \cdot z \neq 0$ .

Indeed, consider the diagram

$$\begin{array}{ccccccc}
\longrightarrow & H_1(\partial\bar{\mathcal{C}}) & \xrightarrow{i_*} & H_1(\bar{\mathcal{C}}) & \xrightarrow{j_*} & H_1(\bar{\mathcal{C}}, \partial\bar{\mathcal{C}}) & \longrightarrow \\
& & & \uparrow \varphi_* & & \uparrow D & \\
& & & H_1(\mathcal{C}) & & H^{n-1}(\mathcal{C}, \mathbb{Z}) & \\
& & & \uparrow D & & \uparrow \varphi_* & \\
\longrightarrow & H^{n-1}(\bar{\mathcal{C}}, \partial\bar{\mathcal{C}}, \mathbb{Z}) & \xrightarrow{j_*} & H^{n-1}(\bar{\mathcal{C}}, \mathbb{Z}) & \xrightarrow{i_*} & H^{n-1}(\partial\bar{\mathcal{C}}, \mathbb{Z}) & \longrightarrow
\end{array}$$

where  $\mathcal{C} = \mathcal{C}^{\min}$ , the lines are exact sequences of pairs,  $D$  is the Poincaré isomorphism defined by the cap-product, and the homomorphisms  $\varphi_*$  and  $\varphi^*$  are induced by the inclusion  $\varphi: \mathcal{C} \hookrightarrow \bar{\mathcal{C}}$ .

Let  $c \subset \mathcal{C}$  be a closed curve such that  $\int_c \omega \neq 0$  and  $u = [c]$ . Then  $k\varphi_*u \notin \text{im } i_*$  for any  $k \in \mathbb{Z}$  and  $j_*\varphi_*u \neq 0$ ; let  $\alpha = D^{-1}j_*\varphi_*u \neq 0$ . Since  $\alpha \in H^{n-1}(\mathcal{C}, \mathbb{Z})$  is of infinite order, it can be viewed as an element of  $\text{Hom}(H_{n-1}(\mathcal{C}), \mathbb{R})$ . So there exists a cycle  $z \in H_{n-1}(\mathcal{C})$  such that  $\alpha(z) \neq 0$ . By construction,  $u \cdot z \neq 0$ .

(ii) For each minimal component  $\mathcal{C}_i = \mathcal{C}_i^{\min}$  consider the cycles  $u_i \in H_1(M)$ ,  $z_i \in H_{n-1}(M)$  such that  $u_i \cdot z_i \neq 0$ .

Consider a maximal system  $\{\gamma_j\}$  of homologically independent compact leaves. The system  $\{z_i\} \cup \{[\gamma_j]\}$  is independent. Indeed, suppose that

$$\sum p_i z_i + \sum q_j [\gamma_j] = 0.$$

Since  $u_k \cdot [\gamma_j] = 0$  and  $u_k \cdot z_i = 0$  for all  $i \neq k$ , we have all  $p_i = 0$ . Then all  $q_j = 0$  since  $[\gamma_j]$  are linearly independent.

Finally, on  $\{\mathcal{D}z_i\} \cup \{\mathcal{D}[\gamma_j]\}$ , where  $\mathcal{D}: H_{n-1}(M) \rightarrow H^1(M, \mathbb{Z})$  is the Poincaré duality, the cup-product  $\smile$  is trivial, since all  $\mathcal{C}_i$  and  $\gamma_j$  do not intersect.  $\square$

**Example 3.3.** For a torus  $T^n$ ,  $m(\omega) + c(\omega) \leq 1$ . We have all the conclusions of Example 3.1 (with an obvious correction of its last sentence).

**Remark 3.2.** The bound on  $m(\omega)$  implied by Theorem 3.1 and its corollary

$$(3.3) \quad m(\omega) \leq \frac{1}{2}b_1(M)$$

is independent of the bound implied by Theorem 3.2 and its corollary

$$(3.4) \quad m(\omega) \leq h(M),$$

as the following examples show. Which one is stronger depends on the structure of the cup-product.

**Example 3.4.** For a torus, we have  $h(T^n) = 1$  and  $b_1(T^n) = n$ . For  $n = 2, 3$ , the bounds (3.3) and (3.4) are equivalent. If  $n \geq 4$ , even the weakened form (3.4) of Theorem 3.2 gives a stronger bound on  $m(\omega)$  than Theorem 3.1.

**Example 3.5.** For  $M = \#_{i=1}^p (S^2 \times S^1)$ ,  $p \geq 2$ , we have  $h(M) = b_1(M) = p$ . Theorem 3.1 gives a stronger bound than Theorem 3.2 and (3.3) stronger than (3.4). Moreover, for a minimal foliation on this  $M$ , (3.3) gives a stronger bound than Theorem 3.2. We leave open the question of existence of manifolds on which (3.3) gives a stronger bound than Theorem 3.2 for *any* foliation.

**3.2. Total number of components.** Denote by  $M(\omega)$  the number of maximal components.

**Theorem 3.3.** *If  $\text{Sing } \omega \neq \emptyset$  then*

$$(3.5) \quad M(\omega) + m(\omega) \leq h(M) + |\text{Sing } \omega| - 1;$$

$$(3.6) \quad M(\omega) + 2m(\omega) \leq b_1(M) + |\text{Sing } \omega| - 1.$$

**Proof.** In the foliation graph  $\Gamma$  with  $M(\omega)$  edges and  $p$  vertices,  $M(\omega) = m(\Gamma) + p - 1$ . Obviously,  $p \leq |\text{Sing } \omega|$ . By Theorem 2.1 and Theorem 3.2,  $m(\Gamma) = c(\omega) \leq h(M) - m(\omega)$ , which gives (3.5); similarly, Theorem 3.1 gives (3.6).  $\square$

**Example 3.6.** For a torus  $T^n$  and  $S^n \times S^1$ , (3.5) gives  $M(\omega) + m(\omega) \leq |\text{Sing } \omega|$ ; for  $M_g^2$ ,  $M(\omega) + m(\omega) \leq g + |\text{Sing } \omega| - 1$ .

The estimates (3.5) and (3.6) are independent; cf. Remark 3.2 and the examples therein.

For a non-singular form, obviously,  $M(\omega) + m(\omega) = 1$ . Note that for such forms, (3.5) does not hold, for instance, on  $T^n$  and (3.6) does not hold, for instance, on  $S^n \times S^1$ .

#### 4. PRESENCE OF MINIMAL COMPONENTS AND HOMOLOGICALLY NON-TRIVIAL LEAVES

A foliation without minimal components is called *compactifiable*. We will present some conditions for compactifiability of  $\mathcal{F}_\omega$  and the presence of homologically non-trivial leaves in it (i.e., whether the foliation graph is a tree) in terms of  $\text{rk } \omega$  and  $\text{Sing } \omega$ .

**4.1. Conditions in terms of the rank of the form.** Consider the form's *rank*  $\text{rk } \omega = \text{rk im } [\omega]$ , where  $[\omega]: H_1(M) \rightarrow \mathbb{R}$  is the integration map. If  $\omega$  is *rational*

( $\text{rk } \omega \leq 1$ ) then  $\mathcal{F}_\omega$  is compactifiable, i.e.  $m(\omega) = 0$  [13]. We will show that the converse is true in the sense that any compactifiable foliation *can* be defined by a rational form; more specifically, the ranks of forms defining such a foliation cover a limited range starting from 0 or 1.

Recall that a maximal component  $\mathcal{C}_j^{\max}$  is  $\gamma_j \times (0, 1)$  for some leaf  $\gamma_j$  and  $M(\omega)$  is the number of maximal components.

**Theorem 4.1.** *Let  $\mathcal{F}_\omega$  be a compactifiable Morse form foliation and  $r \geq 1$ . Then there exists a Morse form of rank  $r$  defining the same foliation iff  $r \leq c(\omega)$ .*

Whether there exists such a form of rank 0 depends on the structure of the directed foliation graph [4].

*Proof.* It is known that

$$H_1(M) = \left\langle D[\gamma_1], \dots, D[\gamma_{M(\omega)}], i_* H_1 \left( M \setminus \bigcup_{j=1}^{M(\omega)} \mathcal{C}_j^{\max} \right) \right\rangle,$$

where  $D$  is a Poincaré duality map (i.e.,  $D[\gamma_k] \cdot [\gamma_l] = \delta_{kl}$ ) and  $i$  is the inclusion: indeed, the cycles  $z \notin \langle D[\gamma_i] \rangle$  in Figure 5 can be moved out of  $\bigcup \mathcal{C}_i^{\max}$ , cf. (2.1); see a detailed proof in [3].




Figure 5. A cycle  $z$  such that  $z \cdot [\gamma_i] = 0$  can be moved out of  $\mathcal{C}_i^{\max}$ .

So  $\text{rk } \omega$  is defined by  $\langle D[\gamma_k] \rangle$ , which by Theorem 2.1 has a basis  $D[\gamma_1], \dots, D[\gamma_{c(\omega)}]$ . Let us realize each  $D[\gamma_i]$  by a closed curve  $\alpha_i$ ; by construction  $\alpha_j \cap \mathcal{C}_i^{\max} = \emptyset$  iff  $i \neq j$ . Without loss of generality we can suppose that  $\alpha_i \cap \mathcal{C}_i^{\max}$  is connected and transverse to the leaves.

Consider on  $M$  a smooth function  $f(x) > 0$  constant on leaves in each cylinder  $\mathcal{C}_i^{\max}$  and equal to 1 outside  $\bigcup \mathcal{C}_i^{\max}$ . The form  $\omega' = f(x)\omega$  defines the same foliation  $\mathcal{F}_\omega$ . By choosing an appropriate  $f(x)$  we can vary its integrals along  $\alpha_i$ ,  $i = 1, \dots, c(\omega)$ , obtaining any desired  $\text{rk } \omega'$ .  $\square$

**Example 4.1.** The compactifiable foliation on  $M_g^2$  (or  $\#_{i=1}^g T^n$ ) shown in Figure 6 can be defined by a rational form with  $\text{rk } \omega = 1$ . It can also be defined by forms with  $\text{rk } \omega = 2, \dots, g$ .




Figure 6. (a) The foliation from Example 4.1 and (b) its foliation graph.

**Corollary 4.1.** *For a compactifiable foliation  $\mathcal{F}_\omega$  it holds  $\text{rk } \omega \leq c(\omega)$ .*

**Corollary 4.2.** *Any compactifiable Morse form foliation can be defined by a rational form.*

On the other hand, in any cohomology class with  $\text{rk } \omega > 1$  there exists a form defining a minimal foliation [1]. What is more:

**Corollary 4.3.** *If  $\text{rk } \omega > h(M)$  then  $\mathcal{F}_\omega$  has a minimal component [3]. In particular, if  $\text{rk } \omega = b_1(M)$  (maximal) and the cup-product  $\smile$  on  $H^1(M, \mathbb{Z})$  is non-trivial then  $\mathcal{F}_\omega$  has a minimal component.*

**Proposition 4.1.** *If  $\text{rk } \omega = b_1(M)$ ,  $H_1(M)$  has no torsion, and  $\smile$  is non-degenerate, then  $c(\omega) = 0$ , i.e., all compact leaves are homologically trivial.*

Note that under these conditions, harmonic Morse forms have no compact leaves [2].

**Proof.** Suppose that for some compact leaf  $\gamma$  it holds  $[\gamma] \neq 0$ . We can construct a (non-Morse) form  $\varphi$  trivial outside a cylindrical neighborhood of  $\gamma$ , in which  $\mathcal{F}_\varphi = \mathcal{F}_\omega$ ; thus  $\varphi \wedge \omega = 0$ . Since  $[\gamma] \neq 0$ , the form  $\varphi$  can be chosen such that  $[\varphi] \in H^1(M, \mathbb{Z})$ . Thus  $[\varphi] \smile_{\mathbb{R}} [\omega] = \sum \alpha_i ([\varphi] \smile \xi_i) = 0$ , where  $\smile_{\mathbb{R}}$  is the cup-product on  $H^1(M, \mathbb{R})$  and  $\xi_i$  is a basis in  $H^1(M, \mathbb{Z})$ . Since  $H^2(M, \mathbb{Z})$  has no torsion,  $[\varphi] \smile \xi_i = 0$  for all  $i$ , which contradicts the non-degenerateness of  $\smile$ .  $\square$

#### 4.2. Conditions in terms of the form's singularities.

In Section 3, we have shown that  $m(\omega) + c(\omega) \leq h(M)$ . We will show that if there are more centers than conic singularities, then  $m(\omega) + c(\omega) = 0$ , i.e.,  $\mathcal{F}_\omega$  is compactifiable and all its compact leaves are homologically trivial.

No necessary condition for compactifiability of  $\mathcal{F}_\omega$  can be obtained in terms of  $\text{Sing } \omega$ : for any  $\omega$  there exists a rational Morse form with the same singularities with

their indices [14], [15]; its foliation is compactifiable. However,  $\text{Sing } \omega$  can define the topology of  $M$ .

If  $\text{Sing } \omega = \emptyset$  then  $M$  is a bundle over  $S^1$  [15]. The topology of  $\mathcal{F}_\omega$  is defined by  $\text{rk } \omega$ : if  $\text{rk } \omega = 1$  then  $\mathcal{F}_\omega$  is compact (all its leaves are compact), otherwise it is minimal.

If  $\text{Sing } \omega \neq \emptyset$  but all its singularities are centers then  $M = S^n$  and  $\mathcal{F}_\omega$  is compact. We will generalize this fact: if there are more centers than conic singularities then  $b_1(M) = 0$ .

Denote by  $\Omega_k = \Omega_k(\omega)$ ,  $k \leq \frac{n}{2}$ , the set of singularities of index  $k$  or  $n - k$ . Let  $\mu_i = \mu_i(\omega)$  be the number of singularities of index  $i$ , then  $|\Omega_k| = \mu_k + \mu_{n-k}$  for  $k \neq \frac{n}{2}$  and  $|\Omega_{\frac{n}{2}}| = \mu_{\frac{n}{2}}$ .

**Theorem 4.2.** *If  $|\Omega_0| > |\Omega_1|$  then  $b_1(M) = 0$ .*

*P r o o f.* (i) If  $|\Omega_0| > |\Omega_1|$  then  $\omega = df$ .

Indeed, there exists a rational Morse form with the same singularities and indices [14]; its foliation is compactifiable. Without loss of generality we can assume that each its leaf adjoins at most one singularity [6]. Thus its foliation graph  $\Gamma$  has no vertices  $P$  with  $\deg P > 3$ ; the singularity in any vertex with  $\deg P = 3$  belongs to  $\Omega_1$ , and the set of vertices with  $\deg P = 1$  is  $\Omega_0$ .

For a connected graph,  $2m(\Gamma) = \sum (k - 2)p_k + 2 \geq 0$ , where  $m(\Gamma)$  is the cycle rank and  $p_k$  is the number of vertices with  $\deg P = k$ ;  $p_1 = |\Omega_0|$ ,  $p_3 \leq |\Omega_1|$ . Thus  $|\Omega_0| > |\Omega_1|$  implies  $m(\Gamma) = 0$ ; by Theorems 2.1 and 4.1,  $\omega = df$ .

(ii) If  $\dim M \geq 3$  and  $\omega = df$  then

$$(4.1) \quad |\Omega_0| - |\Omega_1| \leq 2 - 2b_1(M).$$

Indeed, consider the Morse inequality

$$F(\lambda) = \sum_{k=0}^{\lambda} (-1)^{\lambda-k} (\mu_k(f) - b_k(M)) \geq 0,$$

which turns into equality for  $\lambda = n$ , where  $n = \dim M$ . Applying the last inequality to  $F(1) + F(n - 2) - F(n)$  gives

$$-(\mu_0 - b_0) + (\mu_1 - b_1) + (\mu_{n-1} - b_{n-1}) - (\mu_n - b_n) \geq 0,$$

which gives (4.1) since  $|\Omega_0| = \mu_0(f) + \mu_n(f)$  and  $|\Omega_1| = \mu_1(f) + \mu_{n-1}(f)$ .

(iii) On  $M_g^2$  it holds  $|\Omega_0| - |\Omega_1| = 2 - b_1(M)$ , which together with (ii) finishes the proof.  $\square$

Now, Theorem 3.1 gives:

**Theorem 4.3.** *If  $|\Omega_0| - |\Omega_1| > 0$  then  $\mathcal{F}_\omega$  is compactifiable ( $m(\omega) = 0$ ), all its compact leaves are homologically trivial ( $c(\omega) = 0$ ), and the foliation graph is a tree.*

#### References

- [1] *P. Arnoux and G. Levitt:* Sur l'unique ergodicité des 1-formes fermées singulières. *Invent. Math.* *84* (1986), 141–156.
- [2] *M. Farber, G. Katz and J. Levine:* Morse theory of harmonic forms. *Topology* *37* (1998), 469–483.
- [3] *I. Gelbukh:* Presence of minimal components in a Morse form foliation. *Diff. Geom. Appl.* *22* (2005), 189–198.
- [4] *I. Gelbukh:* Ranks of collinear Morse forms. Submitted.
- [5] *F. Harary:* Graph theory. Addison-Wesley Publ. Comp., Massachusetts, 1994.
- [6] *K. Honda:* A note on Morse theory of harmonic 1-forms. *Topology* *38* (1999), 223–233.
- [7] *H. Imanishi:* On codimension one foliations defined by closed one forms with singularities. *J. Math. Kyoto Univ.* *19* (1979), 285–291.
- [8] *A. Katok:* Invariant measures of flows on oriented surfaces. *Sov. Math. Dokl.* *14* (1973), 1104–1108.
- [9] *G. Levitt:* 1-formes fermées singulières et groupe fondamental. *Invent. Math.* *88* (1987), 635–667.
- [10] *G. Levitt:* Groupe fondamental de l'espace des feuilles dans les feuilletages sans holonomie. *J. Diff. Geom.* *31* (1990), 711–761.
- [11] *I. Mel'nikova:* A test for non-compactness of the foliation of a Morse form. *Russ. Math. Surveys* *50* (1995), 444–445.
- [12] *I. Mel'nikova:* Maximal isotropic subspaces of skew-symmetric bilinear map. *Vestnik MGU* *4* (1999), 3–5.
- [13] *S. Novikov:* The Hamiltonian formalism and a multivalued analog of Morse theory. *Russian Math. Surveys* *37* (1982), 1–56.
- [14] *A. Pashitnov:* The incidence coefficients in the Novikov complex are generically rational functions. *Sankt-Petersbourg Math. J.* *9* (1998), 969–1006.
- [15] *D. Tischler:* On fibering certain foliated manifolds over  $S^1$ . *Topology* *9* (1970), 153–154.

*Author's address:* I. Gelbukh, CIC-IPN, 07738, DF, Mexico, e-mail: [gelbukh@member.ams.org](mailto:gelbukh@member.ams.org).