

Acta Universitatis Palackianae Olomucensis. Facultas Rerum  
Naturalium. Mathematica

---

Raad J. K. al Lami

Holomorphically projective mappings of compact semisymmetric manifolds

*Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica*, Vol. 49 (2010), No. 1, 49--53

Persistent URL: <http://dml.cz/dmlcz/140736>

**Terms of use:**

© Palacký University Olomouc, Faculty of Science, 2010

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

# Holomorphically Projective Mappings of Compact Semisymmetric Manifolds

Raad J. K. Al LAMY

*Basic Education College, Misan University, Iraq*  
*e-mail: raadjaka@yahoo.com*

(Received February 10, 2010)

## Abstract

In this paper we consider holomorphically projective mappings from the compact semisymmetric spaces  $A_n$  onto (pseudo-) Kählerian spaces  $\bar{K}_n$ . We proved that in this case space  $A_n$  is holomorphically projective flat and  $\bar{K}_n$  is space with constant holomorphic curvature. These results are the generalization of results by T. Sakaguchi, J. Mikeš, V. V. Domashev, N. S. Sinyukov, E. N. Sinyukova, M. Škodová, which were done for holomorphically projective mappings of symmetric, recurrent and semisymmetric Kählerian spaces.

**Key words:** Holomorphically projective mapping, equiaffine space, affine-connected space, semisymmetric space, Riemannian space, Kählerian space.

**2000 Mathematics Subject Classification:** 53B20, 53B30, 53B35

## 1 Introduction

Diffeomorphisms and automorphisms of geometrically generalized spaces constitute one of the current main directions in differential geometry. A large number of papers are devoted to geodesic, quasigeodesic, almost geodesic, holomorphically projective and other mappings (see [1], [6], [8], [9], [10], [14], [16], [18], [20], [21], [22], [23]). On the other hand, one line of thought is now the most important one, namely, the investigation of special affine-connected, Riemannian, Kählerian and Hermitian spaces.

As we know, Kählerian spaces are the special case of Hermitian spaces [23]. In many papers, holomorphically projective mappings and transformations of Hermitian spaces are studied (for example see [1], [3], [4], [10], [12], [15], [16], [13], [17], [19], [20], [21], [23]). These are special cases of  $F_1$ -planar mappings.

In [8], [10],  $F_1$ -planar mappings from the space  $A_n$  with affine connection onto a Riemannian space  $\bar{V}_n$  are defined and studied.

In this paper, we present some new results obtained for holomorphically projective mappings from compact equiaffine semisymmetric  $A_n$  onto Kählerian spaces  $\bar{K}_n$ .

## 2 Holomorphically projective mappings

J. Mikeš and O. Pokorná [12] considered holomorphically projective mappings from equiaffine spaces  $A_n$  onto Kählerian spaces  $\bar{K}_n$ .

A space  $A_n$  with the affine connection  $\Gamma$  is *equiaffine* if in  $A_n$  the Ricci tensor is symmetric. A (pseudo-) Riemannian space  $\bar{K}_n$  is called a *Kählerian space* if it contains, along with the metric tensor  $\bar{g}$ , an affine structure  $F$  satisfying the following relations

$$F^2 = -Id, \quad \bar{g}(X, FY) + \bar{g}(Y, FX) = 0, \quad \bar{\nabla}F = 0,$$

where  $X, Y \in T\bar{K}_n$ ,  $\bar{\nabla}$  is the connection of  $\bar{K}_n$ .

The following criteria from the paper [12] hold for holomorphically projective mappings from an equiaffine space  $A_n$  onto a Kählerian space  $\bar{K}_n$ .

Consider concrete mappings  $f: A_n \rightarrow \bar{K}_n$ , both spaces being referred to the general coordinate system  $x$  with respect to this mapping. This is a coordinate system where two corresponding points  $M \in A_n$  and  $f(M) \in \bar{K}_n$  have equal coordinates  $x = (x^1, x^2, \dots, x^n)$ ; the corresponding geometric objects in  $\bar{K}_n$  will be marked with a bar. For example,  $\nabla$  and  $\bar{\nabla}$  are connections on  $A_n$  and  $\bar{K}_n$ , respectively.

The equiaffine space  $A_n$  admits a *holomorphically projective mapping*  $f$  onto the Kählerian space  $\bar{K}_n$  if and only if the following conditions in the common coordinate system  $x$  hold

$$\bar{\nabla}(X, Y) = \nabla(X, Y) + \psi(X)Y + \psi(Y)X - \psi(FX)FY - \psi(FY)FX.$$

where  $\psi$  is a linear form. We suppose that  $\psi$  is a gradient form, i.e.  $\psi = d\Psi$ . In this space  $A_n$  there is a complex structure  $F$  covariantly constant.

If  $\psi \neq 0$  then a holomorphically projective mapping is called *nontrivial*; otherwise it is said to be *trivial* or *affine*.

Hereafter we shall assume that in the equiaffine space  $A_n$  the Ricci tensor *Ric with respect to the structure  $F$  will be preserved*, i.e.  $Ric(X, Y) = Ric(FX, FY)$ .

In this case  $\bar{P} = P$  holds, where

$$P_{ijk}^h = R_{ijk}^h - \frac{1}{n+2} (\delta_k^h R_{ij} - \delta_j^h R_{ik} + (F_j^h R_{\alpha k} - F_k^h R_{\alpha j}) F_i^\alpha + 2 F_i^h R_{\alpha k} F_j^\alpha)$$

is the *tensor of the holomorphically projective curvature* of  $A_n$ . This tensor is an invariant object of the holomorphically projective mappings, see [10], [20], [21], [23]. This tensor is  $F$ -traceless [5].

If an equiaffine space  $A_n$  with condition  $P_{ijk}^h = 0$  (*holomorphically projective flat space*) admits a holomorphically projective mapping onto a Kählerian space  $\bar{K}_n$ , then  $\bar{K}_n$  has the constant holomorphic curvature [12].

### 3 Holomorphically projective mappings from semisymmetric equiaffine spaces

It is known, *semisymmetric* spaces  $A_n$  are characterized by the condition  $R \circ R = 0$  [2]. These spaces were characterized by N. S. Sinyukov [20] by the following differential conditions on the Riemannian tensor:  $R_{ijk,[lm]}^h = 0$ . On base of the Ricci identity, this condition is written as follows

$$R_{\alpha jk}^h R_{ilm}^\alpha + R_{i\alpha k}^h R_{jtm}^\alpha + R_{ij\alpha}^h R_{klm}^\alpha - R_{ijk}^\alpha R_{\alpha lm}^h = 0.$$

We have the following [6]:

**Theorem 1** *If an equiaffine semisymmetric space  $A_n$ , where the Ricci tensor  $R_{ij}$  with respect to the structure  $F$  will be preserved, admits a holomorphically projective mapping onto a Kählerian space  $\bar{K}_n$ , then  $\bar{K}_n$  is the space with the constant holomorphically projective curvature or  $\bar{K}_n$  admits a  $K$ -concircular vector field  $\psi_i$ , which satisfies*

$$\nabla_Y \psi(X) - \psi(X)\psi(Y) + \psi(FX)\psi(FY) = \frac{\Delta}{n} \bar{g}(X, Y), \quad \Delta = \text{const.} \quad (3.1)$$

### 4 Holomorphically projective mappings from compact semisymmetric equiaffine spaces

We have the following:

**Theorem 2** *If a compact equiaffine semisymmetric space  $A_n$ , where the Ricci tensor  $R_{ij}$  with respect to the structure  $F$  will be preserved, admits a holomorphically projective mapping onto a Kählerian space  $\bar{K}_n$  then  $\bar{K}_n$  is the space with the constant holomorphically projective curvature (and  $K_n$  is holomorphically projective) or mapping is trivial (affine).*

**Proof** Let  $A_n$  be a compact semisymmetric space of the type described above which admits a holomorphically projective mapping onto a Kählerian space  $\bar{K}_n$ .

According to the Theorem 1, we have formulas (3.1) which may be written on all coordinate neighbourhood  $(x)$  in the followig form

$$\psi_{i,j} - \psi_i \psi_j + \psi_\alpha \psi_\beta F_i^\alpha F_j^\beta = \frac{\Delta}{n} \bar{g}_{hi}, \quad (4.1)$$

where “,” is denote covariant derivative with respect to the connection  $\nabla$  on  $A_n$ ,  $\psi_i$ ,  $F_i^h$  and  $\bar{g}_{ij}$  are components of form  $\psi$ , structure  $F$  and metric tensor  $\bar{g}$ , respectively.

Let us suppose that there exists some solution of the equation (4.1), let be denoted by  $\varphi$ . Using  $\varphi$  we obtain the following formulation of these equations:

$$\psi_{ij} \equiv \psi_{i,j} - \psi_i \varphi_j + \psi_\alpha \varphi_\beta F_i^\alpha F_j^\beta = \frac{\Delta}{n} \bar{g}_{hi}, \quad (4.2)$$

Since the metric tensor  $\bar{g}_{ij}$  is regular ( $\det(\bar{g}) \neq 0$ ), in a neighbourhood of every point there exists a positive form  $A^{ij} z_i z_j$  such that

$$\bar{g}_{ij} A^{ij} \geq 0 \quad (\text{or } \bar{g}_{ij} A^{ij} \leq 0).$$

Contracting formula (4.2) we have  $\psi_{ij} A^{ij} \geq 0$  (or  $\psi_{ij} A^{ij} \leq 0$ ), which may be, in detail, written in the form:

$$(\Psi_{,ij} - \Psi_{,i} \varphi_j + \Psi_{,\alpha} \varphi_\beta F_i^\alpha F_j^\beta) A^{ij} \geq 0 \quad (\text{or } \leq 0) \quad (4.3)$$

where the symbol  $\geq$  (or  $\leq 0$ ) evidently depends on the signum of the constant  $\Delta$  and  $\psi = d\Psi$ .

The formula (4.3) may be written in the following form:

$$\frac{\partial^2 \Psi(x)}{\partial x^i \partial x^j} A^{ij}(x) + \frac{\partial \Psi(x)}{\partial x^i} B^i(x) \geq 0 \quad (\text{or } \leq 0)$$

where  $B^i(x)$  are function on coordinate neighbourhood of manifold.

It follows from the Hopf's Theorem (see [11], [24]) that this second order differential equations has only trivial solution  $\Psi = \text{const}$ . This implies  $\psi = 0$ .

Herewith the proof is complete.  $\square$

## References

- [1] Beklemishev, D. V.: *Differential geometry of spaces with almost complex structure*. Geometria. Itogi Nauki i Tekhn., *All-Union Inst. for Sci. and Techn. Information (VINITI)*, Akad. Nauk SSSR, Moscow, (1965), 165–212.
- [2] Boeckx, E., Kowalski, O., Vanhecke, L.: *Riemannian manifolds of conullity two*. *World Sci.*, 1996.
- [3] Domashev, V. V., Mikeš, J.: *Theory of holomorphically projective mappings of Kählerian spaces*. *Math. Notes* **23** (1978), 160–163, transl. from *Mat. Zametki* **23**, 2 (1978), 297–304.
- [4] Kurbatova, I. N.: *HP-mappings of H-spaces*. *Ukr. Geom. Sb., Kharkov* **27** (1984), 75–82.
- [5] Lakomá, L., Jukl, M.: *The decomposition of tensor spaces with almost complex structure*. *Suppl. Rend. Circ. Mat. (Palermo)* **72**, II (2004), 145–150.
- [6] Al Lamy, R. J. K., Škodová, M., Mikeš, J.: *On holomorphically projective mappings from equiaffine generally recurrent spaces onto Kählerian spaces*. *Arch. Math. (Brno)* **42**, 5 (2006), 291–299.
- [7] Mikeš, J.: *Geodesic mappings onto semisymmetric spaces*. *Russ. Math.* **38**, 2 (1994), 35–41, transl. from *Izv. Vyssh. Uchebn. Zaved., Mat.* **381**, 2 (1994), 37–43.
- [8] Mikeš, J.: *On special F-planar mappings of affine-connected spaces*. *Vestn. Mosk. Univ.* **3** (1994), 18–24.
- [9] Mikeš, J.: *Geodesic mappings of affine-connected and Riemannian spaces*. *J. Math. Sci., New York* **78**, 3 (1996), 311–333.

- [10] Mikeš, J.: *Holomorphically projective mappings and their generalizations*. J. Math. Sci., New York **89**, 3 (1998), 1334–1353.
- [11] Mikeš, J., Chodorová, M.: *On concircular and torse-forming vector fields on compact manifolds*. Acta Acad. Paedagog. Nyregyházi., Mat.-Inform. Közl. (2010).
- [12] Mikeš, J., Pokorná, O.: *On holomorphically projective mappings onto Kählerian spaces*. Suppl. Rend. Circ. Mat. (Palermo) **69**, II (2002), 181–186.
- [13] Mikeš, J., Radulović, Ž, Haddad, M.: *Geodesic and holomorphically projective mappings of  $m$ -pseudo- and  $m$ -quasisymmetric Riemannian spaces*. Russ. Math. **40**, 10 (1996), 28–32, transl. from Izv. Vyssh. Uchebn., Mat 1996, 10(413), 30–35.
- [14] Mikeš, J., Sinyukov, N. S.: *On quasiplanar mappings of spaces of affine connection*. Sov. Math. **27**, 1 (1983), 63–70, transl. from Izv. Vyssh. Uchebn. Zaved., Mat., 1983, 1(248), 55–61.
- [15] Mikeš, J., Starko, G. A.: *K-concircular vector fields and holomorphically projective mappings on Kählerian spaces*. Circ. Mat. di Palermo, Suppl. Rend. Circ. Mat. (Palermo) **46**, II (1997), 123–127.
- [16] Mikeš, J., Vanžurová, A., Hinterleitner, I.: *Geodesic Mappings and some Generalizations*. Palacký Univ. Publ., Olomouc, 2009.
- [17] Otsuki, T., Tashiro, Y.: *On curves in Kaehlerian spaces*. Math. J. Okayama Univ. **4** (1954), 57–78.
- [18] Petrov, A. Z.: *Simulation of physical fields*. In: Gravitation and the Theory of Relativity, **4–5**, Kazan' State Univ., Kazan, 1968, 7–21.
- [19] Sakaguchi, T.: *On the holomorphically projective correspondence between Kählerian spaces preserving complex structure*. Hokkaido Math. J. **3** (1974), 203–212.
- [20] Sinyukov, N. S.: *Geodesic mappings of Riemannian spaces*. Nauka, Moscow, 1979.
- [21] Sinyukov, N. S.: *Almost geodesic mappings of affinely connected and Riemannian spaces*. J. Sov. Math. **25** (1984), 1235–1249.
- [22] Sobchuk, V. S., Mikeš, J., Pokorná, O.: *On almost geodesic mappings  $\pi_2$  between semisymmetric Riemannian spaces*. Novi Sad J. Math. **29**, 3 (1999), 309–312.
- [23] Yano, K.: *Differential Geometry on Complex and Almost Complex Spaces*. Pergamon Press, Oxford–London–New York–Paris–Frankfurt, 1965.
- [24] Yano, K., Bochner, S.: *Curvature and Betti Numbers*. Annals of Mathematics Studies **32**, Princeton University Press, Princeton, 1953.