

Ludvík Janoš

Some cohomological aspects of the Banach fixed point principle

Mathematica Bohemica, Vol. 136 (2011), No. 3, 333–336

Persistent URL: <http://dml.cz/dmlcz/141653>

Terms of use:

© Institute of Mathematics AS CR, 2011

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SOME COHOMOLOGICAL ASPECTS OF THE
BANACH FIXED POINT PRINCIPLE

LUDVÍK JANOŠ, Claremont

(Received September 20, 2010)

Abstract. Let $T: X \rightarrow X$ be a continuous selfmap of a compact metrizable space X . We prove the equivalence of the following two statements: (1) The mapping T is a Banach contraction relative to some compatible metric on X . (2) There is a countable point separating family $\mathcal{F} \subset \mathcal{C}(X)$ of non-negative functions $f \in \mathcal{C}(X)$ such that for every $f \in \mathcal{F}$ there is $g \in \mathcal{C}(X)$ with $f = g - g \circ T$.

Keywords: Banach contraction, cohomology, cocycle, coboundary, separating family, core

MSC 2010: 54H25, 54H20

1. INTRODUCTION AND NOTATION

The object of our study is a continuous selfmap T of a compact metrizable space X . Let $\mathcal{C}(X)$ and $\mathcal{M}(X)$ denote the set of all continuous real-valued functions and the set of all compatible metrics on X , respectively. Regarding $\mathcal{C}(X)$ as an abelian group, we convert it to an $(\mathbb{N}, +)$ -module defining the action of $n \in \mathbb{N}$ on $f \in \mathcal{C}(X)$ by

$$n \cdot f = f \circ T^n,$$

where T^n is the n th iteration of T . In most textbooks on homological algebra only group-actions are treated, whereas $(\mathbb{N}, +)$ is only a commutative monoid reflecting the fact that T need not have an inverse. But it was shown by S. MacLane, H. Cartan, A. Bakakhanian [1] and others that the homology and cohomology theory of monoids is almost identical with that of groups. In our case only one-cohomology will be relevant. By a *one-chain* we understand a map

$$\varphi: \mathbb{N} \rightarrow \mathcal{C}(X)$$

with $\varphi(0) = 0 \in \mathcal{C}(X)$. The cochain φ is a cocycle if it is of the form

$$\varphi(n) = f + 1 \cdot f + 2 \cdot f + \dots + (n-1) \cdot f, \quad n \in \mathbb{N},$$

where f is a function in $\mathcal{C}(X)$. If f is of the form $g - 1 \cdot g$ with $g \in \mathcal{C}(X)$ then the cocycle is a *coboundary*. In this case we have

$$\sum_{k=0}^{n-1} k \cdot f = g - n \cdot g, \quad n \in \mathbb{N}$$

and by abuse of language we call the function itself a coboundary. If $\mathcal{F} \subset \mathcal{C}(X)$ is a family of functions we say \mathcal{F} is *point separating* if for any $x, y \in X$, $x \neq y$, there is $f \in \mathcal{F}$ with $f(x) \neq f(y)$. A selfmap $T: X \rightarrow X$ is a *Banach contraction* if for some $c \in [0, 1)$ and $d \in M(X)$ we have

$$d(Tx, Ty) \leq cd(x, y), \quad x, y \in X.$$

We will prove

Theorem 1. *The selfmap T is a Banach contraction if and only if there is a countable point separating family $\mathcal{F} \subset \mathcal{C}(X)$ of nonnegative coboundaries.*

2. PROOF OF THE THEOREM

We prove first the easier “only if” part. Thus let T be a c -contraction on a compact metric space (X, d) . Let $\{x_k\}_{k \in \mathbb{N}}$ be a countable dense subset of X and for every $k \in \mathbb{N}$ consider the orbit $\mathcal{O}(x_k) = \{T^n x_k : n \in \mathbb{N}\}$. We define the function f_k as the distance from $x \in X$ to the orbit $\mathcal{O}(x_k)$, i.e. $f_k(x) = d(x, \mathcal{O}(x_k))$. It is obvious that $f_k \in \mathcal{C}(X)$ and that $\sum_{n=0}^{\infty} f_k(T^n x)$ converges uniformly to some continuous function $g_k \in \mathcal{C}(X)$, so that $f_k(x) = g_k(x) - g_k(Tx)$, $x \in X$, showing that this f_k is a coboundary. It is also easy to see that the family $\mathcal{F} = \{f_k : k \in \mathbb{N}\}$ is point separating, which completes the “only if” part of the proof.

To prove the “if” part suppose $\mathcal{F} \subset \mathcal{C}(X)$ is a family with the above mentioned properties. Let $f \in \mathcal{F}$. Since f is a coboundary there is $g \in \mathcal{C}(X)$ with $f(x) = g(x) - g(Tx)$, $x \in X$. The equation

$$\sum_{k=0}^{n-1} f(T^k x) = g(x) - g(T^n x)$$

shows that the infinite sum $\sum_{k=0}^{\infty} f(T^k x)$ exists due to the fact that $f \geq 0$, g is continuous and the space X is compact. This implies that

$$(2.1) \quad \lim_{k \rightarrow \infty} f(T^k x) = 0, \quad x \in X.$$

We will show that the orbit

$$\mathcal{O}(x) = x, Tx, T^2x, \dots, T^n x, \dots$$

converges for every $x \in X$. Assume that for some x it is not true. Then there are two distinct points, say $y_1 \neq y_2$ towards which two subsequences of $\mathcal{O}(x)$ converge. But then 1 implies that $f(y_1) = f(y_2) = 0$. Since f is an arbitrary element of \mathcal{F} this implies that \mathcal{F} fails to separate y_1 from y_2 , a desired contradiction. This yields that the orbit $\mathcal{O}(x)$ converges and since $\lim[\mathcal{O}(x)] = T \lim[\mathcal{O}(x)]$ the limit is a fixed point, say x^* . From the equality

$$f(x^*) = g(x^*) - g(Tx^*) = 0$$

it follows that T cannot have more fixed points than one since otherwise \mathcal{F} would not separate them. Thus T has a unique fixed point x^* toward which every orbit converges. We will show also that T has no periodic points. Suppose a point $x \in X$ has period $p \geq 2$, i.e. $T^p x = x$. Since

$$f(x) + f(Tx) + \dots + f(T^{p-1}x) = g(x) - g(T^p x) = 0$$

we obtain that

$$f(x) + f(Tx) + \dots + f(T^{p-1}x) = 0$$

implying that $f(T^k x) = 0$ for $k = 0, 1, \dots, p-1$ which would again clash with the separation property of \mathcal{F} . This also shows how strong the argument of separation is.

In the final stage of our proof the function g appearing in $f(x) = g(x) - g(Tx)$ will play an important role. Note that it is not uniquely determined by f , since $g + c$, with c any constant, can replace g . Thus we can choose the function g which corresponds to $f \in \mathcal{F}$ by setting $g(x^*) = 0$. Since

$$\sum_{k=0}^{n-1} f(T^k x) = g(x) - g(T^n x)$$

and the orbit converges to x^* it follows that the infinite sum $\sum_{k=0}^{\infty} f(T^k x)$ converges to the function g .

One of our previous results (see [1]) is that T is a Banach contraction if and only if the core of T , $\text{core}(T) = \bigcap \{T^n(X) : n \in \mathbb{N}\}$ is a singleton. It is known that the core is a nonempty compact T -invariant subset of X and that the restriction of T to the core is surjective. Thus, to conclude our proof we will assume that $|\text{core}(T)| \geq 2$ and deduce from it a contradiction. Since the fixed point x^* is in the core there is in the core another point, say y , distinct from x^* . Since T maps the core onto itself every point in it has at least one pre-image in the core. Since there are no periodic points we can construct an inverse orbit from y , i.e. a sequence $\{y_k\}_{k=0}^{\infty}$ of distinct points with $Ty_{k+1} = y_k$, $k \in \mathbb{N}$. We observe that there must be at least one $f \in \mathcal{F}$ with $f(y) > 0$ since otherwise \mathcal{F} would not separate y from x^* . Let g be the function corresponding to f and consider the sequence $\{g(y_k)\}_{k=0}^{\infty}$. Since $g(y_{k+1}) \geq g(y_k)$, $k \in \mathbb{N}$ and $g(x) \geq f(x)$ for every $x \in X$ the limit of this sequence exists and is positive. Let $m > 0$ be the limit. Let Z denote the set of all points in the core obtainable as limits of subsequences from $\{y_k\}_{k=0}^{\infty}$. It follows that for any $z \in Z$ we have $g(z) = m > 0$. We consider the orbit $\mathcal{O}(z)$. Let $\{y_{k(i)}\}$ where $k(1) < k(2) < \dots \rightarrow \infty$ be the subsequence of $\{y_k\}$ the limit of which is z . Then Tz is obtained as the limit of $\{y_{k(i)-1}\}$. From this we see that $\mathcal{O}(z) \subset Z$ and the sequence $\{g(T^n z)\}_{n=0}^{\infty}$ is the sequence of constants $m > 0$. But since $\{T^n z\}$ converges to x^* we obtain the desired contradiction since $g(x^*) = 0$. Thus the core of T is the singleton $\{x^*\}$ which concludes the proof.

References

- [1] *A. Bakakhanian*: Cohomological Methods in Group Theory. Marcel Dekker, New York, 1972.
- [2] *L. Janoš*: The Banach contraction mapping principle and cohomology. Comment. Math. Univ. Carolin. 41 (2000), 605–610.

Author's address: Ludvík Janoš, PO Box 1563, Claremont, CA 91711, USA.