

Václav Flaška; Jaroslav Ježek; Tomáš Kepka
Transitive closures of binary relations. III.

Acta Universitatis Carolinae. Mathematica et Physica, Vol. 49 (2008), No. 1, 25--31

Persistent URL: <http://dml.cz/dmlcz/142771>

Terms of use:

© Univerzita Karlova v Praze, 2008

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Transitive Closures of Binary Relations III

VÁCLAV FLAŠKA, JAROSLAV JEŽEK and TOMÁŠ KEPKA

Praha

Received 2nd October 2007

Transitive closures of the covering relation in lattices are investigated.
Vyšetřují se tranzitivní uzávěry pokrývací relace ve svazech.

This extremely short expository note collects a few more or less notoriously known results on the covering relation β in lattices. Special attention is paid to the property that any two β -sequences connecting two given elements are of the same length. We refer to [1] and [2] as for terminology, notation, further references, etc.

1. The covering relation in lattices

Throughout the note, let $L = L(+, \cdot)$ be a lattice (i.e., both $L(+)$ and $L(\cdot)$ are semilattices and $a(a + b) = a = a + (ab)$ for all $a, b \in L$). Define a relation α on L by $(a, b) \in \alpha$ if and only if $a + b = b$.

1.1 Proposition.

- (i) *The relation α is a stable (reflexive) ordering of the lattice and $(a, b) \in \alpha$ if and only if $ab = a$.*
- (ii) *$(a, a + b) \in \alpha$, $(b, a + b) \in \alpha$, $(ab, a) \in \alpha$ and $(ab, b) \in \alpha$ for all $a, b \in L$. (In fact, $a + b = \sup_{\alpha}(a, b)$ and $ab = \inf_{\alpha}(a, b)$.)*

Department of Algebra, MFF UK, Sokolovská 83, 186 75 Praha 8, Czech Republic

The work is a part of the research project MSM 0021620839 financed by MŠMT and partly supported by the Grant Agency of Czech Republic, grant GAČR-201/05/0002.

E-mail adress: flaska@karlin.mff.cuni.cz

E-mail adress: jezek@karlin.mff.cuni.cz

E-mail adress: kepka@karlin.mff.cuni.cz

- (iii) An element $a \in L$ is maximal in $L(\alpha)$ (i.e., a is right α -isolated) if and only if $a = 1_L$ is an absorbing element of $L(+)$ if and only if a is a neutral element of $L(\cdot)$. (Then a is the (unique) greatest element of $L(\alpha)$.)
- (iv) An element $a \in L$ is minimal in $L(\alpha)$ (i.e., a is left α -isolated) if and only if $a = 0_L$ is a neutral element of $L(+)$ if and only if a is an absorbing element of $L(\cdot)$. (Then a is the (unique) smallest element of $L(\alpha)$.)

Proof. It is obvious. □

1.2. Lemma.

- (i) Every weakly pseudoirreducible finite α -sequence is pseudoirreducible.
- (ii) Every weakly pseudoirreducible right (left, resp.) directed infinite α -sequence is pseudoirreducible.
- (iii) If there exists no pseudoirreducible right (left, resp.) directed infinite α -sequence then $1_L \in L$ ($0_L \in L$, resp.).

Proof. It is obvious. □

1.3 Lemma. Let $(a, b) \in \alpha$ and $I = \text{Int}_x(a, b) = \{c \in L \mid (a, c) \in \alpha \text{ and } (c, b) \in \alpha\}$.

Then:

- (i) I is a sublattice of L and $\{a, b\} \subseteq I$.
- (ii) $a = 0_I$ and $b = 1_I$.
- (iii) $\alpha_I = \alpha_L \upharpoonright I$.

Proof. It is obvious. □

In the sequel, put $\beta = \sqrt{\alpha}$ and $\gamma = \mathbf{rt}(\beta)$, so that β is the covering relation of L and γ is its reflexive and transitive closure. Notice that $\mathbf{i}(\gamma) = \mathbf{t}(\beta)$.

1.4 Proposition.

- (i) β is totally antitransitive.
- (ii) $\beta \subseteq \gamma \subseteq \alpha$.
- (iii) γ is an ordering of L .
- (iv) If $(a, b) \in \alpha$ and $\text{Int}_x(a, b)$ is finite then $(a, b) \in \gamma$.

Proof. It is obvious. □

We say that the lattice L is resuscitable if so is the ordering α (i.e., $\alpha = \gamma$).

1.5 Proposition. The lattice L is resuscitable, provided that the following two conditions are satisfied:

- (1) no right (left, resp.) directed infinite $\mathbf{i}(\alpha)$ -sequence is right (left, resp.) bounded in $L(\alpha)$;
- (2) no left (right, resp.) directed infinite β -sequence is left (right, resp.) bounded in $L(\alpha)$.

Proof. See II.1.8. □

1.6 Corollary. The lattice L is resuscitable, provided that it is finite.

1.7 Example. The boolean lattice of all subsets of an infinite set is not resuscitable.

1.8 Example. A chain is resuscitable if and only if it can be embedded into the chain of integers (with respect to the usual ordering of integers).

1.9 Example. Consider the lattice $L_1 = \{1, a_0, a_1, a_2, \dots, b_0, b_1, b_2, \dots\}$ with $(x, y) \in \alpha$ if and only if either $x = y$, or $x = a_0$, or $y = 1$, or $(x, y) = (a_i, a_j)$ where $i \leq j$, or $(x, y) = (a_i, b_j)$ where $i \leq j$. This infinite lattice L_1 is resuscitable, while its sublattice $\{1, a_0, a_1, a_2, \dots\}$ is not. It follows that the class of resuscitable lattices is not closed under sublattices.

2. On when the covering relation is right/left confluent (or weakly semimodular lattices)

The lattice L is called

- upwards (downwards, resp.) weakly semimodular if the semilattice $L(+)$ ($L(\cdot)$, resp.) is weakly semimodular;
- weakly semimodular if it is both upwards and downwards weakly semimodular.

2.1 Lemma. *The lattice L is upwards (downwards, resp.) weakly semimodular if and only if the relation β is right (left, resp.) confluent.*

Proof. See II.2.1. □

2.2 Lemma. *Assume that L is upwards (downwards, resp.) weakly semimodular. If $a, b, c \in L$ are such that $(a, b) \in \gamma$ and $(a, c) \in \gamma$ ($(b, a) \in \gamma$ and $(c, a) \in \gamma$, resp.) then $(b, b + c) \in \gamma$ and $(c, b + c) \in \gamma$ ($(bc, b) \in \gamma$ and $(bc, c) \in \gamma$, resp.).*

Proof. See II.2.3. □

2.3 Corollary. *If L is upwards (downwards, resp.) weakly semimodular then the ordering γ is right (left, resp.) strictly confluent.*

2.4 Lemma. *Assume that L is upwards (downwards, resp.) weakly semimodular. If $(a, b) \in \gamma$ then there exists no right (left, resp.) directed infinite $\mathbf{i}(\gamma)$ -sequence (a_0, a_1, a_2, \dots) ((\dots, b_2, b_1, b_0) , resp.) such that $a_0 = a$ ($b_0 = b$, resp.) and $(a_i, b) \in \alpha$ ($(a, b_i) \in \alpha$, resp.) for every $i \geq 1$.*

Proof. See II.2.6. □

2.5 Lemma. *Assume that L is weakly semimodular. If $(a, b) \in \gamma$ then:*

- (i) $K = \text{Int}_\gamma(a, b)$ is a sublattice of L , $a = 0_K$ and $b = 1_K$.
- (ii) K is resuscitable.
- (iii) *If $c \in \text{Int}_x(a, b)$ and either $(a, c) \in \gamma$ or $(c, b) \in \gamma$ then $c \in K$.*

Proof. See II.2.7. □

2.6 Example. Consider the lattice L_2 with seven elements $0, 1, a, b, c, d, e$ and the covering relation $\beta = \{(0, a), (0, b), (a, c), (a, d), (b, d), (b, e), (c, 1), (d, 1), (e, 1)\}$. (A finite lattice is uniquely determined by its covering relation.) Clearly, L_2 is upwards weakly semimodular but not downwards weakly semimodular.

2.7 Example. Consider the lattice \mathbf{N} with five elements $0, 1, a, b, c$ and the covering relation $\beta = \{(0, a), (0, b), (a, 1), (b, c), (c, 1)\}$. Clearly, \mathbf{N} is neither upwards nor downwards weakly semimodular.

3. Semimodular lattices

The lattice L is called

- upwards (downwards, resp.) semimodular if the semilattice $L(+)$ ($L(\cdot)$, resp.) is semimodular;
- semimodular if it is both upwards and downwards semimodular.

3.1 Lemma.

- (i) If L is (upwards, downwards) semimodular then it is (upwards, downwards) weakly semimodular.
- (ii) If L is semimodular then γ is a stable ordering of L .

Proof. See II.3.2. □

3.2 Proposition. Assume that L is resuscitable. Then L is (upwards, downwards) semimodular if and only if it is (upwards, downwards) weakly semimodular.

Proof. See II.3.3. □

3.3 Corollary. If L is finite then L is (upwards, downwards) semimodular if and only if it is (upwards, downwards) weakly semimodular.

3.4 Proposition. Assume that L is weakly semimodular. Let $(a, b) \in \gamma$ and $K = \text{Int}_\gamma(a, b)$. Then:

- (i) K is a sublattice of L , $a = 0_K$ and $b = 1_K$.
- (ii) K is semimodular and resuscitable.
- (iii) Every subchain of K (α) is finite and of length at most $\text{dist}_\gamma(a, b)$.
- (iv) $K \subseteq \text{Int}_\alpha(a, b)$ and $c \in K$, provided that $c \in \text{Int}_\alpha(a, b)$ and either $(a, c) \in \gamma$ or $(c, b) \in \gamma$.
- (v) If L is upwards or downwards semimodular then $K = \text{Int}_\alpha(a, b)$.

Proof. Combine 2.5 and II.6.3. □

3.5 Proposition. The following four conditions are equivalent:

- (i) L is upwards (downwards, resp.) weakly semimodular, no right directed infinite $\mathbf{i}(\alpha)$ -sequence is right bounded in $L(\alpha)$ and no left directed infinite β -sequence is left bounded in $L(\alpha)$.

- (ii) L is upwards (downwards, resp.) weakly semimodular, no left difrected infinite $\mathbf{i}(\alpha)$ -sequence is left bounded in $L(\alpha)$ and no right directed infinite β -sequence is right bounded in $L(\alpha)$.
- (iii) L is upwards (downwards, resp.) semimodular and resuscitable.
- (iv) L is upwards (downwards, resp.) weakly semimodular and every right and left bounded subchain of $L(\alpha)$ is finite.

Proof. See II.6.4. □

3.6 Example. The lattice L_2 from 2.6 is upwards semimodular but not downwards weakly semimodular.

3.7 Example. Consider the lattice $L_3 = \{0, 1, a, b_1, b_2, \dots\}$ with $(x, y) \in \alpha$ if and only if either $x = y$ or $x = 0$ or $y = 1$ or $(x, y) = (b_i, b_j)$ where $i < j$. This infinite lattice L_3 is weakly semimodular but neither upwards nor downwards semimodular. Moreover, $(0, 1) \in \gamma$, $\text{dist}_\gamma(0, 1) = 2$ and $\text{Int}_\gamma(0, 1) = \{0, \alpha, 1\} \neq L_3 = \text{Int}_\alpha(0, 1)$.

4. Modular lattices

The lattice L is called modular if no sublattice of L is a copy of the pentagon (the lattice \mathbf{N} from 2.7).

4.1 Proposition. *If L is modular then it is semimodular.*

Proof. It is obvious. □

4.2 Proposition. *A resuscitable lattice is modular if and only if it is weakly semimodular.*

Proof. The direct implication follows from 4.1. Let L be a resuscitable, weakly semimodular lattice. By 3.2, L is semimodular. Let $x < y$ stand for $(x, y) \in \mathbf{i}(\alpha)$ and $x \prec y$ stand for $(x, y) \in \beta$. Suppose that L is not modular, so that it contains a subpentagon $\{o, a, b, c, i\}$ (o is its smallest element, i is the largest, and $b < c$). Choose these five elements in such a way that the interval $\text{Int}(o, i)$ has minimal possible length. (Since L is resuscitable and semimodular, every interval I of L has a finite length n and every maximal chain in I is of length n).

Suppose $o < b$. Then $a < i$ by the upwards semimodularity, from which we get $o < c$ by the downwards semimodularity, a contradiction. Thus o is not covered by b and there exists an element $d \in L$ with $o \prec d < b$. Put $e = a + d$. By the upwards semimodularity we have $a < e$; since a is not covered by i , we get $a \prec e < i$. Thus $b \not\leq e$ and e is incomparable with both b and c . By the minimality of $\text{Int}(o, i)$, the elements d, e, b, c, i do not form a subpentagon. Since $e + b = i$, we get $ec \not\leq d$. Put $f = ec$. Thus $d < f < e$. But then the elements o, a, d, f, e form a subpentagon of L , a contradiction with the minimality of $\text{Int}(o, i)$. □

4.3 Corollary. *A finite lattice is modular if and only if it is semimodular.*

4.4 Example. Proposition 4.2 cannot be generalized to arbitrary lattices. Let L be any infinite lattice such that its covering relation is empty. Then L is semimodular. Of course, such a lattice need not to be modular. Thus a semimodular lattice is not necessarily modular.

The lattice L is called

- upwards (downwards, resp.) strongly modular if the semilattice $L(+)$ ($L(\cdot)$, resp.) is strongly modular;
- strongly modular if it is both upwards and downwards strongly modular.

4.5 Example. For every cardinal number $\kappa > 0$ denote by M_κ the (unique up to isomorphism) lattice of length 2 with κ atoms (elements covering the least element). Clearly, each M_κ is a strongly modular lattice. We see that a strongly modular lattice is not necessarily distributive.

4.6 Example. Denote by L_4 the lattice with six elements a, b, c, d, e, f , such that $\beta = \{(a,b), (b,c), (c,f), (a,d), (d,e), (b,e), (e,f)\}$. (The product of the two-element chain with the three-element chain.) Clearly, L_4 is neither downwards nor upwards strongly modular. On the other hand, it is distributive.

4.7 Proposition. *The following conditions are equivalent:*

- (i) L is upwards strongly modular;
- (ii) L is downwards strongly modular;
- (iii) L is strongly modular;
- (iv) neither N nor L_4 can be embedded into L .

Proof. By 4.6, each of the first three conditions implies (iv). Thus it is sufficient to prove that (iv) implies (i). Let L be a modular lattice not containing a sublattice isomorphic with L_4 and suppose that L is not upwards strongly modular, so that it contains four distinct elements a, b, c, i such that a is incomparable with b , $i = a + b$ and $b < c < i$. If $ac < i$ then these four elements together with ac form a subpentagon, a contradiction. Thus ac is incomparable with b . Put $d = ac$ and $e = ab = db$; we have $e < d < a < i$. Also, put $f = d + b$, so that $b < f \leq c < i$. It can be easily checked that the elements e, d, a, b, f, i . It can be easily checked that the elements e, d, a, b, f, i form a sublattice isomorphic with L_4 , a contradiction. \square

4.8 Example. For two finite lattices P and Q we define a lattice $L = P \oplus Q$, called their glued ordinal sum, as follows. We can assume that $P \cap Q = \{1_P\} = \{0_Q\}$. In that case put $L = P \cup Q$ and $\alpha_L = \alpha_P \cup \alpha_Q \cup (P \times Q)$. Similarly, we can define $R_1 \oplus \dots \oplus R_n$ for any finite nonempty sequence of lattices R_1, \dots, R_n . It follows from 4.7 that a finite lattice is strongly modular if and only if it can be expressed as the glued ordinal sum of a finite sequence

of finite lattices, each of which is either a chain or isomorphic to M_n for some $n \geq 2$.

5. On when the covering relation is regular

5.1 Proposition. *If L is upwards or downwards weakly semimodular then its covering relation β is regular.*

Proof. See II.5.1. □

References

- [1] FLÁŠKA, V., JEŽEK, J., KEPKA, T. AND KORTELAINEN, J.: *Transitive closures of binary relations I.* Acta Univ. Carolinae 48 (2007), 55–69.
- [2] FLÁŠKA, V., JEŽEK, J. AND KEPKA, T.: *Transitive closures of binary relations II.* Acta Univ. Carolinae 48 (2007), 71–80.