

# Applications of Mathematics

---

Pigong Han

Interior regularity of weak solutions to the perturbed Navier-Stokes equations

*Applications of Mathematics*, Vol. 57 (2012), No. 5, 427--444

Persistent URL: <http://dml.cz/dmlcz/142908>

## Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

INTERIOR REGULARITY OF WEAK SOLUTIONS TO THE  
PERTURBED NAVIER-STOKES EQUATIONS\*

PIGONG HAN, Beijing

(Received July 7, 2009)

*Abstract.* In this paper we establish interior regularity for weak solutions and partial regularity for suitable weak solutions of the perturbed Navier-Stokes system, which can be regarded as generalizations of the results in *L. Caffarelli, R. Kohn, L. Nirenberg*: Partial regularity of suitable weak solutions of the Navier-Stokes equations, *Commun. Pure. Appl. Math.* 35 (1982), 771–831, and *S. Takahashi*, On interior regularity criteria for weak solutions of the Navier-Stokes equations, *Manuscr. Math.* 69 (1990), 237–254.

*Keywords:* perturbed Navier-Stokes equations, interior regularity, partial regularity

*MSC 2010:* 35Q30, 76D05, 35B65

1. INTRODUCTION AND MAIN RESULTS

Let  $\Omega \subset \mathbb{R}^N$  ( $N = 2, 3$ ) be an open domain with smooth boundary  $\partial\Omega$ . The dynamical behavior of the viscous incompressible fluid flow in  $\Omega$  is governed by the following perturbed Navier-Stokes system (see page 140 in [1]):

$$\begin{cases} \partial_t v - \Delta v + (v \cdot \nabla)v + \nabla \pi_1 = -\partial_t v_0 - (\partial_t \omega_0) \times x - \omega_0 \times (\omega_0 \times x) - 2\omega_0 \times v, \\ \nabla \cdot v = 0, \end{cases}$$

where the translational velocity  $v_0 = v_0(t)$  and the angular velocity  $\omega_0 = \omega_0(t) = (0, 0, \dots, 0, \theta(t))$  are  $N$ -vectors depending only on the time-variable  $t$ ;  $v = (v^1(x, t), v^2(x, t), \dots, v^N(x, t))$  and  $p = p(x, t)$  denote the unknown velocity vector and the pressure, respectively.

---

\* This work was completed with the support in part by Key Laboratory of Random Complex Structures and Data Science, Chinese Academy of Sciences; National Natural Science Foundation of China under grant No. 11071239.

Set  $u = v + v_0 + \omega_0 \times x$ ,  $\pi = \pi_1 - v_0 \cdot (\omega_0 \times x)$ . Then the above equations can be rewritten as follows:

$$(1.1) \quad \partial_t u - \Delta u + (u \cdot \nabla)u + \nabla \pi = v_0 \cdot \nabla u + (\omega_0 \times x) \cdot \nabla u - \omega_0 \times u, \\ (x, t) \in \Omega \times (0, T),$$

$$(1.2) \quad \nabla \cdot u = 0, \quad (x, t) \in \Omega \times (0, T).$$

**Definition.** The function  $u$  is called a *weak solution* of (1.1), (1.2) if  $u \in L^\infty(0, T; L^2(\Omega))$  with  $\nabla u \in L^2(0, T; L^2(\Omega))$  satisfies

$$- \int_0^T \int_\Omega u \partial_\tau v \, dx \, d\tau + \int_0^T \int_\Omega \nabla u \cdot \nabla v \, dx \, d\tau + \int_0^T \int_\Omega (u \cdot \nabla)u \cdot v \, dx \, d\tau \\ = \int_0^T \int_\Omega [v_0 \cdot \nabla u + (\omega_0 \times x) \cdot \nabla u - \omega_0 \times u]v \, dx \, d\tau \\ \text{for all } v \in C_0^\infty((0, T); C_{0,\sigma}^\infty(\Omega)).$$

Furthermore, we say that the weak solution  $u$  is a *suitable weak solution* of (1.1), (1.2) if the associated pressure  $\pi$  is in  $L^{N/2}(\Omega \times (0, T))$ , and the following generalized energy inequality holds:

$$\int_\Omega |u(x, t)|^2 \phi(x, t) \, dx + 2 \int_0^t \int_\Omega |\nabla u|^2 \phi \, dx \, d\tau \\ \leq \int_0^t \int_\Omega [|u|^2(\phi_\tau + \Delta \phi) + (|u|^2 + 2\pi)u \cdot \nabla \phi] \, dx \, d\tau \\ - 2 \int_0^t \int_\Omega |u|^2 (v_0(\tau) + \omega_0(\tau) \times x) \cdot \nabla \phi \, dx \, d\tau \\ \text{for each } 0 \leq \phi \in C_0^\infty(\Omega \times (0, T)).$$

In the absence of the terms on the right-hand side of (1.1), problem (1.1), (1.2) is reduced to the standard incompressible Navier-Stokes equation. In the three dimensional case, a large gap remains between the regularity available in the existence results and the additional regularity required in the sufficient conditions to guarantee the smoothness of weak solutions of the standard Navier-Stokes equations. This gap has been narrowed by the works of Iskauriaza-Seregin-Sverak [18], Ladyzhenskaya-Seregin [19], Scheffer [25], Serrin [27], Struwe [29], see also [2], [3], [4], [5], [6], [8], [9], [10], [13], [14], [15], [16], [20], [22], [23], [24], [26], [31], [32] and the references therein, which bring about a deeper understanding of the regularity. In particular, some local partial regularity results and Hausdorff dimension estimates on the possible singular set have been obtained for a class of suitable weak solutions defined and constructed

in [7], where the principal tools are the so-called generalized energy inequality and a scaling argument.

The exterior problem of (1.1), (1.2) was studied in [11] from the point of view of mechanics. Hishida [17] considered the problem (1.1), (1.2) with  $v_0(t) \equiv 0$ ,  $\omega_0(t) \equiv \text{const}$ ,  $0 < |\omega_0(t)| \ll 1$ , and established an existence result in a three-dimensional exterior domain. Chen-Miyakawa [12] discussed the existence of a global weak solution in the whole space  $\mathbb{R}^N$  ( $N = 2, 3$ ) and the algebraic decay rates for the kinetic energy of the weak solutions constructed. The global existence of weak solutions of (1.1), (1.2) with inhomogeneous boundary values in smooth bounded domains can be constructed by using Galerkin methods and weak convergence theories, see [21] for example. In general domains, it is difficult to construct a global weak solution for (1.1), (1.2), because the perturbed terms contain the space variable  $x$ , which causes many difficulties at infinity (see Hishida's paper [17] for example). However, in this paper we mainly focus on the regularity of weak solutions, not the existence.

As far as we know, there are few results on the regularity of weak solutions of (1.1), (1.2). In the present paper, we try to establish regularity criteria for (1.1), (1.2), which are similar to Serrin's class in [27] and Takahashi's criteria in [30]. In addition, we also discuss partial regularity for suitable weak solutions of (1.1), (1.2), which can be regarded as an extension of results in [7]. The regularity of the solution at the point  $(x_0, t_0)$  seems to follow almost directly from the corresponding results for the non-perturbed Navier-Stokes equations, but we cannot find exact references explaining these questions. This is why we give strict and detailed proofs in the present paper.

In this paper, we always suppose that  $v_0(t)$ ,  $\omega_0(t)$  are smooth bounded on  $[0, T)$ ,  $0 < T < \infty$ . Our main results read as follows:

**Theorem 1.1.** *Suppose that  $u$  is a weak solution of (1.1), (1.2) with  $N = 2, 3$ . Let one of the following conditions hold:*

$$\|u\|_{L^q(0,T;L^p(\Omega))} < \infty \quad \text{for any } N < p \leq \infty \text{ with } \frac{N}{p} + \frac{2}{q} \leq 1,$$

or

$$\|u\|_{L^\infty(0,T;L^N(\Omega))} \quad \text{is suitable small.}$$

Then, for any  $\Omega'$  with  $\overline{\Omega'} \subset \Omega$  and any  $0 < \sigma < T$ ,

$$u, \text{curl } u \in L^\infty(\Omega' \times (\sigma, T)).$$

Now we recall the Lorentz space  $L^{(q)}$  for  $1 < q < \infty$ ,

$$L^{(q)}(0, T) = \left\{ f \in L^1(0, T) : \sup_{\lambda > 0} \lambda \{ \mu \{ t \in (0, T) : |f(t)| > \lambda \} \}^{1/q} \right\},$$

where  $\mu$  denotes the Lebesgue measure on  $\mathbb{R}$ .

Define  $g \in L^{(q),p}(Q)$ ,  $Q = \Omega \times (0, T)$  if

$$\|g\|_{L^{(q),p}(Q)} = \{ \|g(\cdot, t)\|_{L^p(\Omega)} \}_{L^{(q)}(0, T)} < \infty.$$

Set  $Q_R(x, t) = \{(y, \tau) \in \Omega \times (0, T) : |y - x| < R, t - R^2 < \tau < t\}$ .

**Theorem 1.2.** *Let  $1 \leq p, q \leq \infty$  with  $N/p + 2/q \leq 1$  and  $p > N$  ( $N = 2, 3$ ). Assume that  $u$  is a weak solution of (1.1), (1.2) in  $Q_R = Q_R(x_0, t_0) \subset \Omega \times (0, T)$ . If there exists a positive number  $\varepsilon = \varepsilon(N, p) \ll 1$  such that*

$$\|u\|_{L^{(q),p}(Q_R)} \leq \varepsilon,$$

then

$$u, \operatorname{curl} u \in L^\infty(Q_{R/2}).$$

Next we deal with the partial regularity of suitable weak solutions of (1.1), (1.2). As in [7], we call a point  $(x, t)$  singular if  $u$  is not  $L^\infty_{\text{loc}}$  in any neighborhood of  $(x, t)$ ; the remaining points, where  $u$  is locally essentially bounded, will be called regular points.

For any  $X \subset \mathbb{R}^N \times \mathbb{R}$  and  $k \geq 0$ , define (see [7])

$$\mathcal{F}^k(X) = \lim_{\delta \rightarrow 0} \mathcal{F}_\delta^k(X), \quad \text{where } \mathcal{F}_\delta^k(X) = \inf \left\{ \sum_{i=1}^{\infty} r_i^k : X \subset \bigcup_{i=1}^{\infty} Q_{r_i}, r_i < \delta \right\}.$$

Then  $\mathcal{F}^k$  is a Borel regular measure. Hausdorff measure  $\mathcal{H}^k$  is defined in an entirely similar manner, but with  $Q_{r_i}$  replaced by an arbitrary closed subset in  $\mathbb{R}^N \times \mathbb{R}$  with diameter at most  $r_i$ . Clearly,  $\mathcal{H}^k \leq C\mathcal{F}^k$ .

It could be easily verified that any weak solution  $u$  is in  $L^q(0, T; L^p(\Omega))$  with  $N/p + 2/q = 1$ ,  $p > N = 2$ . Therefore, based on Theorem 1.1, we always take the dimension  $N = 3$  in the next arguments.

**Theorem 1.3.** *Let  $\varepsilon_1 > 0$  be an absolute constant. Suppose that  $(u, \pi)$  is a suitable weak solution of (1.1), (1.2) on  $Q_R(x_0, t_0) \subset \Omega \times (0, T)$  such that*

$$R^{-2} \iint_{Q_R(x_0, t_0)} (|u|^3 + |u||\pi|) \, dx \, dt + R^{-13/4} \int_{t_0 - R^2}^{t_0} \left( \int_{|x - x_0| < R} |\pi| \, dx \right)^{5/4} \, dt \leq \varepsilon_1.$$

*Then  $|u(x, t)| \leq c(R)$  for almost all  $(x, t) \in Q_{R/2}(x_0, t_0)$ . In particular,  $\mathcal{F}^{5/3}(S) = 0$ , where  $S$  denotes the set of singular points.*

Compared to Proposition 1 in [7], Theorem 1.3 shows that the perturbed terms in problem (1.1) have little effect on the regularity of the standard Navier-Stokes flows.

**Theorem 1.4.** *Let  $\varepsilon_2 > 0$  be an absolute constant. Suppose that  $(u, \pi)$  is a suitable weak solution of (1.1), (1.2) on  $Q_{R_1}(x_0, t_0) \subset \Omega \times (0, T)$  such that*

$$\limsup_{r \rightarrow 0} \iint_{Q_r(x_0, t_0)} |\nabla u|^2 \, dx \, dt \leq \varepsilon_2.$$

*Then  $(x_0, t_0)$  is a regular point of  $u$ . Furthermore,  $\mathcal{F}^1(S) = 0$ , which implies that  $S$  has Hausdorff dimension at most 1.*

Compared to Proposition 2 in [7], Theorem 1.4 reveals that the perturbed Navier-Stokes system has the same Hausdorff dimensional estimates at the singular points of suitable weak solutions of (1.1).

The local theory of partial regularity which we develop to prove Theorem 1.4 has other applications as well. We use it to study the initial value problem on  $\mathbb{R}^3$  in case the initial velocity  $u_0$  satisfies either

$$(1.3) \quad \frac{1}{2} \int_{\mathbb{R}^3} |x| |u_0(x)|^2 \, dx \triangleq G < \infty$$

or

$$(1.4) \quad \frac{1}{2} \int_{\mathbb{R}^3} |u_0(x)|^2 |x|^{-1} \, dx \triangleq L < L_0.$$

Condition (1.3) stipulates that  $u_0$  should “decay sufficiently rapidly at  $\infty$ ”, and under such assumption we can show that  $u$  is regular for  $|x|$  large enough.

**Theorem 1.5.** *Suppose that  $(u, \pi)$  is a suitable weak solution of (1.1), (1.2) on  $\mathbb{R}^3 \times (0, T)$  with initial data  $u_0 \in L^2_\sigma(\mathbb{R}^3)$ , and suppose (1.3) holds. Then  $u$  is regular in the region  $\{(x, t) \in \mathbb{R}^3 \times (0, T) : |x|^2 t > K_1\}$ , where  $K_1 = K_1(E, G)$  is a constant depending only on  $G$  and  $E \triangleq \frac{1}{2} \int_{\mathbb{R}^3} |u_0(x)|^2 dx$ .*

In the limit  $t \rightarrow 0$ , the region  $\{(x, t) \in \mathbb{R}^3 \times (0, T) : |x|^2 t > K_1\}$  in Theorem 1.5 expands to include all space. However, if  $u_0$  is regular enough (for example,  $u_0 \in H^1(\mathbb{R}^3)$ ),  $u$  is regular for a short time interval; in this case, Theorem 1.5 shows that the set of possible singular points is bounded.

**Theorem 1.6.** *Suppose that  $(u, \pi)$  is a suitable weak solution of (1.1), (1.2) on  $\mathbb{R}^3 \times (0, T)$  with initial data  $u_0 \in L^2_\sigma(\mathbb{R}^3)$ . Let there exist an absolute constant  $L_0 > 0$  such that (1.4) holds. Then  $u$  is regular in the region*

$$Y \triangleq \left\{ (a, s) \in \mathbb{R}^3 \times (0, T) : |a|^2 < \frac{s(L_0 - L)}{2M^2} - \left(4 + \frac{M_0}{2M}\right) s^2 \right\}.$$

Here  $M_0 \triangleq \|u\|_{L^\infty(0, T; L^2(\mathbb{R}^3))}^2$ ,  $M \triangleq 2 \sup_{0 \leq t < T} (|v_0(t)| + (1+t)|\omega_0(t)|) < \infty$ .

**Remark.** If  $|a| > 0$ ,  $s > 0$  in Theorem 1.6 are small enough, we know that the set  $Y \neq \emptyset$ . Condition (1.4) represents a restriction that  $u_0$  “not be too singular” at the origin. As it turns out, the argument works only if  $L$  is sufficiently small. Theorem 1.6 shows, for an appropriate choice of the constant  $L_0$ , that a suitable weak solution with initial data satisfying (1.4) is regular in a parabolic domain above the origin.

Let  $\Omega = \mathbb{R}^3$  or an exterior domain in  $\mathbb{R}^3$ . If the assumptions (1.3) and (1.4) are removed, then any suitable weak solution of (1.1), (1.2) is still regular if the space variable  $x$  is large enough and the time variable  $t$  is bounded below by a positive small number.

**Theorem 1.7.** *Let  $\Omega = \mathbb{R}^3$  or an exterior domain in  $\mathbb{R}^3$ , and let  $\delta \in (0, T)$ . Suppose that  $(u, \pi)$  is a suitable weak solution of (1.1), (1.2) on  $\Omega \times (0, T)$ . Then there exist constants  $K = K(\delta, T, u, \pi) > 0$ ,  $C = C(\delta, T, u, \pi) > 0$  such that  $|u(x, t)| \leq C$  for almost all  $(x, t) \in \Omega \times [\delta, T)$  with  $|x| \geq K$ .*

**Remark.** The results in Theorems 1.1–1.7 are similar to those obtained for the standard Navier-Stokes equations, and reveal that the perturbed terms (even if they include the velocity and its first derivatives) in problem (1.1) have little effects on the regularity of (suitable) weak solutions of (1.1), (1.2). This is what the present paper shows to the interested readers.

This paper is organized as follows: Section 2 is devoted to the proofs of Theorems 1.1, 1.2, and we establish regularity criteria for weak solutions of (1.1), (1.2). In Section 3, we focus on the partial regularity of weak solutions of (1.1), (1.2) in the 3D whole space.

## 2. INTERIOR REGULARITY FOR WEAK SOLUTIONS OF (1.1), (1.2)

Set

$$\begin{aligned} C_{0,\sigma}^\infty(\Omega) &= \{u \in C_0^\infty(\Omega) : \operatorname{div} u = 0\}, \\ L_\sigma^2(\Omega) &= \text{the closure of } C_{0,\sigma}^\infty(\Omega) \text{ in } L^2(\Omega), \\ B_r(x_0) &= \{x \in \mathbb{R}^N : |x - x_0| < r\}, \end{aligned}$$

and  $\bar{u}_D = 1/|D| \int_D u(y) \, dy$ .

Consider the problem

$$(2.1) \quad \begin{cases} \partial_t v - \Delta v + \nabla b v = F & \text{in } \mathbb{R}^N \times (0, T), \\ v(x, 0) = 0, \end{cases}$$

where  $v = (v^1, v^2, \dots, v^d)$ ,  $d = N(N-1)/2$ ,  $N \geq 2$ ,  $F = \nabla g + h$ ,  $\nabla g = \left( \sum_{j=1}^N \partial g^{ij} / \partial x_i \right)_{i=1}^d$ ,  $h = (h^1, h^2, \dots, h^d)$ ,  $b = (b_{jk}^i(x, t))_{i=1}^d$  for  $1 \leq i, k \leq d$ ,  $1 \leq j \leq N$ , and  $\nabla b v = \left\{ \sum_{j=1}^N \sum_{k=1}^d (\partial / \partial x_j)(b_{jk}^i(x, t)v^k(x, t)) \right\}_{i=1}^d$ .

The function  $v \in L^{s'}(0, T; L^s(\mathbb{R}^N))$  ( $1 < s, s' \leq \infty$ ) is said to be a *weak solution* of (2.1) if

$$\begin{aligned} & - \int_0^T \int_{\mathbb{R}^N} (\partial_t \varphi + \Delta \varphi + b \nabla \varphi) v \, dx \, dt \\ & = \int_0^T \int_{\mathbb{R}^N} F \varphi \, dx \, dt \quad \text{for any } \varphi \in C_0^\infty(\mathbb{R}^N \times (0, T)). \end{aligned}$$

The following lemma can be found in [30].

**Lemma 2.1.** *Assume that  $1 \leq p, q \leq \infty$  and  $N/p + 2/q = 1$  ( $N \geq 2$ ),  $l, l'$  satisfy  $1/l = 1/p + 1/m$ ,  $1/l' = 1/q + 1/m'$ , where  $2 \leq m \leq r \leq \infty$ ,  $2 \leq m' \leq r' < \infty$  satisfy  $2/m + 2/m' \leq N/r + 2/r' + 1$ . Suppose that  $v \in L^2(0, T; L^2(\mathbb{R}^N))$  is a weak solution of (2.1) with  $F = \nabla g + h$  and  $g, h \in L^2(0, T; L^2(\mathbb{R}^N))$ . There exists a positive constant  $\varepsilon$  such that*

(i) *if  $\|b\|_{L^q(0, T; L^p(\mathbb{R}^N))} \leq \varepsilon$ , then*

$$\|b\|_{L^{r'}(0, T; L^r(\mathbb{R}^N))} \leq C(\|g\|_{L^{m'}(0, T; L^m(\mathbb{R}^N))} + \|h\|_{L^{l'}(0, T; L^l(\mathbb{R}^N))});$$



or

(ii) if  $\|b\|_{L^{(q)}(0,T;L^p(\mathbb{R}^N))} \leq \varepsilon$ , then

$$\|b\|_{L^{(r')}(0,T;L^r(\mathbb{R}^N))} \leq C(\|g\|_{L^{(m')}(0,T;L^m(\mathbb{R}^N))} + \|h\|_{L^{(l')}(0,T;L^l(\mathbb{R}^N))}).$$

Here  $C = C(N, d, T, r, r', m, m')$  and  $\varepsilon = \varepsilon(N, d, p, r')$  if  $p > N$ , and  $\varepsilon = \varepsilon(N, d, r, r')$  if  $p = N$ . The exponent  $r$  should be

- (1)  $1 < r < \infty$  if  $p = N$ ,  $q = \infty$ , or
- (2)  $m' = r'$  and  $1/m = 1/r + 1/N$ .

**P r o o f** of Theorem 1.1. Set  $\omega = \text{curl } u$ . Then  $\omega = (\partial_2 u_3 - \partial_3 u_2, -\partial_1 u_3 + \partial_3 u_1, \partial_1 u_2 - \partial_2 u_1)$  for  $N = 3$ , and  $\omega = \partial_1 u_2 - \partial_2 u_1$  for  $N = 2$ . Moreover,

$$(2.2) \quad \partial_t \omega - \Delta \omega + \text{curl}((u \cdot \nabla)u) = \text{curl } g(v_0(t) \cdot \nabla u + (\omega_0 \times x) \cdot \nabla u - \omega_0 \times u).$$

Note that for  $N = 3$ ,

$$(2.3) \quad \text{curl}(v_0(t) \cdot \nabla u) = \text{curl}\left(\sum_{i=1}^3 v_0^i(t) \partial_i u\right) = (v_0(t) \cdot \nabla) \omega,$$

$$(2.4) \quad \text{curl}((u \cdot \nabla)u) = (u \cdot \nabla) \omega - (\omega \cdot \nabla)u,$$

and

$$(2.5) \quad \begin{aligned} & \text{curl}[(\omega_0 \times x) \cdot \nabla u - \omega_0 \times u] \\ &= \text{curl}[(-x_2 \theta(t) \partial_1 + x_1 \theta(t) \partial_2)u - (-u_2 \theta(t), u_1 \theta(t), 0)] \\ &= ((\omega_0 \times x) \cdot \nabla \omega_1 - \theta(t) \partial_1 u_3, (\omega_0 \times x) \cdot \nabla \omega_2 - \theta(t) \partial_2 u_3, \\ & \quad (\omega_0 \times x) \cdot \nabla \omega_3 + \theta(t) (\partial_2 u_2 + \partial_2 u_2)) \\ & \quad - (-\theta(t) \partial_3 u_1, -\theta(t) \partial_3 u_2, 0) \\ &= (\omega_0 \times x) \cdot \nabla \omega - \omega_0 \times \omega. \end{aligned}$$

Inserting (2.3)–(2.5) into (2.2), we obtain

$$(2.6) \quad \partial_t \omega - \Delta \omega + (u \cdot \nabla) \omega - (\omega \cdot \nabla) u = (v_0(t) \cdot \nabla) \omega + (\omega_0 \times x) \cdot \nabla \omega - \omega_0 \times \omega,$$

which also holds for  $N = 2$ . For any  $(x_0, t_0) \in \Omega \times (0, T)$  we have  $\overline{Q_R(x_0, t_0)} \subset \Omega \times (0, T)$  with some small number  $R > 0$ . Set  $v = \omega \psi$ , where  $\psi \in C_0^\infty(Q_R(x_0, t_0))$  with  $\psi \equiv 1$  in  $Q_{\theta R}(x_0, t_0)$ ,  $\theta \in (0, 1)$ . Then  $v = \omega$  in  $Q_{\theta R}(x_0, t_0)$ , and  $v \in L^{2,2}(\mathbb{R}^N \times (-R^2 + t_0, t_0))$  solves in the sense of distribution

$$\begin{cases} \partial_t v - \Delta v + \nabla b v = F(\omega) & \text{in } \mathbb{R}^N \times (t_0 - R^2, t_0), \\ v(x, t_0 - R^2) = 0 & \text{in } \mathbb{R}^N, \end{cases}$$

where  $b = (b_{jk}^i(x, t))_{i=1}^d$  with  $d = 1$  if  $N = 2$ , and  $d = 3$  if  $N = 3$ ,  $b_{jk}^i(x, t) = u^j \delta_{ik} - u^i \delta_{jk}$ ,  $F(\omega) = \nabla g + h$ ,

$$g = -2\omega \otimes (\nabla \psi) + \psi(v_0(t) + \omega_0(t) \times x) \otimes \omega,$$

and

$$h = -\omega \sum_{i=1}^N \partial_i(\psi v_0^i) - \psi \omega_0 \times \omega + \omega \psi_t + \omega \Delta \psi + \left( \sum_{j,k=1}^N \omega^k b_{jk}^i \frac{\partial \psi}{\partial x_j} \right)_{i=1}^N.$$

Applying Lemma 2.1 and using the assumptions in Theorem 1.1, we get

$$(2.7) \quad \begin{aligned} \|v\|_{L^{r'}(0,T;L^r(\mathbb{R}^N))} &\leq C(\|g\|_{L^{m'}(0,T;L^m(\mathbb{R}^N))} + \|h\|_{L^{l'}(0,T;L^l(\mathbb{R}^N))}) \\ &\leq C\|\omega\|_{L^{m',m}(Q_R)}. \end{aligned}$$

Here  $L^{m',m}(Q_R) = L^{m'}(t_0 - R^2, t_0; L^m(B_R(x_0)))$ . The norm  $\|b\|_{L^q(0,T;L^p(\mathbb{R}^N))}$  in the above formula is sufficiently small, and the exponents  $p, q, l, m, r, l', m', r'$  satisfy the relations in Lemma 2.1.

In particular, by the choice of the cut-off function  $\psi$ , one has from (2.7) that

$$(2.8) \quad \|\omega\|_{L^{r',r}(Q_{R/2})} \leq C\|\omega\|_{L^{m',m}(Q_R)}.$$

*Step 1.* Take  $m = r = m' = 2$ ,  $r' = \beta$  in (2.8),  $\beta \in [2, \infty)$ . Since  $\omega \in L^{2,2}(Q_R)$ , we deduce  $\omega \in L^{\beta,2}(Q_{R/2})$  for any  $\beta \in [2, \infty)$ .

*Step 2.* Take  $m = 2$ ,  $m' = r' = \beta$  in (2.8). Then it follows from (2.8) that  $\omega \in L^{\beta,r_0}(Q_{R/4})$ , where  $r_0 = N/(1 - 2\varepsilon_0/\beta(\beta + \varepsilon_0))$ , and  $\varepsilon_0 > 0$  is a small number. Moreover,  $1 + N/r_0 = 2 - 2\varepsilon_0/\beta(\beta + \varepsilon_0) > N/2$  ( $N = 2, 3$ ) if  $\varepsilon_0 > 0$  is small enough.

*Step 3.* Take  $r = \infty$ ,  $m = r_0$ ,  $m' = \beta$ ,  $r' = \beta + \varepsilon_0 > m'$  in (2.8). Then

$$\frac{N}{m} + \frac{2}{m'} - \left( \frac{N}{r} + \frac{2}{r'} + 1 \right) = \frac{2}{\beta} - \frac{2}{(\beta + \varepsilon_0)} - \frac{2}{\beta(\beta + \varepsilon_0)} = 0.$$

Hence, from (2.8) we obtain  $\omega \in L^{\beta,\infty}(Q_{R/8})$  for any  $\beta \in [2, \infty)$ .

For any open region  $G$  with compact closure in  $\Omega$  we have (see [27])

$$(2.9) \quad u(x, t) = \int_G (\nabla_x H(x - \xi)) \times \omega(\xi, t) \, d\xi + A(x, t),$$

where  $H(x)$  is the fundamental solution of the Laplace equation, and  $A(x, t)$  is harmonic in  $G$ . Moreover, in any region  $S = G \times (T_1, T_2)$  with compact closure in  $\Omega \times (0, T)$ , we also have

$$(2.10) \quad \omega(x, t) = \nabla K * g + B.$$

Here  $K$  is the fundamental solution of the heat equation, which is given by  $K = K(x, t) = (4\pi t)^{-N/2} e^{-|x|^2/(4t)}$  for  $t > 0$ ; and  $K = K(x, t) = 0$  for  $t \leq 0$ . The function  $B = B(x, t)$  is a solution of  $\partial_t B - \Delta B = 0$ , and  $|g| \leq C(|\omega||u| + |\nabla u| + |u|)$ .

Since  $\omega \in L^{\beta, \infty}(Q_{R/8})$  for any  $\beta \in [2, \infty)$ , we infer from (2.9) that  $u, \nabla u \in L^{\beta, \infty}(Q_{R_1})$  for some  $R_1 \in (0, R/8)$ , and  $g \in L^{q', q}(Q_{R_1})$  with  $q, q' \in [2, \infty)$ . As in the proof of the main theorem with non-conservative force in [27], we, together with (2.9), (2.10), obtain  $u, \omega \in L^{\infty, \infty}(Q_{R_2})$  for some  $R_2 \in (0, R_1)$ . Using a standard covering argument, we complete the proof of Theorem 1.1.  $\square$

**P r o o f** of Theorem 1.2. As in the proof of (2.8), we also have

$$\|v\|_{L^{(r')}(0, T; L^r(\mathbb{R}^N))} \leq \|\omega\|_{L^{m', m}(Q_R)}.$$

Note that  $\omega \in L^{2, 2}(Q_R)$  yields  $\omega \in L^{(2), 2}(Q_{R/2})$ . Repeating the proof of Theorem 1.1, we complete the proof of Theorem 1.2.  $\square$

### 3. PROOFS OF THEOREMS 1.3–1.7

**Proof of Theorem 1.3.** We rewrite our hypothesis as follows:

$$(3.1) \quad R^{-2} \iint_{Q_R(x_0, t_0)} (|u|^3 + |u||\pi|) \, dx \, dt \\ + R^{-13/4} \int_{t_0 - R^2}^{t_0} \left( \int_{|x - x_0| < R} |\pi| \, dx \right)^{5/4} \, dt \leq \varepsilon_1.$$

Note that  $Q_{R/2}(a, s) \subset Q_R(x_0, t_0)$  for any  $(a, s) \in Q_{R/2}(x_0, t_0)$ . From (3.1) we have

$$(3.2) \quad R^{-2} \iint_{Q_{R/2}(a, s)} (|u|^3 + |u||\pi|) \, dx \, dt \\ + R^{-13/4} \int_{-R^2/4 + s}^s \left( \int_{B_{R/2}(a)} |\pi| \, dx \right)^{5/4} \, dt \leq \varepsilon_1.$$

Set  $r_n = 2^{-n}R$ ,  $Q^n = Q_{r_n}(a, s)$ . Now we verify by induction that inequalities (3.3) $_n$  and (3.4) $_n$  below hold:

$$(3.3_{n(n \geq 3)}) \quad \left( \oint_{Q^n} |u|^3 \, dx \, dt \right)^{4/3} + r_n^{3/5} \oint_{Q^n} |u||\pi - \bar{\pi}_n| \, dx \, dt \leq \varepsilon_1^{2/3},$$

and

$$(3.4_{n(n \geq 2)}) \quad \sup_{-r^2 + s < t \leq s} \oint_{|x - a| < r_n} |u(x, t)|^2 \, dx + r_n^{-3} \iint_{Q^n} |\nabla u|^2 \, dx \, dt \leq C_0 \varepsilon_1^{2/3}.$$

Here  $\oint_{Q^n} |f| dx dt = |Q^n|^{-1} \iint_{Q^n} |f| dx dt$ , and  $\bar{g}_n = \bar{g}_n(t) = \oint_{|x-a| < r_n} g dx = 1/|B_{r_n}(a)| \int_{|x-a| < r_n} g dx$ .

*Step 1:*  $(3.3)_k$  ( $3 \leq k \leq n$ ) implies  $(3.4)_{n+1}$ . Set  $\phi_n = \chi \psi_n$ , where  $\chi \in C^\infty$  with  $0 \leq \chi \leq 1$  and  $\chi \equiv 1$  on  $Q^2 = Q_{r_2}(a, s)$ ;  $\chi \equiv 0$  out of  $Q_{R/3}(a, s)$ ;  $\psi_n(x, t) = (r_n^2 + s - t)^{-N/2} e^{-|x-a|^2/4(r_n^2+s-t)}$  for  $t < s + r_n^2$ .

Note that

$$\begin{aligned} \partial_t \phi_n + \Delta \phi_n &= 0 \quad \text{on } Q^2, \\ |\partial_t \phi_n + \Delta \phi_n| &\leq C \quad \text{on } Q^1, \\ C_1 r_n^{-3} &\leq \phi_n(x, t) \leq C_2 r_n^{-3} \quad \text{and} \quad |\nabla \phi_n(x, t)| \leq C r_n^{-4}, \quad \forall (x, t) \in Q^n, \\ |\phi_n(x, t)| &\leq C r_k^{-3} \quad \text{and} \quad |\nabla \phi_n(x, t)| \leq C r_k^{-4} \quad \text{for any } (x, t) \in Q^{k-1} \setminus Q^k. \end{aligned}$$

Taking  $\phi = \phi_n$  in the generalized energy inequality, we obtain

$$\sup_{s-r^2 < t \leq s} \oint_{|x-a| < r_{n+1}} |u(x, t)|^2 dx + r_{n+1}^{-3} \iint_{Q^n} |\nabla u|^2 dx dt \leq C(I + II + III),$$

where

$$\begin{aligned} I &= \left| \iint_{Q^1} |u|^2 (\partial_t \phi_n + \Delta \phi_n) dx dt - 2 \iint_{Q^1} |u|^2 (v_0(t) + \omega_0(t) \times x) \cdot \nabla \phi_n dx dt \right|, \\ II &= \iint_{Q^1} |u|^3 |\nabla \phi_n| dx dt, \end{aligned}$$

and

$$III = \left| \iint_{Q^1} \pi u \cdot \nabla \phi_n dx dt \right|.$$

Therefore,

$$\begin{aligned} (3.5) \quad I &\leq C \iint_{Q^1} |u|^2 dx dt + C \sum_{k=1}^n \left[ (r_k^{-4} + r_k^{-3}) \iint_{Q^k} |u|^2 dx dt \right] \\ &\leq C \varepsilon_1^{2/3} + C \sum_{k=1}^n \left[ (r_k^{-4} + r_k^{-3}) r_k^{5/3} \left( \iint_{Q^k} |u|^3 dx dt \right)^{2/3} \right] \\ &\leq C \varepsilon_1^{2/3} + C \sum_{k=1}^n [(r_k^{-4} + r_k^{-3}) r_k^{5/3} (\varepsilon_1^{1/2} r_k^5)^{2/3}] \\ &\leq C \varepsilon_1^{2/3} + C \varepsilon_1^{1/3} \sum_{k=1}^n (r_k + r_k^2) \leq C \varepsilon_1^{2/3}. \end{aligned}$$

The estimates of *II*, *III* are given in [7], namely

$$(3.6) \quad II + III \leq C\varepsilon_1^{2/3}.$$

Combining (3.5) and (3.6), we infer that (3.4)<sub>n+1</sub> ( $n \geq 3$ ) holds.

*Step 2:* (3.4)<sub>k</sub> ( $2 \leq k \leq n$ ) implies (3.3)<sub>n+1</sub> ( $n \geq 2$ ). As in [7], we have

$$\int_{Q^{n+1}} |u|^3 \, dx \, dt \leq C^* \varepsilon_1,$$

where  $C^*$  is independent of  $n$ .

Hence,

$$\left( \int_{Q^{n+1}} |u|^3 \, dx \, dt \right)^{4/3} \leq (C^* \varepsilon_1)^{4/3} \leq \frac{1}{2} \varepsilon_1^{2/3},$$

provided that  $(C^* \varepsilon_1)^{4/3} \leq \frac{1}{2}$ , which is possible if  $\varepsilon_1$  is small enough. The rest of the proof is the same as that in [7] due to the following two facts:

- (i) The generalized energy inequality is not used in Step 2.
- (ii) It is not difficult to verify that

$$\operatorname{div}\{(v_0(t) + \omega_0(t) \times x) \cdot \nabla u - \omega_0(t) \times u\} = 0 \quad \text{in the sense of distribution.}$$

Therefore,  $-\Delta\pi = \operatorname{div} \operatorname{div}(u \otimes u)$  holds in the sense of distribution, and then the pressure  $\pi$  has the same formula as in [7].

From the above arguments, we conclude that for any  $n \geq 2$

$$\oint_{|x-a| < r_n} |u(x, t)|^2 \, dx \leq C_0 \varepsilon_1^{2/3} \quad \text{for each } (a, s) \in Q_{R/2}(x_0, t_0).$$

Then  $|u(a, s)|^2 \leq C\varepsilon_1^{2/3}$  provided that  $(a, s)$  is a Lebesgue point for  $u$ , hence almost everywhere on  $Q_{R/2}(x_0, t_0)$ . The proof of  $\mathcal{F}^{5/3}(S) = 0$  is the same as that in [7], so we omit the details here.  $\square$

**Proof of Theorem 1.4.** For any  $0 < r \leq \frac{1}{2}\varrho$  with  $\varrho \leq R_1$ , a test function  $\phi$  in the generalized energy inequality will be chosen satisfying  $\phi \in C_0^\infty(Q_\varrho(x_0, t_0))$ ,  $0 \leq \phi \leq 1$ , and  $\phi \equiv 1$  on  $Q_r(x_0, t_0)$ ,  $|\nabla\phi| \leq C\varrho^{-1}$ ,  $|\partial_t\phi| + |\Delta\phi| \leq C\varrho^{-2}$ . Then

$$\begin{aligned} & \left| -2 \iint_{Q_\varrho(x_0, t_0)} |u|^2 (v_0(t) + \omega_0(t) \times x) \cdot \nabla\phi \, dx \, dt \right| \\ & \leq C(1 + \varrho)\varrho^{-1} \iint_{Q_\varrho(x_0, t_0)} |u|^2 \, dx \, dt \\ & \leq C\varrho \left( \varrho^{-2} \iint_{Q_\varrho(x_0, t_0)} |u|^3 \, dx \, dt \right)^{2/3}. \end{aligned}$$

The estimates of the other terms on the right-hand side of the generalized energy inequality can be obtained by following the proof of Lemma 5.5 in [7]. So Lemma 5.5 in [7] also holds for the suitable weak solution of (1.1), (1.2). Since the next arguments are the same as those of Proposition 3 in [7], we omit the details here.  $\square$

**P r o o f** of Theorem 1.5. Let  $(u, \pi)$  be a suitable weak solution of (1.1), (1.2) with  $G \triangleq \frac{1}{2} \int_{\mathbb{R}^3} |x| |u_0(x)|^2 dx < \infty$ . Using a sequence of test functions converging to  $\phi \equiv 1$  in the generalized energy inequality, we conclude that for each  $t > 0$

$$(3.7) \quad \frac{1}{2} \int_{\mathbb{R}^3} |u(x, t)|^2 dx + \int_0^t \int_{\mathbb{R}^3} |\nabla u|^2 dy ds \leq \frac{1}{2} \int_{\mathbb{R}^3} |u_0(x)|^2 dx \triangleq E.$$

As in the proofs of Lemmas 8.1, 8.2 in [7], one has for almost all  $t > 0$

$$(3.8) \quad \begin{aligned} \frac{1}{2} \int_{\mathbb{R}^3} |x| |u(x, t)|^2 dx + \int_0^t \int_{\mathbb{R}^3} |x| |\nabla u|^2 dx ds \\ \leq A(t) + \frac{C}{2} \int_0^t \int_{\mathbb{R}^3} (1 + |x|) |u|^2 dx ds, \end{aligned}$$

where  $A(t) = G + CEt^{1/2} + Ct^{1/4}E^{3/2}$ .

Set  $g(t) = \frac{1}{2} \int_0^t \int_{\mathbb{R}^3} (1 + |x|) |u|^2 dx ds$ . Then (3.7), (3.8) imply

$$\frac{d}{dt} g(t) \leq B(t) + Cg(t), \quad \text{where } B(t) = A(t) + E.$$

After a direct calculation, we get for each  $t > 0$

$$(3.9) \quad g(t) \leq e^{Ct} \int_0^t e^{-Cs} B(s) ds.$$

Inserting (3.9) into (3.8), we conclude that

$$\frac{1}{2} \int_{\mathbb{R}^3} |x| |u(x, t)|^2 dx + \int_0^t \int_{\mathbb{R}^3} |x| |\nabla u|^2 dx ds \leq A(t) + Ce^{Ct} \int_0^t e^{-Cs} B(s) ds.$$

The rest of the proof is similar to that of Theorem C in [7], so we omit the details here.  $\square$

**P r o o f** of Theorem 1.6. Define  $v(y, t) = u(x, t)$ , where  $y = x - t\xi$ . Then for any  $\xi \in \mathbb{R}^3$ ,

$$\partial_t v - \Delta v + (v - \xi) \cdot \nabla v + \nabla \pi = (v_0(t) + \omega_0(t) \times (y + t\xi)) \cdot \nabla v - \omega_0(t) \times v.$$

Moreover, as in the proof of Lemma 8.3 in [7], one has for each  $t > 0$

$$(3.10) \quad \begin{aligned} & \psi(t)a(t) - a(0) + \frac{1}{k}(1 - e^{-kB_0(t)}) \\ & \leq -k \int_0^t \psi a \partial_\tau B_0 \, d\tau + \tilde{C} \int_0^t \psi a \partial_\tau B_0 \, d\tau + \theta^2 \int_0^t \psi(\tau) \, d\tau + I(t), \end{aligned}$$

where  $a(t) = \int_{\mathbb{R}^3} \sigma_\varepsilon |v|^2 \, dx$ ,  $B_0(t) = \int_0^t \int_{\mathbb{R}^3} \sigma_\varepsilon |\nabla v|^2 \, dx \, d\tau$ ,  $\psi(t) = e^{-kB_0(t)}$ ,  $\theta = |\xi|$ ,  $k > 0$  is an absolute constant to be chosen later,  $\sigma_\varepsilon(x) = (\varepsilon + |x|^2)^{-1/2}$ ,  $\varepsilon > 0$ , and

$$I(t) = -2 \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 (v_0(\tau) + \omega_0(\tau) \times (y + \tau\xi)) \cdot \nabla \sigma_\varepsilon \, dy \, d\tau.$$

Set  $M \triangleq \sup_{0 \leq t < T} 2\{|v_0(t)| + (1+t)|\omega_0(t)|\} < \infty$ . Then

$$(3.11) \quad \begin{aligned} |I(t)| & \leq 2 \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 (|v_0(\tau)| + |\omega_0(\tau)|(|y| + \tau|\xi|)) |\nabla \sigma_\varepsilon| \, dy \, d\tau \\ & \leq M \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 (1 + |y| + |\xi|) \sigma_\varepsilon^2 \, dy \, d\tau. \end{aligned}$$

Note that

$$(3.12) \quad \begin{aligned} & \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 |y| \sigma_\varepsilon^2(y) \, dy \, d\tau \\ & = \int_0^t \int_{\{|y| \geq 1\}} \psi |v|^2 |y| \sigma_\varepsilon^2(y) \, dy \, d\tau + \int_0^t \int_{\{|y| < 1\}} \psi |v|^2 |y| \sigma_\varepsilon^2(y) \, dy \, d\tau \\ & \leq \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 \, dy \, d\tau + \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 \sigma_\varepsilon^2(y) \, dy \, d\tau \\ & \leq \|v\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2 \int_0^t \psi \, d\tau + \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 \sigma_\varepsilon^2(y) \, dy \, d\tau. \end{aligned}$$

Inserting (3.12) into (3.11), we conclude that

$$(3.13) \quad \begin{aligned} |I(t)| & \leq M(2 + \theta) \int_0^t \int_{\mathbb{R}^3} \psi |v|^2 \sigma_\varepsilon^2(y) \, dy \, d\tau + M \|u\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2 \int_0^t \psi \, d\tau \\ & \leq CM(2 + \theta) \int_0^t \psi (a \partial_\tau B)^{1/2} \, d\tau + Mt \|u\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2 \\ & \leq C^2 \int_0^t \psi a \partial_\tau B \, d\tau + [2M^2(4 + \theta^2) + M \|u\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2] t, \end{aligned}$$

where we have used the fact (see [7]) that  $\int_{\mathbb{R}^3} \sigma_\varepsilon^2 |v|^2 \, dx \leq C(a \partial_\tau B)^{1/2}$ .

Combining (3.10) and (3.13), we obtain

$$\psi(t)a(t) - a(0) + \frac{1}{k}(1 - e^{-kB(t)}) \leq (-k + \tilde{C} + C^2) \int_0^t \psi a \partial_\tau B \, d\tau + D(\theta, u)t = D(\theta, u)t.$$

Here we choose  $k = \tilde{C} + C^2$  and  $D(\theta, u) = 2M^2(4 + \theta^2) + M\|u\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2$ .

Hence,

$$(3.14) \quad a(t) - \frac{1}{k} + \left(\frac{1}{k} - a(0) - D(\theta, u)t\right)e^{kB(t)} \leq 0.$$

Passing to the limit  $\varepsilon \rightarrow 0$  in (3.14), replacing the variable  $y$  by  $x - t\xi$ , and taking  $L_0 = k^{-1}$ , we obtain

$$\int_{\mathbb{R}^3 \times \{t\}} \frac{|u(x, t)|^2}{|x - t\xi|^2} \, dx + (L_0 - L - D(\theta, u)t)e^{1/L \int_0^t \int_{\mathbb{R}^3} |\nabla u(x, \tau)|^2 / |x - t\xi|^2 \, dx \, d\tau} \leq L_0.$$

In particular,

$$(3.15) \quad \int_0^t \int_{\mathbb{R}^3} \frac{|\nabla u(x, \tau)|^2}{|x - t\xi|^2} \, dx \, d\tau < \infty,$$

provided that

$$(3.16) \quad 0 < t < \frac{L_0 - L}{D(\theta, u)} = \frac{L_0 - L}{2M^2(4 + |\xi|^2) + \widetilde{M}(u)},$$

$$\widetilde{M}(u) = M\|u\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2.$$

Note that

$$|a|^2 < \frac{(L_0 - L)s}{2M^2} - \left(4 + \frac{1}{2M}\|u\|_{L^\infty(0,T;L^2(\mathbb{R}^3))}^2\right)s^2$$

is equivalent to

$$\frac{L_0 - L}{D(|\xi|, u)} > s, \quad \xi = s^{-1}a.$$

Thus one can choose  $t \in (s - r^2, s)$  such that (3.16) holds if  $r > 0$  is sufficiently small. Consequently,

$$|x - t\xi| \leq |x - a| + |a - t\xi| = |x - a| + |\xi||s - t| \leq r + |\xi|r^2 \leq 2r \quad \text{if } |\xi|r \leq 1,$$

and so

$$\limsup_{r \rightarrow 0} r^{-1} \iint_{Q_r(a,s)} |\nabla u(x, \tau)|^2 \, dx \, d\tau \leq \frac{1}{4} \limsup_{r \rightarrow 0} \iint_{Q_r(a,s)} \frac{|\nabla u(x, \tau)|^2}{|x - t\xi|^2} \, dx \, d\tau = 0.$$

From Theorem 1.4 we infer that  $u$  is regular at  $(a, s)$ . The proof of  $\mathcal{F}^1(S) = 0$  is the same as that in [7], so we omit the details here.  $\square$



**Proof of Theorem 1.7.** Let  $\delta > 0$  and  $\Omega = \mathbb{R}^3$ . For any  $(x, t) \in \mathbb{R}^3 \times (\delta, T)$  we can find a small number  $R > 0$  such that  $Q_R(x, t) \subset \mathbb{R}^3 \times (\delta/2, T)$ . Thus,

$$\begin{aligned} J(x, t) &= R^{-2} \iint_{Q_R(x, t)} (|u|^3 + |u||\pi|) \, dy \, d\tau + R^{-13/4} \int_{t-R^2}^t \left( \int_{B_R(x)} |\pi| \, dy \right)^{5/4} \, d\tau \\ &\leq C(R) \left( \iint_{Q_R(x, t)} |\nabla u|^2 \, dx \, d\tau \right)^{3/4} \left( \int_{t-R^2}^t \left( \int_{B_R(x)} |u|^2 \, dy \right)^3 \, d\tau \right)^{1/4} \\ &\quad + C(R) \int_{t-R^2}^t \left( \int_{B_R(x)} |u|^2 \, dy \right)^{3/2} \, d\tau + C(R) \iint_{Q_R(x, t)} |\pi|^{3/2} \, dy \, d\tau \\ &\quad + C(R) \left( \iint_{Q_R(x, t)} |\pi|^{3/2} \right)^{5/6} \, d\tau. \end{aligned}$$

Since  $(u, \pi)$  is a suitable weak solution of (1.1), (1.2), we deduce that for a fixed  $R > 0$ ,  $\lim_{|x| \rightarrow \infty} J(x, t) = 0$  for any  $t \in [\delta, T)$ . Therefore, there exists  $K = K(\delta, T, R, u, \pi) > 0$  such that for any  $t \in [\delta, T)$ ,  $J(x, t) \leq \varepsilon_1$  for every  $x \in \mathbb{R}^3$  with  $|x| \geq K$ . Using Theorem 1.3, one can find  $K_1 > K$  such that  $|u(x, t)| \leq C$  for almost all  $(x, t) \in \mathbb{R}^3 \times [\delta, T)$  with  $|x| \geq K_1$ . If  $\Omega$  is an exterior domain in  $\mathbb{R}^3$ , the proof is similar to the above arguments, so we omit the details here.  $\square$

**Acknowledgment.** The author is grateful to the anonymous referee for his or her many suggestive comments, which improve this manuscript greatly.

#### References

- [1] *G. K. Batchelor*: An Introduction to Fluid Dynamics. Cambridge University Press, Cambridge, 1967.
- [2] *H.-O. Bae, H. J. Choe*: A regularity criterion for the Navier-Stokes equations. *Commun. Partial. Differ. Equations* 32 (2007), 1173–1187.
- [3] *H. Beirão da Veiga*: On the smoothness of a class of weak solutions to the Navier-Stokes equations. *J. Math. Fluid Mech.* 2 (2000), 315–323.
- [4] *H. Beirão da Veiga*: A new regularity class for the Navier-Stokes equations in  $\mathbb{R}^n$ . *Chin. Ann. Math., Ser. B* 16 (1995), 407–412.
- [5] *H. Beirão da Veiga*: A sufficient condition on the pressure for the regularity of weak solutions to the Navier-Stokes equations. *J. Math. Fluid Mech.* 2 (2000), 99–106.
- [6] *L. C. Berselli, G. P. Galdi*: Regularity criteria involving the pressure for the weak solutions of the Navier-Stokes equations. *Proc. Am. Math. Soc.* 130 (2002), 3585–3595.
- [7] *L. Caffarelli, R. Kohn, L. Nirenberg*: Partial regularity of suitable weak solutions of the Navier-Stokes equations. *Commun. Pure Appl. Math.* 35 (1982), 771–831.
- [8] *C. Cao, E. Titi*: Regularity criteria for the three-dimensional Navier-Stokes equations. *Indiana Univ. Math. J.* 57 (2008), 2643–2661.
- [9] *D. Chae, H.-J. Choe*: Regularity of solutions to the Navier-Stokes equation. *Electron. J. Differ. Equ.* 5 (1999), 1–7.

- [10] *D. Chae, J. Lee*: Regularity criterion in terms of pressure for the Navier-Stokes equations. *Nonlinear Anal., Theory Methods Appl.* *46* (2001), 727–735.
- [11] *W. Chester*: A general theory for the motion of a body through a fluid at low Reynolds number. *Proc. R. Soc. Lond., Ser. A* *430* (1990), 89–104.
- [12] *Z. Chen, T. Miyakawa*: Decay properties of weak solutions to a perturbed Navier-Stokes system in  $\mathbb{R}^n$ . *Adv. Math. Sci. Appl.* *7* (1997), 741–770.
- [13] *R. Farwig, C. Komo*: Regularity of weak solutions to the Navier-Stokes equations in exterior domains. *NoDEA, Nonlinear Differ. Equ. Appl.* *17* (2010), 303–321.
- [14] *R. Farwig, H. Kozono, H. Sohr*: Local in time regularity properties of the Navier-Stokes equations. *Indiana Univ. Math. J.* *56* (2007), 2111–2132.
- [15] *P. Han*: Regularity of weak solutions to 3D incompressible Navier-Stokes equations. *J. Evol. Equ.* *10* (2010), 195–204.
- [16] *C. He*: Regularity for solutions to the Navier-Stokes equations with one velocity component regular. *Electron J. Differ. Equ.* *29* (2002), 1–13.
- [17] *T. Hishida*: An existence theorem for the Navier-Stokes flow in the exterior of a rotating obstacle. *Arch. Ration. Mech. Anal.* *150* (1999), 307–348.
- [18] *L. Iskauriaz, G. Seregin, V. Shverak*:  $L_{3,\infty}$ -solutions of Navier-Stokes equations and backward uniqueness. *Uspekhi Mat. Nauk.* *58* (2003), 3–44 (In Russian.); translation in *Russ. Math. Surv.* *58* (2003), 211–250.
- [19] *O. Ladyzhenskaya, G. A. Seregin*: On partial regularity of suitable weak solutions to the three-dimensional Navier-Stokes equations. *J. Math. Fluid Mech.* *1* (1999), 356–387.
- [20] *F. Lin*: A new proof of the Caffarelli-Kohn-Nirenberg theorem. *Commun. Pure Appl. Math.* *51* (1998), 241–257.
- [21] *K. Masuda*: Weak solutions of the Navier-Stokes equations. *Tôhoku Math. J., II. Ser.* *36* (1984), 623–646.
- [22] *J. Neustupa, A. Novotný, P. Penel*: An interior regularity of a weak solution to the Navier-Stokes equations in dependence on one component of velocity. *Quaderni di Matematica, Vol. 10: Topics in Mathematical Fluid Mechanics* (G.P. Galdi, R. Rannacher, eds.). 2003, pp. 168–183.
- [23] *J. Neustupa, P. Penel*: Regularity of a suitable weak solution to the Navier-Stokes equations as a consequence of regularity of one velocity component. *Applied Nonlinear Anal.* (A. Sequeira et al., ed.). Kluwer Academic/Plenum Publishers, New York, 1999, pp. 391–402.
- [24] *P. Penel, M. Pokorný*: Some new regularity criteria for the Navier-Stokes equations containing gradient of the velocity. *Appl. Math.* *49* (2004), 483–493.
- [25] *V. Scheffer*: Partial regularity of solutions to the Navier-Stokes equations. *Pac. J. Math.* *66* (1976), 535–552.
- [26] *G. Seregin*: Navier-Stokes equations: almost  $L_{3,\infty}$ -case. *J. Math. Fluid Mech.* *9* (2007), 34–43.
- [27] *J. Serrin*: On the interior regularity of weak solutions of the Navier-Stokes equations. *Arch. Ration. Mech. Anal.* *9* (1962), 187–195.
- [28] *V. A. Solonnikov*: Estimates of the solutions of a nonstationary linearized system of Navier-Stokes equations. *Am. Math. Soc., Transl., II. Sér.* *75* (1968), 1–116.
- [29] *M. Struwe*: On partial regularity results for the Navier-Stokes equations. *Commun. Pure Appl. Math.* *41* (1988), 437–458.
- [30] *S. Takahashi*: On interior regularity criteria for weak solutions of the Navier-Stokes equations. *Manuscr. Math.* *69* (1990), 237–254.
- [31] *Y. Zhou*: Regularity criteria in terms of pressure for the 3-D Navier-Stokes equations in a generic domain. *Math. Ann.* *328* (2004), 173–192.

- [32] *Y. Zhou*: A new regularity criterion for weak solutions to the Navier-Stokes equations. *J. Math. Pures Appl.*, IX. Sér. 84 (2005), 1496–1514.

*Author's address*: *P. Han*, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, P. R. China, e-mail: [pghan@amss.ac.cn](mailto:pghan@amss.ac.cn).