

María Jesús de la Puente

On tropical Kleene star matrices and alcoved polytopes

*Kybernetika*, Vol. 49 (2013), No. 6, 897--910

Persistent URL: <http://dml.cz/dmlcz/143578>

## Terms of use:

© Institute of Information Theory and Automation AS CR, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

# ON TROPICAL KLEENE STAR MATRICES AND ALCOVED POLYTOPES

MARÍA JESÚS DE LA PUENTE

In this paper we give a short, elementary proof of a known result in tropical mathematics, by which the convexity of the column span of a zero-diagonal real matrix  $A$  is characterized by  $A$  being a Kleene star. We give applications to alcoved polytopes, using normal idempotent matrices (which form a subclass of Kleene stars). For a normal matrix we define a norm and show that this is the radius of a hyperplane section of its tropical span.

*Keywords:* tropical algebra, Kleene star, normal matrix, idempotent matrix, alcoved polytope, convex set, norm

*Classification:* 15A80,52C07,15A60

## 1. INTRODUCTION

Tropical algebra (also called max-algebra, extremal algebra, etc.) is a linear algebra performed with the so called tropical operations:  $\max$  (for addition) and  $+$  (for multiplication) — though some variations use  $\min$  instead of  $\max$ , or ordinary multiplication as tropical multiplication. The study of tropical algebra began in the 60's and 70's with the works of Cuninghame-Green, Gondran-Minoux, Vorobyov, Yoeli and K. Zimmermann and has received a fabulous push since the 90's. Today it ramifies into other areas such as algebraic geometry and mathematical analysis. Tropical algebra began as a means to mathematically model processes which involve synchronization of machines. Applications to such practical problems are still pursued today.

A basic problem in tropical algebra is to determine the properties (classical or tropical) of the set  $V$  spanned (by means of tropical operations) by  $m$  given points  $a_1, \dots, a_m$  in  $\mathbb{R}^n$ . The properties of  $V$  follow from the properties of the  $n \times m$  real matrix  $A$  given by the coordinates of the  $a_j$  written in columns. In this setting,  $V$  is denoted  $\text{span}(A)$ . It is always a connected, compact set, and most often it is non-convex, in the classical sense. Convexity-related questions about  $\text{span}(A)$  have drawn the attention of various authors; see [12, 14, 16, 23], as well as [15, 13].

Assume  $m = n$ . Kleene operators (also called Kleene stars or Kleene closures) are well-known in mathematical logic and computer science. For matrices in tropical algebra, Kleene stars (meaning matrices which are Kleene stars of other matrices) form a particularly well-behaved class. They are simply characterized in terms of linear equalities

and inequalities. For a given matrix  $A$ , it is customary for authors to obtain properties of  $A$  (and  $\text{span}(A)$ ) from properties of the directed graph  $G_A$  associated to  $A$ ; see [1, 3, 6, 9, 10, 28]. For example, the tropical (or max-algebraic) principal eigenvalue  $\lambda(A)$  of  $A$  is the maximum cycle mean of  $G_A$ . But if  $A$  is a Kleene star, then properties of  $\text{span}(A)$  follow directly from  $A$ : we need not consider  $G_A$ .

Alcoved polytopes form a very natural class of generally non-regular convex polytopes, including hypercubes. They have been studied in [18, 19, 26]. An alcoved polytope directly arises from a Kleene star matrix.

In this note we prove, by elementary handling of inequalities, the following known result: for any zero-diagonal real matrix  $A$ ,  $A$  is a Kleene star if and only if  $\text{span}(A)$  is convex. Since a certain hyperplane section of  $\text{span}(A)$  is an alcoved polytope, we are able to obtain some applications to these. One application is the possibility of using tropical operations in order to compute the numerous extremals (vertices and pseudoverties) of a given alcoved polytope. Another application is a way to improve the presentation of an alcoved polytope. A third application is the computation of the radius of an alcoved polytope.

## 2. KLEENE STARS, COLUMN SPANS AND NORMAL IDEMPOTENT MATRICES

Write  $\oplus = \max$  and  $\odot = +$ . These are the tropical operations addition and multiplication. For  $n \in \mathbb{N}$ , set  $[n] := \{1, 2, \dots, n\}$ . Let  $\mathbb{R}^{n \times m}$  denote the set of real matrices having  $n$  rows and  $m$  columns. Define tropical sum and product of matrices following the same rules of classical linear algebra, but replacing addition (multiplication) by tropical addition (multiplication). We will never use classical sum or multiplication of matrices, in this note; therefore,  $A \odot B, A \odot A$  will be written  $AB, A^2$ , respectively, for matrices  $A, B$ . Besides, we will never use the classical linear span.

We will write the coordinates of points in  $\mathbb{R}^n$  in columns. Let  $A \in \mathbb{R}^{n \times m}$  and denote by  $a_1, \dots, a_m \in \mathbb{R}^n$  the columns of  $A$ . The *tropical column span* of  $A$  is, by definition,

$$\begin{aligned} \text{span}(A) : &= \{(\lambda_1 + a_1) \oplus \dots \oplus (\lambda_m + a_m) \in \mathbb{R}^n : \lambda_1, \dots, \lambda_m \in \mathbb{R}\} \\ &= \max\{\lambda_1 + a_1, \dots, \lambda_m + a_m : \lambda_1, \dots, \lambda_m \in \mathbb{R}\} \end{aligned} \tag{1}$$

where maxima are computed coordinatewise. For instance,

$$\begin{aligned} \left(3 + \begin{bmatrix} -2 \\ 1 \end{bmatrix}\right) \oplus \left(0 + \begin{bmatrix} 2 \\ 1 \end{bmatrix}\right) &= \begin{bmatrix} 1 \\ 4 \end{bmatrix} \oplus \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}, \\ \text{so that } \begin{bmatrix} 2 \\ 4 \end{bmatrix} &\in \text{span} \begin{bmatrix} -2 & 2 \\ 1 & 1 \end{bmatrix}. \end{aligned}$$

Notice that, by definition, the set  $\text{span}(A)$  is closed under classical addition of the vector  $(\lambda, \dots, \lambda)$ , for  $\lambda \in \mathbb{R}$ . Therefore, a hyperplane section of it, such as  $\text{span}(A) \cap \{x_n = 0\}$  determines  $\text{span}(A)$ .

We will mostly consider real zero-diagonal square matrices, in this paper. The set of such matrices will be denoted  $\mathbb{R}_{zd}^{n \times n}$ . For  $A = (a_{ij}) \in \mathbb{R}_{zd}^{n \times n}$ , consider the matrix  $A_0 = (\alpha_{ij})$ , where

$$\alpha_{ij} = a_{ij} - a_{nj}, \tag{2}$$

whence  $\text{col}(A_0, j) = -a_{nj} + \text{col}(A, j)$ . The columns of  $A_0$  belong to the hyperplane  $\{x_n = 0\}$  and are tropical scalar multiples of the columns of  $A$ , so that

$$\text{span}(A) = \text{span}(A_0). \tag{3}$$

Thus,  $x \in \text{span}(A) \cap \{x_n = 0\}$  if and only if there exist  $\mu_1, \dots, \mu_n \in \mathbb{R}$  such that

$$x_j = \max_{k \in [n]} \{\alpha_{jk} + \mu_k\}, \quad j \in [n - 1], \tag{4}$$

$$0 = \max_{k \in [n]} \mu_k, \tag{5}$$

so that  $x$  is a combination of the columns of  $A_0$  with coefficients  $\mu_j$  (tropically) adding up to zero.

By definition (see [7, 23, 25]),  $A \in \mathbb{R}_{zd}^{n \times n}$  is a *Kleene star* if  $A = A^2$  (i.e.,  $A$  is zero-diagonal and idempotent, tropically). If each diagonal entry of  $A = (a_{ij})$  vanishes, then  $A \leq A^2$ , because for each  $i, j \in [n]$ , we have

$$a_{ij} \leq \max_{k \in [n]} a_{ik} + a_{kj} = (A^2)_{ij}.$$

Therefore, being a Kleene star is characterized by the following  $n$  linear equalities and  $\binom{n}{2} + \binom{n}{3} = \frac{n^3 - n}{6}$  linear inequalities:

$$a_{ii} = 0, \quad a_{ik} + a_{kj} \leq a_{ij}, \quad i, j, k \in [n], \quad \text{card}\{i, j, k\} \geq 2. \tag{6}$$

In particular,  $a_{ik} + a_{ki} \leq 0$ , for  $i, k \in [n]$ .

By definition, an *alcoved polytope*  $\mathcal{P}$  in  $\mathbb{R}^{n-1}$  is a convex polytope defined by inequalities  $c_i \leq x_i \leq b_i$  and  $c_{ik} \leq x_i - x_k \leq b_{ik}$ , for some  $i, k \in [n - 1], i \neq k$ , and  $c_i, b_i, c_{ik}, b_{ik} \in \mathbb{R} \cup \{\pm\infty\}$ . The polytope  $\mathcal{P}$  may have up to  $\binom{2n-2}{n-1}$  extremals (in the sense of classical convexity) and this bound is sharp; see [12]. This is a fast-growing number, since

$$\binom{2n}{n} \simeq \frac{4^n}{\sqrt{\pi n}},$$

as  $n \rightarrow \infty$ , by Stirling’s formula. For instance, for  $n = 10$ ,  $\mathcal{P}$  may have up to 48.620 extremals.

A matrix  $A \in \mathbb{R}_{zd}^{n \times n}$  induces the following (possibly empty!) alcoved polytope in  $\mathbb{R}^{n-1}$

$$C_A := \left\{ x \in \mathbb{R}^{n-1} : \begin{array}{l} a_{in} \leq x_i \leq -a_{ni} \\ a_{ik} \leq x_i - x_k \leq -a_{ki} \end{array} ; i, k \in [n - 1], i \neq k \right\}. \tag{7}$$

Throughout the paper, we *identify*  $\mathbb{R}^{n-1}$  with the hyperplane  $\{x_n = 0\}$  in  $\mathbb{R}^n$ . Our main result is

**Theorem 2.1.** For any  $A \in \mathbb{R}_{zd}^{n \times n}$ , the following are equivalent:

1.  $A$  is a Kleene star,
2.  $C_A = \text{span}(A) \cap \{x_n = 0\}$ .

To prove this theorem we need two lemmas. Given two points  $x, y \in \mathbb{R}^n$ , let  $B \in \mathbb{R}^{n \times 2}$  be the matrix whose columns are  $x$  and  $y$ . The set  $\text{span}(B)$  is called the *tropical segment* joining  $x$  and  $y$  (not to be confused with the tropical line determined by  $x$  and  $y$ ).

**Lemma 2.2.** If  $A \in \mathbb{R}_{zd}^{n \times n}$ , then  $C_A \subseteq \text{span}(A) \cap \{x_n = 0\}$ .

**Proof.** Given  $x = (x_1, \dots, x_{n-1})^t \in C_A$ , write  $x_n = 0$  and consider scalars  $\mu_n = 0$  and  $\mu_i = x_i + a_{ni} \leq 0$ , for  $i \in [n - 1]$ . Then (4) and (5) hold true, due to (2) and to the  $n(n - 1)$  inequalities defining  $C_A$ . Thus,  $x \in \text{span}(A) \cap \{x_n = 0\}$ . □

**Lemma 2.3. (Tropical convexity of  $C_A$ )** If  $A \in \mathbb{R}_{zd}^{n \times n}$ , then  $\text{span}(B) \cap \{x_n = 0\} \subseteq C_A$ , for every  $x, y$  in  $C_A$ .

**Proof.** Assume that  $x, y \in C_A$ . A point  $z$  in  $\text{span}(B) \cap \{x_n = 0\}$  has coordinates  $z_n = 0 = \max\{\lambda, \mu\}$  and

$$z_i = \max\{\lambda + x_i, \mu + y_i\}, \quad i \in [n - 1],$$

for some  $\lambda, \mu \in \mathbb{R}$ .

Say  $\lambda = 0, \mu \leq 0$ ; then

$$x_i \leq \max\{x_i, \mu + y_i\} = z_i \leq \max\{x_i, y_i\}, \quad i \in [n - 1],$$

so that

$$a_{in} \leq z_i \leq -a_{ni}, \quad i \in [n - 1].$$

Moreover, if  $i, k \in [n - 1], i \neq k$ , we have

$$z_i - z_k = \begin{cases} x_i - x_k, & \text{if } x_i = z_i, x_k = z_k, \\ y_i - y_k, & \text{if } \mu + y_i = z_i, \mu + y_k = z_k, \end{cases}$$

and

$$x_i - x_k \leq z_i - z_k = \mu + y_i - x_k \leq y_i - y_k,$$

if  $\mu + y_i = z_i, x_k = z_k$ . In any case, we get

$$a_{ik} \leq z_i - z_k \leq -a_{ki}.$$

□

Now we go to the proof of Theorem 2.1, showing that (i) and (ii) are also equivalent to

3. each column of  $A_0$  belongs to  $C_A$ .

Proof. Recall that  $A_0 = (\alpha_{ij})$ , where  $\alpha_{ij} = a_{ij} - a_{nj}$ . Then, for  $i, j \in [n]$ ,

- (a)  $\alpha_{ni} = 0$ ,  $\alpha_{in} = a_{in}$  and  $\alpha_{ii} = -a_{ni}$ ,
- (b)  $\alpha_{ij} - \alpha_{jj} = a_{ij}$ .

If  $A$  is a Kleene star, then  $a_{ii} = 0$  and  $a_{ik} + a_{kj} \leq a_{ij}$ , so that

- (c)  $a_{in} \leq \alpha_{ij} \leq -a_{ni}$ ,
- (d)  $a_{ik} \leq \alpha_{ij} - \alpha_{kj} = a_{ij} - a_{kj} \leq -a_{ki}$ .

Items (c) and (d) mean precisely that each column of  $A_0$  belongs to  $C_A$ , so we have that 1 is equivalent to 3.

The coordinates  $(x_1, \dots, x_{n-1}, 0)^t$  of a point  $x$  in  $\text{span}(A) \cap \{x_n = 0\}$  satisfy  $x_j = \max_{k \in [n]} \{\alpha_{jk} + \mu_k\}$ , with  $0 = \max_{k \in [n]} \mu_k$ . Say, without loss of generality,  $\mu_1 = 0$  and write

$$x = z \oplus (\mu_3 + \text{col}(A_0, 3)) \oplus \dots \oplus (\mu_n + \text{col}(A_0, n)),$$

with  $z = \text{col}(A_0, 1) \oplus (\mu_2 + \text{col}(A_0, 2))$ . Assuming 3, then  $z$  lies in  $C_A$ , by lemma 2.3. Again by lemma 2.3, in finitely many steps, we show that  $x$  lies in  $C_A$ . Thus, 3 implies 2, by lemma 2.2. And 2 implies 3, because  $\text{span}(A) = \text{span}(A_0)$ .  $\square$

Theorem 2.1 and its proof deal with linear inequalities and maxima, because the equivalence between conditions 1 and 2 can be restated as

$$(6) \Leftrightarrow [x \in C_A \Leftrightarrow \exists \mu_1, \dots, \mu_n \text{ such that (4) and (5)}]$$

and  $x \in C_A$  (see (7)) depends on inequalities.

The convex set  $C_A \subseteq \mathbb{R}^{n-1} = \{x_n = 0\}$  gives rise to another convex subset in  $\mathbb{R}^n$  as follows:  $\overline{C_A} = \{(x, 0) + (\lambda, \dots, \lambda) : x \in C_A, \lambda \in \mathbb{R}\}$ , the Minkowski sum of  $C_A$  and a line. It is obvious that

$$4. \overline{C_A} = \text{span}(A)$$

is equivalent to 2 in theorem 2.1.

Theorem 2.1 (and its equivalent item 4) is closely related to Sergeev’s section 3.1 in [23] (please note that the notation in [23] is multiplicative — i.e.,  $\odot$  is the usual multiplication). In particular, see top of p. 324 and propositions 3.4, 3.5 and 3.6. In terms of that work, we are proving that a zero-diagonal matrix  $A$  is a Kleene star if and only if its column span equals its subeigenvector cone (denoted  $V^*(A)$  in [23] and  $\overline{C_A}$  here). In proposition 3.4 in [23], the assumption is that  $A$  is definite, meaning that  $\lambda(A) = 0$ . In proposition 3.5, the assumption is that  $A$  is strongly definite, meaning that  $\lambda(A) = 0$  and  $a_{ii} = 0$ ,  $i \in [n]$ . There,  $\lambda(A)$  denotes the maximum cycle mean of  $A$ , the cycles referring to the directed graph  $G_A$ . And  $\lambda(A)$  happens to be the unique eigenvalue of  $A$ . Sergeev’s result and proof can also be found in p.26 of [6]. Unlike in [6, 23], we are not using the terminology of max-plus spectral theory or multi-order

convexity to present or explain our main result (although this is possible too). Moreover, we are not assuming anything about  $\lambda(A)$ .

Theorem 2.1 is also related to proposition 3.6 in [26], where a different concept of generating set for an alcoved polytope is considered (please note that in [26],  $\oplus$  means minimum).

A *first application* to alcoved polytopes  $\mathcal{P} \subset \mathbb{R}^{n-1}$  goes as follows. Remember that  $\mathcal{P}$  is a convex set (in the classical sense) having a large number  $s$  of extremals:  $s \leq \binom{2n-2}{n-1}$ . If  $\mathcal{P} = C_A$  for some Kleene star  $A \in \mathbb{R}_{\geq d}^{n \times n}$ , we know that  $\mathcal{P}$  is tropically spanned by the  $n$  columns of  $A_0$ . The columns of  $A_0$  are extremals of  $\mathcal{P}$  of course, the advantage being that the remaining  $s - n$  extremals of  $\mathcal{P}$  can be computed from  $A_0$ , using a tropical algorithm, such as [2]. Some authors call *vertices* to the columns of  $A_0$  and *pseudoverties* to the remaining  $s - n$  extremals of  $\mathcal{P}$ .

**Example 2.4.** The alcoved polytope  $\mathcal{P} \subset \mathbb{R}^2$  (see Figure 1, left) given by

$$-1 \leq x \leq 3, \quad -2 \leq y \leq 6, \quad -4 \leq y - x \leq 5$$

satisfies  $\mathcal{P} = C_A$ , with

$$A = \begin{bmatrix} 0 & -5 & -1 \\ -4 & 0 & -2 \\ -3 & -6 & 0 \end{bmatrix}, \quad A_0 = \begin{bmatrix} 3 & 1 & -1 \\ -1 & 6 & -2 \\ 0 & 0 & 0 \end{bmatrix}.$$

Since  $A = A^2$ , then  $\mathcal{P}$  is spanned by the columns of  $A_0$ . In particular, the three columns of  $A_0$  are extremals of  $\mathcal{P}$ . The other three extremals of  $\mathcal{P}$  are combinations of these. To be precise,

$$\begin{aligned} \begin{bmatrix} 3 \\ 6 \\ 0 \end{bmatrix} &= \begin{bmatrix} 3 \\ -1 \\ 0 \end{bmatrix} \oplus \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} -1 \\ 4 \\ 0 \end{bmatrix} = -2 + \begin{bmatrix} 1 \\ 6 \\ 0 \end{bmatrix} \oplus \begin{bmatrix} -1 \\ -2 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix} \\ &= \begin{bmatrix} -1 \\ -2 \\ 0 \end{bmatrix} \oplus -1 + \begin{bmatrix} 3 \\ -1 \\ 0 \end{bmatrix}. \end{aligned}$$

**Example 2.5.** Let  $\mathcal{P} = C_A \subset \mathbb{R}^3$  (see Figure 2), where

$$A = \begin{bmatrix} 0 & -6 & -10 & -5 \\ -8 & 0 & -5 & -3 \\ -3 & -5 & 0 & -6 \\ -5 & -3 & -6 & 0 \end{bmatrix}, \quad A_0 = \begin{bmatrix} 5 & -3 & -4 & -5 \\ -3 & 3 & 1 & -3 \\ 2 & -2 & 6 & -6 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since  $A = A^2$ , then the columns of  $A_0$  span  $\mathcal{P}$ , i.e, they are extremals of  $\mathcal{P}$  and every other extremal of  $\mathcal{P}$  can be computed tropically from them (as tropical combinations).


It can be checked (with the help of a computer program) that  $C_A$  has  $17 < \binom{6}{3} = 20$  extremals: the coordinates of the remaining 13 extremals are the columns of the matrix

$$\begin{bmatrix} -5 & -3 & 5 & 5 & 1 & 5 & -3 & -3 & -4 & -5 & -5 & -5 & -5 \\ 1 & -1 & 3 & 3 & 3 & 1 & -3 & 3 & 2 & 1 & -1 & 0 & -3 \\ 5 & -6 & 6 & 2 & -2 & 6 & -6 & 6 & 6 & -4 & -6 & 5 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$


Theorem 2.1 deals with Kleene stars, but we prefer to work with a subclass of particularly nice matrices. These are the *normal idempotent matrices* (NI, for short). By definition, a real matrix  $A = (a_{ij})$  is *normal* if  $a_{ii} = 0$ ,  $a_{ij} \leq 0$ , all  $i, j \in [n]$ ; see [6]. Notice that if  $A$  is NI, then  $a_{ik} + a_{kj} \leq a_{ij}$ , for all  $i, j, k \in [n]$ , so that  $A$  is a Kleene star, by (6). The converse is not true; for instance,  $A = \begin{bmatrix} 0 & -2 \\ 1 & 0 \end{bmatrix}$  is a Kleene star but not a normal matrix. A NI matrix  $A$  satisfies  $\lambda(A) = 0$ , although we do not need this.

Clearly,  $A$  is normal if and only if  $C_A$  contains the origin, in which case, by lemma 2.2,  $\text{span}(A)$  does too. Informally speaking, a matrix  $A$  is normal if the columns of  $A_0$  are set around the origin of  $\mathbb{R}^{n-1}$ , and they follow a precise order — and this order is a kind of orientation in  $\mathbb{R}^{n-1}$ .

Due to the *Hungarian method* (see [17, 22]), any order  $n$  real matrix  $A$  can be normalized, meaning that there exist (non necessarily unique) order  $n$  matrices  $P, Q, N$  such that  $N = QAP$  and  $N$  is normal. Moreover,  $\text{span}(N)$  has the same properties of  $\text{span}(A)$ , since multiplication by  $P$  amounts to a relabeling of columns, and multiplication by  $Q$  amounts to performing a translation. (Here are a few words on the properties



**Fig. 1.** Alcoved polytopes in examples 2.4, 2.7 and 2.10. Generators are rounded (in blue), other extremals are squared (in black), the origin is marked (in white).



**Fig. 2.** Alcoved polytope from example 2.5. The columns of  $A_0$  are marked with digits 1, 2, 3 and 4.

of  $P$  and  $Q$ . The matrices  $P$  and  $Q$  are *generalized permutation matrices*. Here we extend  $\mathbb{R}$  to  $\mathbb{R} \cup \{-\infty\}$ . A *diagonal matrix* is  $D = (d_{ij})$  with  $d_{ii} \in \mathbb{R}$  and  $d_{ij} = -\infty$ . In particular, if  $d_{ii} = 0$  for all  $i \in [n]$ , we get a matrix which acts as an *identity* for matrix multiplication, since  $-\infty$  acts as a neutral element for  $\oplus = \max$ . A *generalized permutation matrix* is the result of applying a permutation  $\sigma$  to the rows and columns of a diagonal matrix). We need *not* use matrices over  $\mathbb{R} \cup \{-\infty\}$  in this paper, because when normalizing a given  $A \in \mathbb{R}_{zd}^{n \times n}$ , every instance of  $-\infty$  in the matrices  $P$  and  $Q$  above can be replaced by  $-t$ , for some real number  $t \geq 0$  big enough, yielding real matrices  $P'$  and  $Q'$  with  $N = Q'AP'$ ; see remark 2 in p. 907 for a bound on  $t$ .) The matrix  $N$  can be obtained from  $A$  with  $O(n^3)$  elementary tropical operations (max and +); see [6] and therein.

A pioneer paper dealing with normal matrices is [27] (although another terminology is used there). If  $A$  is normal, then clearly  $A \leq A^2 \leq A^3 \leq \dots$  and Yoeli proved in [27] that  $A^{n-1} = A^n = A^{n+1} = \dots$ , so that  $A^{n-1}$  is NI, so is a Kleene star. Denote this matrix by  $A^*$  and call it *the Kleene star of A*. More generally, for any real square matrix  $A$ , define  $A^*$  as  $A \oplus A^2 \oplus A^3 \oplus \dots$ , if this limit exists in  $\mathbb{R}^{n \times n}$ .

**Lemma 2.6.** If  $A^*$  exists, then  $C_A = C_{A^*}$ .

*Proof.* By the Hungarian method, we may suppose that  $A$  is normal, so that  $A^* = A^{n-1}$ . Clearly,  $C_A \supseteq C_{A^{n-1}}$ , because  $A \leq A^{n-1}$ . To prove the converse, assume that  $A < A^2$ . Then there exist pairwise different  $i, j, k \in [n]$  such that  $a_{ik} < a_{ij} + a_{jk} = \max_s a_{is} + a_{sk}$ . Suppose that  $x \in C_A$ ; then

$$a_{ij} \leq x_i - x_j \leq -a_{ji}, \tag{8}$$

$$a_{kj} \leq x_k - x_j \leq -a_{jk}, \tag{9}$$

$$a_{ik} \leq x_i - x_k \leq -a_{ki}. \tag{10}$$

Subtracting (9) from (8), we get

$$(A^2)_{ik} = a_{ij} + a_{jk} \leq x_i - x_k$$

which improves (10) to

$$(A^2)_{ik} \leq x_i - x_k \leq -a_{ki}.$$

By going through every entry for which  $A$  and  $A^2$  differ and improving the inequalities as we just did, we get  $C_A = C_{A^2}$ . In a finite number of steps, we get the desired result. □

Lemma 2.6 provides a *second application* to alcoved polytopes  $\mathcal{P}$ . A given presentation  $C_A$  of  $\mathcal{P}$  can be improved to a tight presentation  $\mathcal{P} = C_{A^*}$ .

**Example 2.7.** The alcoved polytope  $\mathcal{P} \subset \mathbb{R}^2$  (see Figure 1, center) determined by

$$-1 \leq x \leq 3, \quad -2 \leq y \leq 6, \quad y - x \leq 5$$

gives rise to the matrix

$$A = \begin{bmatrix} 0 & -5 & -1 \\ -\infty & 0 & -2 \\ -3 & -6 & 0 \end{bmatrix}$$

or, in order to have a real matrix, we can write

$$A(t) = \begin{bmatrix} 0 & -5 & -1 \\ -t & 0 & -2 \\ -3 & -6 & 0 \end{bmatrix},$$

for  $t \in \mathbb{R}$  big enough. Now,

$$A(t)^2 = \begin{bmatrix} 0 & -5 & -1 \\ -5 & 0 & -2 \\ -3 & -6 & 0 \end{bmatrix}$$

is idempotent and does not depend on  $t$ . Write  $A(t)^2 = A(t)^* = A^*$ . Then, by lemma 2.6,  $\mathcal{P} = C_{A^*}$  and  $A^*$  describes  $\mathcal{P}$  tightly. Moreover, by theorem 2.1,  $\mathcal{P}$  is spanned by the columns of

$$(A^*)_0 = \begin{bmatrix} 3 & 1 & -1 \\ -2 & 6 & -2 \\ 0 & 0 & 0 \end{bmatrix}.$$

Notice that in the proof of proposition 3.6 of [26], the authors assume that an alcoved polytope  $C_A$  is described by tight inequalities and then they show that  $A$  is a Kleene star (without explicitly mentioning it).

We close this note by pointing out some some nice *features of normal and NI matrices*.

If  $A$  is NI, then the columns of  $(-A^T)_0$  are extremals of  $\text{span}(A) \cap \{x_n = 0\}$ . A proof of this fact is found in [14] for  $n = 4$ , but the proof works in general. This can be checked out in our examples 2.4 and 2.7 (see also the corresponding figures):

$$(-A^T)_0 = \begin{bmatrix} -1 & 2 & 3 \\ 4 & -2 & 6 \\ 0 & 0 & 0 \end{bmatrix}, \quad (-(A(t)^2)^T)_0 = \begin{bmatrix} -1 & 3 & 3 \\ 4 & -2 & 6 \\ 0 & 0 & 0 \end{bmatrix}$$

and in Example 2.5, where the first four columns of the  $4 \times 13$  matrix are precisely the columns of  $(-A^T)_0$ .


For  $p \in \mathbb{R}^n$ , set

$$\|p\| := \max_{i,j \in [n]} \{|p_i|, |p_i - p_j|\}.$$

This is a *seminorm* in  $\mathbb{R}^n$  (meaning that the property  $\|\lambda + p\| = |\lambda| + \|p\|$ , for  $\lambda \in \mathbb{R}$  is not required). The seminorm  $\|\cdot\|$  is *invariant* under the embedding of  $\mathbb{R}^{n-1} \simeq \{x_n = 0\} \subset \mathbb{R}^n$ . It gives rise to a *semidistance* in  $\mathbb{R}^n$  (where the property  $d(p, q) = 0 \Rightarrow p = q$  is not required)

$$d(p, q) := \max_{i,j \in [n]} \{|p_i - q_i|, |p_i - q_i - p_j + q_j|\}. \tag{11}$$

This is a *distance* on the hyperplane  $\mathbb{R}^{n-1} \simeq \{x_n = 0\}$ ! It measures the integer length (or lattice length) of the tropical segment  $\text{span}(p, q)$ . In  $\mathbb{R}^2 \simeq \{x_3 = 0\}$ , for example, we have  $d((-2, -2), (0, 0)) = 2$  (not  $2\sqrt{2}$ !),  $d((-5, -2), (-2, -5)) = \max\{3, 6\} = 6 = 3 + 3$  and  $d(-5, -2), (0, 0) = \max\{5, 2, 3\} = 5 = 3 + 2$ . It is a sort of Manhattan distance; see Figure 3.



**Fig. 3.** Tropical line in  $\mathbb{R}^2$  with vertex at the point  $(-2, -2)$ .

Define the *tropical radius* of a subset  $S \subset \mathbb{R}^{n-1}$  containing the origin, as follows:

$$r(S) := \sup_{s \in S} d(s, 0) = \sup_{s \in S} \|s\|. \tag{12}$$

For a matrix  $A$ , consider

$$|||A||| := \max_{i,j} |a_{ij}|. \tag{13}$$

If  $A$  is normal, then  $a_{ii} = 0$  and  $a_{ij} \leq 0$ , so that  $A \leq A^{n-1}$ , whence  $|||A||| \geq |||A^{n-1}|||$ .

Below we prove that the radius of  $C_A$  equals the norm of  $A$ , for a NI matrix  $A$ .

**Theorem 2.8.** If  $A$  is normal, then  $|||A||| = r(\text{span}(A) \cap \{x_n = 0\})$ . If, in addition,  $A$  is idempotent, then  $|||A||| = r(C_A)$ .

*Proof.* We only need prove the first statement.

We know that  $A_0 = (\alpha_{ij})$ , with  $\alpha_{ij} = a_{ij} - a_{nj}$ . Assume that  $A = (a_{ij})$  is normal (i. e.,  $a_{ii} = 0$  and  $a_{ij} \leq 0$ ). We first prove that

$$|||A||| = \max_{k \in [n]} ||\text{col}(A_0, k)||. \tag{14}$$

To do so, write  $M$  for the maximum on the right hand side. We have

$$M = \max_{i,j,k \in [n]} \{|\alpha_{ik}|, |\alpha_{ik} - \alpha_{jk}|\} = \max_{i,j,k \in [n]} |a_{ik} - a_{jk}|. \tag{15}$$

Using  $a_{ii} = 0$ , we get  $|||A||| \leq M$ . On the other hand, the maximum on the right hand side of (15) cannot be achieved for mutually different  $i, j, k$  since  $a_{ik} \leq 0$  and  $a_{jk} \leq 0$ ; thus we get  $|||A||| = M$ .

From equalities (3) and (14), we obtain  $|||A||| \leq r(\text{span}(A) \cap \{x_n = 0\})$ .

Now, assume that  $p, y$  are two columns of  $A_0$  and let  $z = \lambda + p \oplus \mu + y$ , with  $z_n = 0 = \max\{\lambda, \mu\}$ . Say  $\lambda = 0$ . Then


$$z_j = \max\{p_j, \mu + y_j\} \leq \max\{p_j, y_j\} \leq \max\{|p_j|, |y_j|\} \leq \max\{||p||, ||y||\}.$$

Besides, by the same argument used in the proof of lemma 2.3, we get  $p_i - p_k \leq z_i - z_k \leq y_i - y_k$ , proving that  $||z|| \leq \max\{||p||, ||y||\} \leq M = |||A|||$ . □

**Remark 1.** It is easy to check that (13) defines a matrix norm on  $\mathbb{R}_{z^d}^{n \times n}$  endowed with  $\oplus, \odot$ , but we do not use it here.

**Remark 2.** In the Hungarian method mentioned in p. 903, it is customary to write matrices  $P, Q$  with entries in  $\mathbb{R} \cup \{-\infty\}$ , while  $A, N$  are real. However, every instance of  $-\infty$  in  $P, Q$  can be replaced by  $-t \in \mathbb{R}$ , with  $t \gg |||A|||, |||N|||$ , getting  $P', Q'$  real such that  $N = Q'AP'$ .

**Remark 3.** In [11, 24], the range seminorm  $\tau$  in  $\mathbb{R}^n$  is introduced as follows:  $\tau(p) = \max_{i,j \in [n]} p_i - p_j = \max_{i,j \in [n]} |p_i - p_j|$ . In general,  $\tau(p) \leq ||p||$ . The seminorm  $\tau$  is not invariant under the embedding of  $\mathbb{R}^{n-1} \simeq \{x_n = 0\} \subset \mathbb{R}^n$ . The range seminorm gives rise to a semidistance, used in [8, 24], and denoted  $d_H$ . The distances induced by  $d$  and  $d_H$  on  $\{x_n = 0\}$  coincide. It is a tropical version of Hilbert’s projective distance.



**Fig. 4.** span  $A(7)$  from example 2.7 (in black) and balls of radius 2 and 7 fitting inside and outside (in green).

**Example 2.9.** Let

$$B = \begin{bmatrix} 0 & -6 & -10 & -5 \\ -9 & 0 & -5 & -3 \\ -3 & -5 & 0 & -6 \\ -5 & -3 & -6 & 0 \end{bmatrix},$$

then  $B^2 = A$  of example 2.5 and  $\text{span}(B)$  is not convex. We have  $\|B\| = \|A\| = 10$  so that the sets  $\text{span}(B) \cap \{x_4 = 0\}$  and  $C_A$  have both radius 10.

**Example 2.10.** Returning to example 2.7, the radius of  $\text{span} A(t) \cap \{x_3 = 0\}$  is  $t$ , for  $t \geq 6$ , while the radius of  $C_{A(t)} = C_{A^*}$  is 6. This is clear from Figure 1 right, where the non-convex set  $\text{span} A(t)$  has an arbitrary long “antenna”.

**Remark 4.** In section 4 of [24], Sergeev computes the radius of a  $d_H$ -ball inscribed in  $\text{span}(A)$ . Sergeev computes the biggest ball fitting inside  $\text{span}(A)$  and we compute a ball centered at the origin and containing  $\text{span}(A)$ ; see Figure 4.

**ACKNOWLEDGEMENT**

I am indebted to S. Sergeev for very helpful discussions and to two referees for taking an interest on this paper and pointing out some ways of improving the manuscript.

(Received February 21, 2013)

## REFERENCES

- 
- [1] M. Akian, R. Bapat, and S. Gaubert: Max-plus algebra. In: Handbook of Linear Algebra, Chapter 25, (L. Hobgen, ed.), Chapman and Hall, Boca Raton 2007.
- [2] X. Allamigeon, S. Gaubert, and E. Goubault: Computing the vertices of tropical polyhedra using directed hypergraphs. *Discrete Comput. Geom.* *49* (2013), 247–279.
- [3] F. L. Baccelli, G. Cohen, G. J. Olsder, and J. P. Quadrat: Synchronization and Linearity. John Wiley, Chichester 1992.
- [4] P. Butkovič: Max-algebra: the linear algebra of combinatorics? *Linear Algebra Appl.* *367* (2003), 313–335.
- [5] P. Butkovič: Simple image set of  $(\max, +)$  linear mappings. *Discrete Appl. Math.* *105* (2000), 73–86.
- [6] P. Butkovič: Max-plus Linear Systems: Theory and Algorithms. Springer, Berlin 2010.
- [7] P. Butkovič, H. Schneider, and S. Sergeev: Generators, extremals and bases of max-cones. *Linear Algebra Appl.* *421* (2007), 394–406.
- [8] G. Cohen, S. Gaubert, and J. P. Quadrat: Duality and separation theorems in idempotent semimodules. *Linear Algebra Appl.* *379* (2004), 395–422.
- [9] R. Cuninghame-Green: Minimax algebra. Lecture Notes in Econom. and Math. Systems *166*, Springer, Berlin 1970.
- [10] R. A. Cuninghame-Green: Minimax algebra and applications. In: *Adv. Imag. Electr. Phys.* *90*, (P. Hawkes, ed.), Academic Press, San Diego 1995, pp. 1–121.
- [11] R. A. Cuninghame-Green and P. Butkovič: Bases in max-algebra. *Linear Algebra Appl.* *389* (2004), 107–120.
- [12] M. Develin and B. Sturmfels: Tropical convexity. *Doc. Math.* *9* (2004), 1–27; Erratum in *Doc. Math.* *9* (electronic), (2004), 205–206.
- [13] Z. Izhakian, M. Johnson, and M. Kambites: Pure dimension and projectivity of tropical polytopes. arXiv: 1106.4525v2, 2012.
- [14] A. Jiménez and M. J. de la Puente: Six combinatorial classes of maximal convex tropical polyhedra. arXiv: 1205.4162, 2012.
- [15] M. Johnson and M. Kambites: Idempotent tropical matrices and finite metric spaces. To appear in *Adv. Geom.*; arXiv: 1203.2480, 2012.
- [16] M. Joswig and K. Kulas: Tropical and ordinary convexity combined. *Adv. Geom.* *10* (2010), 333–352.
- [17] H. W. Kuhn: The Hungarian method for the assignment problem. *Naval Res. Logist.* *2* (1955), 83–97.
- [18] T. Lam and A. Postnikov: Alcoved polytopes I. *Discrete Comput. Geom.* *38* (2007), 453–478.
- [19] T. Lam and A. Postnikov: Alcoved polytopes II. arXiv:1202.4015v1, 2012.
- [20] G. L. Litvinov and V. P. Maslov, (eds.): Idempotent Mathematics and Mathematical Physics. Proc. Vienna 2003, Amer. Math. Soc. *Contemp. Math.* *377* (2005).
- [21] G. L. Litvinov and S. N. Sergeev, (eds.): Tropical and Idempotent Mathematics. Proc. Moscow 2007, Amer. Math. Soc. *Contemp. Math.* *495* (2009).

- [22] C. H. Papadimitriou and K. Steiglitz: Combinatorial optimization: algorithms and complexity. Corrected unabridged republication by Dover, Mineola 1998.
- [23] S. Sergeev: Multiorder, Kleene stars and cyclic projectors in the geometry of max cones. In: [21].
- [24] S. Sergeev: Max-plus definite matrix closures and their eigenspaces. *Linear Algebra Appl.* *421* (2007), 182–201.
- [25] S. Sergeev, H. Schneider, and P. Butkovič: On visualization, subeigenvectors and Kleene stars in max algebra. *Linear Algebra Appl.* *431* (2009), 2395–2406.
- [26] A. Werner and J. Yu: Symmetric alcoved polytopes. arXiv: 1201.4378v1, 2012.
- [27] M. Yoeli: A note on a generalization of boolean matrix theory. *Amer. Math. Monthly* *68* (1961) 552–557.
- [28] K. Zimmermann: Extremální algebra. (In Czech.) Výzkumná publikace ekonomicko-matematické laboratoře při ekonomickém ústavu ČSAV, *46*, Prague 1976.

*María Jesús de la Puente, Dpto. de Álgebra, Facultad de Matemáticas, Universidad Complutense, Plaza de Ciencias, 3, 28040-Madrid. Spain.*

*e-mail: mpuente@mat.ucm.es*