Czechoslovak Mathematical Journal

Wu Dejun

G-dimension over local homomorphisms with respect to a semi-dualizing complex

Czechoslovak Mathematical Journal, Vol. 64 (2014), No. 2, 567–579

Persistent URL: http://dml.cz/dmlcz/144016

Terms of use:

© Institute of Mathematics AS CR, 2014

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.

This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

Czechoslovak Mathematical Journal, 64 (139) (2014), 567–579

G-DIMENSION OVER LOCAL HOMOMORPHISMS WITH RESPECT TO A SEMI-DUALIZING COMPLEX

Wu Dejun, Lanzhou

(Received July 20, 2013)

Abstract. We study the G-dimension over local ring homomorphisms with respect to a semi-dualizing complex. Some results that track the behavior of Gorenstein properties over local ring homomorphisms under composition and decomposition are given. As an application, we characterize a dualizing complex for R in terms of the finiteness of the G-dimension over local ring homomorphisms with respect to a semi-dualizing complex.

Keywords: Cohen factorization; Gorenstein dimension; Gorenstein homomorphism; semi-dualizing complex

MSC 2010: 13D02, 13D05, 13D07

1. Introduction

Throughout this paper, all rings are commutative and noetherian. It is well known that Gorenstein homological dimensions are refinements of the classical homological dimensions.

Gorenstein dimension (abbreviation G-dimension), which is a homological invariant for modules, was introduced by Auslander and was deeply studied by Auslander and Bridger in [1]. With that as a start, G-dimension has been studied by a lot of algebraists so far. Two of its main features are that it is a finer invariant than the projective dimension and that it satisfies an equality of the Auslander-Buchsbaum type.

Let C be a semi-dualizing complex for R. Christensen introduced G-dimension with respect to C in [7]. More precisely, for $X \in \mathscr{D}_b^f(R)$ the *G-dimension* of X with

The research has been supported by National Natural Science Foundation of China (No. 11301242).

respect to C is defined to be

$$\text{G-dim}_C X = \begin{cases} \inf C - \inf X^{\dagger C}, & X \in {}_C\mathscr{R}(R), \\ \infty, & X \notin {}_C\mathscr{R}(R) \end{cases}$$

(see [7], 3.11).

Iyengar and Sather-Wagstaff in [11] develop a theory of Gorenstein dimension over local ring homomorphisms. More precisely, let $\varphi\colon R \to S$ be a local ring homomorphism and $X \in \mathscr{D}_b^f(S)$, and let $R \xrightarrow{\dot{\varphi}} R' \xrightarrow{\varphi'} \hat{S}$ be a Cohen factorization of $\dot{\varphi}$. The *Gorenstein dimension of X over φ* is defined by $\text{G-dim}_\varphi X = \text{G-dim}_{R'} \hat{X} - \text{edim}(\dot{\varphi})$.

Motivated by this, it is natural to consider G-dimension over local ring homomorphisms with respect to a semi-dualizing complex. In this paper, G-dimension over local ring homomorphisms with respect to a semi-dualizing complex is studied and the corresponding results are generalized.

Transfer of homological properties along ring homomorphisms is a classical field of study. The following stability result generalizes [11], Theorem 5.1, and [7], Theorem 3.17 (b), (see Theorem 3.15).

Theorem A. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ and $\sigma\colon S \to T$ be local ring homomorphisms. If $P \in \mathscr{D}_b^f(T)$ with $\text{pd}_\sigma P$ finite and $X \in \mathscr{D}_b^f(S)$, then we have the equality*

$$\text{G-dim}_{\sigma\varphi}^C(X \otimes_S^{\mathbf{L}} P) = \text{G-dim}_\varphi^C(X) + \text{pd}_\sigma P.$$

In particular, $\text{G-dim}_{\sigma\varphi}^C(X \otimes_S^{\mathbf{L}} P)$ and $\text{G-dim}_\varphi^C(X)$ are simultaneously finite.

As an application, we have the following result which recovers [11], Theorem 6.1, (see Theorem 4.1).

Theorem B. *Let $(R, \mathfrak{m}, k)$ be a local ring and C a semi-dualizing complex for R. Then the following conditions are equivalent.*

(i) *C is dualizing for R.*

(ii) *For every local ring homomorphism $\varphi\colon R \to S$ and $X \in \mathscr{D}_b^f(S)$, $\text{G-dim}_\varphi^C(X) < \infty$.*

(iii) *There is a local ring homomorphism $\varphi\colon R \to S$ and an ideal I of S such that $I \supseteq \mathfrak{m}S$, and $\text{G-dim}_\varphi^C(S/I) < \infty$.*

2. Preliminaries

The derived category is written as $\mathscr{D}(R)$. If M is an R-complex, then the projective dimension of M is abbreviated as $\operatorname{pd}_R M$. The symbols $\sup M$ and $\inf M$ are used for the supremum and infimum of the set $\{i \in \mathbb{Z};\ \operatorname{H}_i(M) \neq 0\}$, with the conventions $\sup \emptyset = -\infty$ and $\inf \emptyset = \infty$. A complex M is called *homologically bounded above* if $\sup M$ is finite, it is called *homologically bounded below* if $\inf M$ is finite, and it is called *homologically bounded* if it is homologically bounded above and below. The full subcategories $\mathscr{D}_{\sqsubset}(R)$ and $\mathscr{D}_{\sqsupset}(R)$ consist of complexes X with, respectively, $\sup X < \infty$ and $\inf X > -\infty$. We set $\mathscr{D}_b(R) = \mathscr{D}_{\sqsubset}(R) \cap \mathscr{D}_{\sqsupset}(R)$. The full subcategory $\mathscr{P}(R)$ of $\mathscr{D}_b(R)$ consists of complexes of finite projective dimensions. We use the superscript f to denote finite (finitely generated) homology.

We use the standard notation $\mathbf{R}\operatorname{Hom}_R(-,-)$ and $- \otimes_R^{\mathbf{L}} -$ for the derived Hom and derived tensor product of complexes.

2.1 (Depth)**.** Let $(R, \mathfrak{m}, k)$ be a local ring and M an R-complex. The *depth* of M is defined as

$$\operatorname{depth}_R M = -\sup \mathbf{R}\operatorname{Hom}_R(k, M).$$

By [9], 1.5.(3), for every R-complex M one has

$$\operatorname{depth}_R M \geqslant -\sup M.$$

If $\sup M = s$ is finite, then equality holds if and only if $\mathfrak{m}$ is an associated prime of the homology module $\operatorname{H}_s(M)$.

2.2 (Auslander-Buchsbaum formula)**.** If R is local and $X \in \mathscr{P}^f(R)$, then we have the equality

$$\operatorname{pd}_R X = \operatorname{depth} R - \operatorname{depth}_R X.$$

3. G-dimension over a local ring homomorphism

In this section, we introduce the G-dimension over a local ring homomorphism with respect to a semi-dualizing complex and study some of its properties. First, we need to recall the following definitions from [7].

Definition 3.1. An R-complex C is said to be *semi-dulizing* for R if and only if $C \in \mathscr{D}_b^f(R)$ and the homothety morphism $\chi_C^R\colon R \to \mathbf{R}\operatorname{Hom}_R(C, C)$ is an isomorphism (see [7], 2.1).

Let R be a local ring. Recall that a *dualizing complex* for R is a semi-dualizing complex with finite injective dimension. For instance, when R is complete, it possesses a dualizing complex.

Definition 3.2. Let C be a semi-dualizing complex for R. For $X \in \mathscr{D}(R)$ the *dagger dual* with respect to C is the complex $X^{\dagger C} = \mathbf{R}\mathrm{Hom}_R(X, C)$, and $-^{\dagger C} = \mathbf{R}\mathrm{Hom}_R(-, C)$ is the corresponding *dagger duality functor*.

An R-complex X is said to be *C-reflexive* if and only if X and the dagger dual $X^{\dagger C}$ belong to $\mathscr{D}_b^f(R)$, and the biduality morphism $\delta_X^C\colon X \to (X^{\dagger C})^{\dagger C}$ is invertible. By ${}_C\mathscr{R}(R)$ we denote the full subcategory of $\mathscr{D}_b^f(R)$ whose objects are the C-reflexive complexes (see [7], 2.7).

Definition 3.3. Let C be a semi-dualizing complex for R. For $X \in \mathscr{D}_b^f(R)$ the *G-dimension* of X with respect to C is defined to be

$$\text{G-dim}_C X = \begin{cases} \inf C - \inf X^{\dagger C}, & X \in {}_C\mathscr{R}(R), \\ \infty, & X \notin {}_C\mathscr{R}(R) \end{cases}$$

(see [7], 3.11).

The change of rings theorems for Gorenstein dimensions of modules have been investigated in [5]. For complexes of modules, change of rings theorem for G-dimension has been given by Christensen (see [6], Theorem 2.3.12). Here we also have the following change of rings theorem for G-dimension over local homomorphism with respect to a semi-dualizing complex C.

Lemma 3.4. *Let R be a local ring and C a semi-dualizing complex for R. Let $\boldsymbol{x} = x_1, x_2, \ldots, x_t$ be an R-sequence and $S = R/(\boldsymbol{x})$. For $X \in \mathscr{D}_b^f(S)$ there is an equality*

$$\text{G-dim}_C X = \text{G-dim}_{C \otimes_R^{\mathbf{L}} S} X + t.$$

In particular, the two dimensions are simultaneously finite.

Proof. It follows from [7], Proposition 5.7, that $C \otimes_R^{\mathbf{L}} S$ is semi-dualizing for S. Since it is straightforward to prove that $X \in {}_C\mathscr{R}(R)$ if and only if $X \in {}_{C \otimes_R^{\mathbf{L}} S}\mathscr{R}(R)$, one has $\text{G-dim}_C X$ and that $\text{G-dim}_{C \otimes_R^{\mathbf{L}} S} X$ are simultaneously finite. Now the result follows from [7], Theorem 3.14, and the Auslander-Buchsbaum formula for projective dimension (see (2.2)). $\square$

Here we also need to recall the definition of Cohen factorizations of local homomorphisms from [4].

3.5. Let $\varphi\colon (R, \mathfrak{m}, k) \to (S, \mathfrak{n}, l)$ be a local ring homomorphism. The *embedding dimension of φ* is

$$\mathrm{edim}(\varphi) := \mathrm{edim}(S/\mathfrak{m}S).$$

A *regular* or *Gorenstein factorization* of φ is a diagram of local ring homomorphisms, $R \xrightarrow{\dot\varphi} R' \xrightarrow{\varphi'} \hat{S}$, where $\varphi = \varphi'\dot\varphi$, with $\dot\varphi$ flat, the closed fibre $R'/\mathfrak{m}R'$ regular or Gorenstein, respectively, and $\varphi'\colon R' \to S$ surjective.

Let $\hat{S}$ denote the completion of S at its maximal ideal and let $\iota\colon S \to \hat{S}$ be the canonical inclusion. By [4], (1.1), the composition $\grave{\varphi} = \iota\varphi$ admits a regular factorization $R \to R' \to \hat{S}$ with R' complete. Such a regular factorization is called a *Cohen factorization* of $\grave{\varphi}$.

In order to introduce the concept of G-dimension over a local ring homomorphism with respect to a semi-dualizing complex, we also need the following result.

Theorem 3.6. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ be a local ring homomorphism and $X \in \mathscr{D}_b^f(S)$. If $R \xrightarrow{\dot\varphi_1} R_1 \xrightarrow{\varphi_1'} \hat{S}$ and $R \xrightarrow{\dot\varphi_2} R_2 \xrightarrow{\varphi_2'} \hat{S}$ are Cohen factorizations of $\grave{\varphi}$, then we have the equality*

$$\text{G-dim}_{C\otimes_R^{\mathbf{L}} R_1}(\hat{X}) - \text{edim}(\dot\varphi_1) = \text{G-dim}_{C\otimes_R^{\mathbf{L}} R_2}(\hat{X}) - \text{edim}(\dot\varphi_2),$$

where $\hat{X} = X \otimes_S^{\mathbf{L}} \hat{S}$.

Proof. It follows from [7], Theorem 5.6, that $C \otimes_R^{\mathbf{L}} R_1 = C \otimes_R R_1$ is semi-dualizing for R_1 and $C \otimes_R^{\mathbf{L}} R_2 = C \otimes_R R_2$ is semi-dualizing for R_2. Now by analogy with the proof of [11], Theorem 3.2, using Lemma (3.4) one obtains the result. $\square$

Definition 3.7. Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ be a local ring homomorphism and $R \xrightarrow{\dot\varphi} R' \xrightarrow{\varphi'} \hat{S}$ a Cohen factorization of $\grave{\varphi}$. For $X \in \mathscr{D}_b^f(S)$ we define the *Gorenstein dimension of X over φ with respect to C*, $\text{G-dim}_\varphi^C(X)$, as

$$\text{G-dim}_\varphi^C(X) := \text{G-dim}_{C\otimes_R^{\mathbf{L}} R'}(\hat{X}) - \text{edim}(\dot\varphi),$$

where $\hat{X} = X \otimes_S^{\mathbf{L}} \hat{S}$. It follows from [7], Theorem 5.6, that $C \otimes_R^{\mathbf{L}} R' = C \otimes_R R'$ is semi-dualizing for R'. Theorem 3.6 shows that $\text{G-dim}_\varphi^C(X)$ does not depend on the choice of Cohen factorization. Note that $\text{G-dim}_\varphi^C(X) \in \{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$, and also that $\text{G-dim}_\varphi^C(X) = -\infty$ if and only if X is acyclic.

The *Gorenstein dimension of φ with respect to C* is defined to be

$$\text{G-dim}_C(\varphi) := \text{G-dim}_\varphi^C(S).$$

Recall the next definition from [11], Definition 4.2.

Definition 3.8. Let $\varphi\colon R \to S$ be a local ring homomorphism and $R \xrightarrow{\dot{\varphi}} R' \xrightarrow{\varphi'} \hat{S}$ a Cohen factorization of $\dot{\varphi}$. For $X \in \mathscr{D}_b^f(S)$ the *projective dimension of X over φ*, $\operatorname{pd}_\varphi X$, is defined by

$$\operatorname{pd}_\varphi X = \operatorname{pd}_{R'} \hat{X} - \operatorname{edim}(\dot{\varphi}),$$

where $\hat{X} = X \otimes_S^{\mathbf{L}} \hat{S}$. The *projective dimension of φ* is defined to be

$$\operatorname{pd}(\varphi) = \operatorname{pd}_\varphi S.$$

The following proposition shows that G-dimension with respect to a semi-dualizing complex is a refinement of projective dimension over a local ring homomorphism and recovers [7], Proposition 3.15, and [11], Proposition 4.6.

Proposition 3.9. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ be a local ring homomorphism and $X \in \mathscr{D}_b^f(S)$. Then we have the inequality*

$$\text{G-dim}_\varphi^C(X) \leqslant \operatorname{pd}_\varphi X,$$

and equality holds if $\operatorname{pd}_\varphi X < \infty$.

Proof. Let $R \xrightarrow{\dot{\varphi}} R' \xrightarrow{\varphi'} \hat{S}$ be a Cohen factorization of $\dot{\varphi}$. Then we have

$$\begin{aligned}\text{G-dim}_\varphi^C(X) &= \text{G-dim}_{C \otimes_R^{\mathbf{L}} R'}(\hat{X}) - \operatorname{edim}(\varphi)\\ &\leqslant \operatorname{pd}_{R'} \hat{X} - \operatorname{edim}(\varphi)\\ &= \operatorname{pd}_\varphi X\end{aligned}$$

with equality if $\operatorname{pd}_\varphi X < \infty$ (see [7], Proposition 3.15). □

The next theorem is an extension of the Auslander-Bridger formula for G-dimension over a local ring homomorphism (see [11], Theorem 3.5) and for G-dimension with respect to a semi-dualizing complex (see [7], Theorem 3.14), which is a special case by putting $C = R$ and $\varphi = \mathrm{id}_R$ respectively.

Theorem 3.10. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ be a local ring homomorphism and $X \in \mathscr{D}_b^f(S)$. If $\text{G-dim}_\varphi^C(X) < \infty$, then*

$$\text{G-dim}_\varphi^C(X) = \operatorname{depth} R - \operatorname{depth}_S X.$$

Proof. By analogy with the proof of [11], Theorem 3.5, and this time using [7], Theorem 3.14, one obtains the result. □

Lemma 3.11. *Let $\varphi\colon R \to R'$ be a ring homomorphism of finite flat dimension and C a semi-dualizing complex for R. Assume that R' has a dualizing complex D. Then $\mathbf{R}\mathrm{Hom}_R(C, D)$ is a semi-dualizing complex for R'.*

Proof. Note that $\mathbf{R}\mathrm{Hom}_R(C, D) \in \mathscr{D}^f_{\sqsubset}(R')$ by [3], 1.2.2. Since $\mathrm{fd}_R R'$ is finite, one has that $\mathrm{id}_R D$ is finite and so $\mathbf{R}\mathrm{Hom}_R(C, D) \in \mathscr{D}^f_b(R')$. It follows from [6], A.4.24, that

$$\begin{aligned}\mathbf{R}\mathrm{Hom}_{R'}(D, D) &= \mathbf{R}\mathrm{Hom}_{R'}(\mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(C, C), D), D)\\ &= \mathbf{R}\mathrm{Hom}_{R'}(\mathbf{R}\mathrm{Hom}_R(C, D) \otimes^{\mathbf{L}}_R C, D).\end{aligned}$$

Now the commutative diagram

$$\begin{tikzcd} R' \arrow[r] \arrow[d, "\simeq"] & \mathbf{R}\mathrm{Hom}_{R'}(\mathbf{R}\mathrm{Hom}_R(C, D), \mathbf{R}\mathrm{Hom}_R(C, D)) \arrow[d, "\simeq"] \\ \mathbf{R}\mathrm{Hom}_{R'}(D, D) \arrow[r, "\simeq"] & \mathbf{R}\mathrm{Hom}_{R'}(\mathbf{R}\mathrm{Hom}_R(C, D) \otimes^{\mathbf{L}}_R C, D) \end{tikzcd}$$

shows that the homothety morphism

$$\chi^{R'}_{\mathbf{R}\mathrm{Hom}_R(C,D)}\colon R' \to \mathbf{R}\mathrm{Hom}_{R'}(\mathbf{R}\mathrm{Hom}_R(C, D), \mathbf{R}\mathrm{Hom}_R(C, D))$$

is an isomorphism. Therefore, $\mathbf{R}\mathrm{Hom}_R(C, D)$ is a semi-dualizing complex for R'. $\square$

We proceed by recalling the definition of the C-Aslander class from [7], 4.1.

3.12. Let C be a semi-dualizing complex of R. The objects in the *C-Auslander class* ${}_C\mathscr{A}(R)$ are the homologically bounded R-complexes X such that $C \otimes^{\mathbf{L}}_R X$ is homologically bounded and the natural morphism $X \to \mathbf{R}\mathrm{Hom}_R(C, C \otimes^{\mathbf{L}}_R X)$ is an isomorphism.

We list some stability properties of ${}_C\mathscr{A}(R)$.

Proposition 3.13. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ be a local ring homomorphism and X an S-complex. Then the following statements hold.*

(i) *$X \in {}_C\mathscr{A}(R)$ if and only if $X \otimes_S \hat{S} \in {}_{C\otimes_R \hat{R}}\mathscr{A}(\hat{R})$.*
(ii) *If φ is of finite flat dimension, then $X \in {}_C\mathscr{A}(R)$ if and only if $X \in {}_{C\otimes^{\mathbf{L}}_R S}\mathscr{A}(S)$.*
(iii) *If $S \to S'$ is a flat local ring homomorphism, then $X \in {}_C\mathscr{A}(R)$ if and only if $X \otimes^{\mathbf{L}}_S S' \in {}_C\mathscr{A}(R)$.*
(iv) *If $\mathfrak{q} \in \mathrm{Spec} S$ and $\mathfrak{p} = \mathfrak{q} \cap R$, then $X \in {}_C\mathscr{A}(R)$ if and only if $X_{\mathfrak{q}} \in {}_{C_{\mathfrak{p}}}\mathscr{A}(R_{\mathfrak{p}})$.*

P r o o f. If φ is of finite flat dimension, then $C \otimes_R^{\mathbf{L}} S$ is a semi-dualizing complex for S by [7], Proposition 5.7. Let C be a semi-dualizing complex of R. Then $C_{\mathfrak{p}}$ a semi-dualizing complex of $R_{\mathfrak{p}}$ by [7], Lemma 2.5. Now by analogy with the proof of [3], Proposition 3.7, one obtains the result. $\square$

Proposition 3.14. *Let $\varphi\colon R \to S$ be a local ring homomorphism and $R \xrightarrow{\dot{\varphi}} R' \xrightarrow{\varphi'} \hat{S}$ a Cohen factorization of $\dot{\varphi}$, and let $X \in \mathscr{D}_b^f(S)$. Let C be a semi-dualizing complex for R and D a dualizing complex for R'. Then the following conditions are equivalent.*

(i) $\text{G-dim}_{\varphi}^{C}(X) < \infty$.

(ii) $\text{G-dim}_{C \otimes_R^{\mathbf{L}} R'}(\hat{X}) < \infty$.

(iii) $\hat{X} \in {}_{\mathbf{R}\mathrm{Hom}_R(C,D)}\mathscr{A}(R')$.

P r o o f. (i) $\Leftrightarrow$ (ii) By definition.

(ii) $\Leftrightarrow$ (iii) By Lemma (3.11), $\mathbf{R}\mathrm{Hom}_R(C, D)$ is a semi-dualizing complex for R'. Now the result follows from [7], Theorem 4.7, as

$$C \otimes_R^{\mathbf{L}} R' = \mathbf{R}\mathrm{Hom}_{R'}(\mathbf{R}\mathrm{Hom}_R(C, D), D)$$

(see [6], A.4.24). $\square$

The following stability result is one of the main results in this paper which generalizes [11], Theorem 5.1, and [7], [Theorem 3.17 (b)].

Theorem 3.15. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ and $\sigma\colon S \to T$ be local ring homomorphisms. If $P \in \mathscr{D}_b^f(T)$ with $\mathrm{pd}_\sigma P$ finite and $X \in \mathscr{D}_b^f(S)$, then we have the equality*

$$\text{G-dim}_{\sigma\varphi}^{C}(X \otimes_S^{\mathbf{L}} P) = \text{G-dim}_{\varphi}^{C}(X) + \mathrm{pd}_\sigma P.$$

In particular, $\text{G-dim}_{\sigma\varphi}^{C}(X \otimes_S^{\mathbf{L}} P)$ *and* $\text{G-dim}_{\varphi}^{C}(X)$ *are simultaneously finite.*

P r o o f. Note that $X \otimes_S^{\mathbf{L}} P \in \mathscr{D}_b^f(T)$ by [11], Lemma 2.11. Passing to the completions of S and T at their respective maximal ideals, and replacing X and P by $\hat{S} \otimes_S X$ and $\hat{T} \otimes_T P$, respectively, one may assume that S and T are complete. In doing so, one uses the isomorphism

$$(\hat{S} \otimes_S X) \otimes_{\hat{S}}^{\mathbf{L}} (\hat{T} \otimes_T P) \simeq \hat{T} \otimes_T (X \otimes_S^{\mathbf{L}} P).$$

The next step is the reduction to the case where φ and σ are surjective. To achieve this, take Cohen factorizations $R \to R' \to S$ and $R' \to R'' \to T$, and expand

to a commutative diagram as in [11], 5.9. Let $X' = S' \otimes_S X$. Since $S' = R'' \otimes_{R'} S$, by construction, $X' \cong R'' \otimes_{R'} X$ and hence $X' \otimes^{\mathbf{L}}_{S'} P \simeq X \otimes^{\mathbf{L}}_S P$. Since $R' \to R''$ is faithfully flat, [7], Corollary 5.11, yields that

$$\text{G-dim}_{C \otimes^{\mathbf{L}}_R R'}(X) = \text{G-dim}_{C \otimes^{\mathbf{L}}_R R''}(X').$$

Also, in conjunction with those in [11], 5.9, we have

$$\begin{aligned}
\operatorname{pd}_\sigma(P) &= \operatorname{pd}_{S'}(P) - \operatorname{edim}(\dot{\varrho}),\\
\text{G-dim}^C_\varphi(X) &= \text{G-dim}_{C \otimes^{\mathbf{L}}_R R''}(X') - \operatorname{edim}(\dot{\varphi}),\\
\text{G-dim}^C_{\sigma\varphi}(X \otimes^{\mathbf{L}}_S P) &= \text{G-dim}_{C \otimes^{\mathbf{L}}_R R''}(X' \otimes^{\mathbf{L}}_{S'} P) - \operatorname{edim}(\dot{\varrho}) - \operatorname{edim}(\dot{\varphi}).
\end{aligned}$$

Therefore, it suffices to verify the equality for the diagram $R'' \to S' \to T$ and complexes X' and P. This places us in the situation where $R \to S$ is surjective and then the equality we seek is

$$\text{G-dim}_C(X \otimes^{\mathbf{L}}_S P) = \text{G-dim}_C(X) + \operatorname{pd}_S(P).$$

It suffices to prove that $\text{G-dim}_C(X \otimes^{\mathbf{L}}_S P)$ and $\text{G-dim}_C(X)$ are simultaneously finite. For, when they are both finite, one has

$$\begin{aligned}
\text{G-dim}_C(X \otimes^{\mathbf{L}}_S P) &= \operatorname{depth} R - \operatorname{depth}_R(X \otimes^{\mathbf{L}}_S P)\\
&= \operatorname{depth} R - \operatorname{depth}_S(X \otimes^{\mathbf{L}}_S P)\\
&= \operatorname{depth} R - \operatorname{depth}_S X - \operatorname{depth}_S P + \operatorname{depth} S\\
&= \operatorname{depth} R - \operatorname{depth}_R X + \operatorname{pd}_S P\\
&= \text{G-dim}_C(X) + \operatorname{pd}_S P
\end{aligned}$$

where the first and the last equalities follow by the Auslander-Bridger formula (see [7], Theorem 3.14), the second by [11], Lemma 2.8, the third a consequence of [10], Theorem 4.1, while the forth is a consequence of [11], Lemma 2.8, and (2.2).

The rest of the proof is dedicated to proving that $\text{G-dim}_C(X)$ and $\text{G-dim}_C(X \otimes^{\mathbf{L}}_S P)$ are simultaneously finite. This is tantamount to proving that

$$X \in {}_C\mathscr{R}(R) \Leftrightarrow X \otimes^{\mathbf{L}}_S P \in {}_C\mathscr{R}(R).$$

First, note that $X \in \mathscr{D}^f_b(R)$ if and only if $X \otimes^{\mathbf{L}}_S P \in \mathscr{D}^f_b(R)$ by [11], Theorem 2.9. Secondly, since $\operatorname{pd}_S \mathbf{R}\operatorname{Hom}_R(P, S) = -\inf P$ is finite, we have the equalities

$$\begin{aligned}
\mathbf{R}\operatorname{Hom}_R(X \otimes^{\mathbf{L}}_S P, C) &= \mathbf{R}\operatorname{Hom}_S(P, \mathbf{R}\operatorname{Hom}_R(X, C))\\
&= \mathbf{R}\operatorname{Hom}_S(P, S) \otimes^{\mathbf{L}}_S \mathbf{R}\operatorname{Hom}_R(X, C),
\end{aligned}$$

where the first equality follows by adjointness and the second by tensor evaluation (see, for example, [6], A.4.21 and A.4.23). Hence $\mathbf{R}\mathrm{Hom}_R(X,C) \in \mathscr{D}_b^f(R)$ if and only if $\mathbf{R}\mathrm{Hom}_R(X \otimes_S^{\mathbf{L}} P, C) \in \mathscr{D}_b^f(R)$ by virtue of [11], Theorem 2.9.

Finally, since $\mathrm{pd}_S P$ is finite, one has that

$$\delta_P^S\colon\ P \to \mathbf{R}\mathrm{Hom}_S(\mathbf{R}\mathrm{Hom}_S(P,S),S)$$

is an isomorphism. Also, since $\mathrm{pd}_S \mathbf{R}\mathrm{Hom}_R(P,S) = -\inf P$ is finite, we have the equalities

$$\begin{aligned}
&\mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X,C),C) \otimes_S^{\mathbf{L}} P \\
&\quad = \mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X,C),C) \otimes_S^{\mathbf{L}} \mathbf{R}\mathrm{Hom}_S(\mathbf{R}\mathrm{Hom}_S(P,S),S) \\
&\quad = \mathbf{R}\mathrm{Hom}_S(\mathbf{R}\mathrm{Hom}_S(P,S),\mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X,C),C)) \\
&\quad = \mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_S(P,S) \otimes_S^{\mathbf{L}} \mathbf{R}\mathrm{Hom}_R(X,C),C).
\end{aligned}$$

Now the commutative diagram

$$\begin{tikzcd}
X \otimes_S^{\mathbf{L}} P \arrow[r, "(\delta_X^C)\otimes_S^{\mathbf{L}} P"] \arrow[dd, equal] & \mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X,C),C) \otimes_S^{\mathbf{L}} P \arrow[d, "\simeq"] \\
& \mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X,C) \otimes_S^{\mathbf{L}} \mathbf{R}\mathrm{Hom}_S(P,S),C) \\
X \otimes_S^{\mathbf{L}} P \arrow[r, "\delta^C_{(X\otimes_S^{\mathbf{L}} P)}"] & \mathbf{R}\mathrm{Hom}_R(\mathbf{R}\mathrm{Hom}_R(X \otimes_S^{\mathbf{L}} P,C),C) \arrow[u, "\simeq"']
\end{tikzcd}$$

shows that δ_X^C is an isomorphism if and only if $\delta^C_{(X\otimes_S^{\mathbf{L}} P)}$ is an isomorphism. This completes the proof. $\square$

The next result is just the special case arising by taking $X = S$ and $P = T$ in Theorem 3.15 and it generalizes [11], Theorem 5.2, by putting $C = R$.

Corollary 3.16. *Let C be a semi-dualizing complex for R. Let $\varphi\colon\ R \to S$ and $\sigma\colon\ S \to T$ be local homomorphisms with $\mathrm{pd}(\sigma)$ finite. Then*

$$\text{G-dim}_C(\sigma\varphi) = \text{G-dim}_C(\varphi) + \mathrm{pd}(\sigma).$$

In particular, $\text{G-dim}_C(\sigma\varphi)$ is finite if and only if $\text{G-dim}_C(\varphi)$ is finite.

The next stability result generalizes [11], Theorem 5.6, and [7], Theorem 3.17 (a).

Corollary 3.17. *Let C be a semi-dualizing complex for R. Let $\varphi\colon R \to S$ be a local ring homomorphism and $P \in \mathscr{D}_b^f(S)$ with $\operatorname{pd}_S P$ finite. For $X \in \mathscr{D}_b^f(S)$ we have the equality*

$$\text{G-dim}_\varphi^C(\mathbf{R}\operatorname{Hom}_S(P, X)) = \text{G-dim}_\varphi^C(X) - \inf P.$$

Thus, $\text{G-dim}_\varphi^C(X)$ *and* $\text{G-dim}_\varphi^C(\mathbf{R}\operatorname{Hom}_S(P, X))$ *are simultaneously finite.*

Proof. By analogy with the proof of [11], Theorem 5.7, and this time using Theorem 3.15, one obtains the result. □

4. Some applications

In this section, we characterize R and a dualizing complex for R in terms of the finiteness of G-dimension over local ring homomorphisms with respect to a semi-dualizing complex.

Let $\varphi\colon (R, \mathfrak{m}, k) \to (S, \mathfrak{n}, l)$ be a local ring homomorphism. Recall that φ is *Gorenstein* (see [2]), or more precisely, *Gorenstein at* $\mathfrak{n}$, if $\operatorname{fd}_R \varphi < \infty$ and $\mu_R^{i+\operatorname{depth} R} = \mu_S^{i+\operatorname{depth} S}$ for all $i \in \mathbb{Z}$. By [2], 4.2, a flat ring homomorphism is Gorenstein if and only if the ring $S/\mathfrak{m}S$ is Gorenstein.

Applying the next result to $C = R$ we recover [11], Theorem 6.1.

Theorem 4.1. *Let $(R, \mathfrak{m}, k)$ be a local ring and C a semi-dualizing complex for R. Then the following conditions are equivalent.*

(i) *C is dualizing for R.*

(ii) *For every local ring homomorphism $\varphi\colon R \to S$ and $X \in \mathscr{D}_b^f(S)$,* $\text{G-dim}_\varphi^C(X) < \infty$.

(iii) *There is a local ring homomorphism $\varphi\colon R \to S$ and an ideal I of S such that $I \supseteq \mathfrak{m}S$, and* $\text{G-dim}_\varphi^C(S/I) < \infty$.

Proof. (i) $\Rightarrow$ (ii) Let $R \xrightarrow{\dot\varphi} R' \xrightarrow{\varphi'} \hat{S}$ be a Cohen factorization of $\dot\varphi$. Then $\dot\varphi$ is Gorenstein. By [3], 2.11, one has that $C \otimes_R^{\mathbf{L}} R'$ is dualizing for R'. It follows from [7], Proposition 8.4, that ${}_{C\otimes_R^{\mathbf{L}} R'}\mathscr{R}(R') = \mathscr{D}_b^f(R')$ and so $\hat{X} \in {}_{C\otimes_R^{\mathbf{L}} R'}\mathscr{R}(R')$. Therefore, $\text{G-dim}_{C\otimes_R^{\mathbf{L}} R'}(\hat{X}) < \infty$. Thus $\text{G-dim}_\varphi^C(X) < \infty$.

(ii) $\Rightarrow$ (iii) It is trivial.

(iii) $\Rightarrow$ (i) Let $R \to R' \to \hat{S}$ be a Cohen factorization. Composing with the surjection $\hat{S} \xrightarrow{\pi} \hat{S}/I\hat{S}$ gives a diagram $R \to R' \to \hat{S}/I\hat{S}$ that is also a Cohen factorization. Since $\text{G-dim}_{C\otimes_R^{\mathbf{L}} R'}(\hat{S}/I\hat{S})$ is finite, so is $\text{G-dim}_C(\pi\dot\varphi)$. The composition $\pi\dot\varphi$ factors

through the residue field k of R, giving the commutative diagram

$$\begin{tikzcd} R \arrow[rr] \arrow[dr] & & \hat{S}/I\hat{S}. \\ & k \arrow[ur] & \end{tikzcd}$$

The map $k \to \hat{S}/I\hat{S}$ has finite projective dimension as k is a field. Therefore, Corollary (3.16) implies that the surjection $R \to k$ has finite G-dimension with respect to C. Thus, C is dualizing for R by [7], Proposition 8.4. $\square$

The next result generalizes [11], Theorem 6.2.

Theorem 4.2. *Let C be a semi-dualizing complex for R and $\varphi\colon R \to S$ a local ring homomorphism such that S is Gorenstein. Then the following conditions are equivalent.*

(i) *C is dualizing for R.*

(ii) *$\text{G-dim}_C(\varphi)$ is finite.*

(iii) *There exists a complex $P \in \mathscr{D}_b^f(S)$ such that $\operatorname{pd}_S P$ is finite and $\text{G-dim}_\varphi^C(P)$ is finite.*

Proof. (i) $\Rightarrow$ (ii) By Theorem 4.1.

(ii) $\Leftrightarrow$ (iii) By Theorem 3.15.

(ii) $\Rightarrow$ (i) Let $\text{G-dim}_C(\varphi)$ be finite. In particular, $\text{G-dim}(\varphi)$ is finite. By [11], Theorem 6.2, R is Gorenstein. Hence the result follows from [7], Corollary 8.6. $\square$

References

[1] *M. Auslander, M. Bridger*: Stable Module Theory. Mem. Am. Math. Soc. 94, Providence, 1969.

[2] *L. L. Avramov, H.-B. Foxby*: Locally Gorenstein homomorphisms. Am. J. Math. *114* (1992), 1007–1047.

[3] *L. L. Avramov, H.-B. Foxby*: Ring homomorphisms and finite Gorenstein dimension. Proc. Lond. Math. Soc. *75* (1997), 241–270.

[4] *L. L. Avramov, H.-B. Foxby, B. Herzog*: Structure of local homomorphisms. J. Algebra *164* (1994), 124–145.

[5] *D. Bennis, N. Mahdou*: First, second, and third change of rings theorems for Gorenstein homological dimensions. Commun. Algebra *38* (2010), 3837–3850.

[6] *L. W. Christensen*: Gorenstein Dimensions. Lecture Notes in Mathematics 1747, Springer, Berlin, 2000.

[7] *L. W. Christensen*: Semi-dualizing complexes and their Auslander categories. Trans. Am. Math. Soc. *353* (2001), 1839–1883.

[8] *L. W. Christensen, H. Holm*: Ascent properties of Auslander categories. Can. J. Math. *61* (2009), 76–108.

[9] *H.-B. Foxby, S. Iyengar*: Depth and amplitude for unbounded complexes. Commutative Algebra. Interactions with Algebraic Geometry (L. L. Avramov et al., eds.). Contemp. Math. 331, American Mathematical Society, Providence, RI, 2003, pp. 119–137.

[10] *S. Iyengar*: Depth for complexes, and intersection theorems. Math. Z. *230* (1999), 545–567.

[11] *S. Iyengar, S. Sather-Wagstaff*: G-dimension over local homomorphisms. Applications to the Frobenius endomorphism. Illinois J. Math. *48* (2004), 241–272.

Author's address: Wu Dejun, Department of Applied Mathematics, Lanzhou University of Technology, The Langongping Road No. 287, Lanzhou, 730 050, Gansu, China, e-mail: `wudj2007@gmail.com, wudj@lut.cn`.