

Felix Goldberg

A spectral bound for graph irregularity

*Czechoslovak Mathematical Journal*, Vol. 65 (2015), No. 2, 375–379

Persistent URL: <http://dml.cz/dmlcz/144276>

## Terms of use:

© Institute of Mathematics AS CR, 2015

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## A SPECTRAL BOUND FOR GRAPH IRREGULARITY

FELIX GOLDBERG, Haifa

(Received November 3, 2014)

*Abstract.* The imbalance of an edge  $e = \{u, v\}$  in a graph is defined as  $i(e) = |d(u) - d(v)|$ , where  $d(\cdot)$  is the vertex degree. The irregularity  $I(G)$  of  $G$  is then defined as the sum of imbalances over all edges of  $G$ . This concept was introduced by Albertson who proved that  $I(G) \leq 4n^3/27$  (where  $n = |V(G)|$ ) and obtained stronger bounds for bipartite and triangle-free graphs. Since then a number of additional bounds were given by various authors. In this paper we prove a new upper bound, which improves a bound found by Zhou and Luo in 2008. Our bound involves the Laplacian spectral radius  $\lambda$ .

*Keywords:* irregularity; Laplacian matrix; degree; Laplacian index

*MSC 2010:* 05C35, 05C50, 05C07

## 1. INTRODUCTION

Albertson [2] has defined the *irregularity* of a graph  $G$  as:

$$I(G) = \sum_{(u,v) \in E(G)} |d(u) - d(v)|,$$

where  $d(u)$  is the degree of vertex  $u$ . Clearly  $I(G)$  is zero if and only if  $G$  is regular and for non-regular graphs  $I(G)$  is a measure of the defect of regularity. Albertson proved the following upper bound:

$$(1.1) \quad I(G) \leq \frac{4n^3}{27}.$$

Abdo, Cohen and Dimitrov [1] improved Albertson's bound:

$$(1.2) \quad I(G) \leq \left\lfloor \frac{n}{3} \right\rfloor \left\lceil \frac{2n}{3} \right\rceil \left( \left\lceil \frac{2n}{3} \right\rceil - 1 \right).$$

Additional upper bounds on  $I(G)$  have been given by various authors: Hansen and Mélot [5], Henning and Rautenbach [6], Zhou and Luo [10], and Fath-Tabar [3]. These bounds are, strictly speaking, noncomparable but the bound of Zhou and Luo seems to be much sharper than the others for most graphs. We obtain here a new upper bound which is always less than the Zhou-Luo bound or equal to it.

To state the results, let us define the quantity  $Z_G = \sum_{u \in V(G)} d(u)^2$ . It is sometimes called the *first Zagreb index* of  $G$  (cf. [3]).

**Theorem 1.1** ([10], Theorem 1). *Let  $G$  be a graph on  $n$  vertices and with  $m$  edges. Then:*

$$I(G) \leq \sqrt{m(nZ_G - 4m^2)}.$$

Let us now recall the definition of the Laplacian matrix  $L$  of the graph  $G = (V, E)$  whose vertices are labelled  $\{1, 2, \dots, n\}$ :

$$L_{ij} = \begin{cases} -1 & \text{if } (i, j) \in E, \\ 0 & \text{if } (i, j) \notin E \text{ and } i \neq j, \\ -\sum_{k \neq i} L_{ik} & \text{if } i = j. \end{cases}$$

It is obvious from the definition that  $L$  is a positive semidefinite matrix. Surveys of its variegated and fascinating properties can be found in [8], [9]. One simple fact will be germane to us here: the largest eigenvalue  $\lambda_{\max}$  of  $L$  satisfies  $\lambda_{\max} \leq n$ .

We can now state our new result which is clearly an improvement upon the Zhou-Luo bound:

**Theorem 1.2.** *Let  $G$  be a graph on  $n$  vertices and with  $m$  edges. Then:*

$$(1.3) \quad I(G) \leq \sqrt{m(nZ_G - 4m^2)(\lambda_{\max}/n)}.$$

## 2. PROOF OF THE MAIN RESULT

The quadratic form defined by  $L$  has the following useful expression (where we identify the vector  $x \in \mathbb{R}^n$  with a function  $x: V(G) \rightarrow \mathbb{R}$ ):

$$(2.1) \quad x^T Lx = \sum_{(u,v) \in E(G)} (x(u) - x(v))^2.$$

We also need Fiedler's [4] well-known characterization of  $\lambda_{\max}$ :

**Lemma 2.1.**

$$\lambda_{\max} = 2n \max_x \frac{\sum_{(u,v) \in E(G)} (x(u) - x(v))^2}{\sum_{u \in V(G)} \sum_{v \in V(G)} (x(u) - x(v))^2},$$

where  $x$  is a nonconstant vector.

**Proof** of Theorem 1.2. The first step is to apply the Cauchy-Schwarz inequality:

$$(2.2) \quad I(G) = \sum_{(u,v) \in E(G)} |d(u) - d(v)| \leq \sqrt{m} \sqrt{\sum_{(u,v) \in E(G)} (d(u) - d(v))^2}.$$

In light of (2.1) we have:

$$I(G) \leq \sqrt{m} \sqrt{d^T L d}.$$

We now turn to estimate  $d^T L d$  using Lemma 2.1 and Lagrange's identity:

$$\begin{aligned} d^T L d &\leq \frac{\lambda_{\max}}{2n} \sum_{u \in V(G)} \sum_{v \in V(G)} (d(u) - d(v))^2 \\ &= \frac{\lambda_{\max}}{n} \left[ n \sum_{v \in V(G)} d(v)^2 - \left( \sum_{v \in V(G)} d(v) \right)^2 \right]. \end{aligned}$$

Clearly, the latter expression is equal to

$$\frac{\lambda_{\max}}{n} (nZ_G - 4m^2).$$

□

### 3. AN EXAMPLE

Consider the *yoke* graph  $G = Y_{n_1, n_2}$  which consists of two cycles of lengths  $n_1$  and  $n_2$  ( $n_1 + n_2 = n$ ), connected by an edge. It is not hard to see that in this case  $I(G) = 4$ .




Figure 1. The graph  $Y_{7,5}$ .

In order to compare the bounds given by Theorems 1.1 and 1.2 we observe that:

$$m = n + 1, \quad Z_G = 4n + 10.$$

Therefore, the bound given by Theorem 1.1 is:

$$(3.1) \quad I(G) \leq \sqrt{2(n+1)(n-2)} = \Theta(n).$$

To estimate  $\lambda_{\max}$  we can use a result due to Merris [7]. To state it, we define  $m(v)$  to be the average degree of the neighbours of a vertex  $v$ .

**Lemma 3.1** ([7]).  $\lambda_{\max} \leq \max\{d(v) + m(v); v \in V(G)\}$ .

Using this lemma we obtain that  $\lambda_{\max} \leq 16/3$  for any yoke graph and therefore the bound of Theorem 1.2 is at least as good as:

$$(3.2) \quad I(G) \leq \sqrt{\frac{32}{3} \frac{(n+1)(n-2)}{n}} = \Theta(\sqrt{n}).$$

Obviously, (3.2) is much nearer to the true value of  $I(G)$  than (3.1), although it still leaves something to be desired.

#### 4. ANOTHER EXAMPLE—TREES

We wish now to compare Theorem 1.2 to a specialized result of Zhou and Luo which gives a different and very interesting bound for trees.

**Theorem 4.1** ([10], Theorem 4). *Let  $T$  be a tree with  $p$  pendant vertices. Then  $I(G) \leq p(p-1)$ .*

Before reporting a comparison between the two bounds, we wish to point out that by an observation of Albertson ([2], Corollary 5)  $I(G)$  must be an even integer. Therefore, any upper bound on  $I(G)$  can be replaced by the largest even integer not exceeding it.

We have computed the bounds of Theorem 1.2 (truncated to an even integer, as explained above) and of Theorem 4.1 (which always produces even integers) for the 106 non-isomorphic trees on ten vertices. The summary of the results is:

- ▷ For 46 trees Theorem 1.2 is better than Theorem 4.1.
- ▷ For 18 trees both bounds agree.
- ▷ For 42 trees Theorem 4.1 is better than Theorem 1.2.

To take some specific examples, consider first the tree in Figure 2. It is easy to compute that  $I(T_{15}) = 22$ . Since the graph has 7 pendant vertices, Theorem 4.1 yields the estimate  $I(T_{15}) \leq 42$ . On the other hand, the right-hand side of (1.3) is 27.8614 and so Theorem 1.2 gives us the estimate  $I(G) \leq 26$ .




Figure 2. The tree  $T_{15}$ .

On the other hand, if we take the path  $P_n$  on  $n$  vertices, then  $I(G) = 2$  and this value coincides with the bound of Theorem 4.1 since  $p = 2$  in this case. However, Theorem 1.2 gives a poor estimate in this case—we have  $m = n - 1$ ,  $Z_G = 4n - 6$ , and  $\lambda_{\max} = 2(1 + \cos(\pi/n))$ . Since  $\lambda_{\max}$  tends to 4 as  $n$  grows, we can calculate that the right-hand side of (1.3) tends to  $\sqrt{8(n-1)(n-2)/n} = \Theta(\sqrt{n})$ .

#### References

- [1] *H. Abdo, N. Cohen, D. Dimitrov*: Bounds and computation of irregularity of a graph. Preprint: <http://arxiv.org/abs/1207.4804> (2012).
- [2] *M. O. Albertson*: The irregularity of a graph. *Ars Comb.* **46** (1997), 219–225.
- [3] *G. H. Fath-Tabar*: Old and new Zagreb indices of graphs. *MATCH Commun. Math. Comput. Chem.* **65** (2011), 79–84.
- [4] *M. Fiedler*: A property of eigenvectors of nonnegative symmetric matrices and its application to graph theory. *Czech. Math. J.* **25** (1975), 619–633.
- [5] *P. Hansen, H. M elot*: Variable neighborhood search for extremal graphs. IX: Bounding the irregularity of a graph. *Graphs and Discovery* (S. Fajtlowicz et al., eds.). Proc. DIMACS working group, Piscataway, USA, 2001. DIMACS Ser. Discrete Math. Theoret. Comput. Sci. **69**, AMS, Providence, 2005, pp. 253–264.
- [6] *M. A. Henning, D. Rautenbach*: On the irregularity of bipartite graphs. *Discrete Math.* **307** (2007), 1467–1472.
- [7] *R. Merris*: A note on Laplacian graph eigenvalues. *Linear Algebra Appl.* **285** (1998), 33–35.
- [8] *R. Merris*: Laplacian matrices of graphs: A survey. *Linear Algebra Appl.* **197–198** (1994), 143–176.
- [9] *B. Mohar*: Some applications of Laplace eigenvalues of graphs. *Graph Symmetry: Algebraic Methods and Applications* (G. Hahn et al., eds.). Proc. NATO Adv. Study Inst., Montr al, Canada, 1996. NATO ASI Ser., Ser. C, Math. Phys. Sci. **497**, Kluwer Academic Publishers, Dordrecht, 1997, pp. 225–275.
- [10] *B. Zhou, W. Luo*: On irregularity of graphs. *Ars Comb.* **88** (2008), 55–64.

*Author’s address:* Felix Goldberg, Caesarea-Rothschild Institute, University of Haifa, Mount Carmel, Haifa 31905, Israel, e-mail: [felix.goldberg@gmail.com](mailto:felix.goldberg@gmail.com).