

Sheza M. El-Deeb; Georgia I. Oros

Fuzzy differential subordinations connected with the linear operator

Mathematica Bohemica, Vol. 146 (2021), No. 4, 397–406

Persistent URL: <http://dml.cz/dmlcz/149256>

Terms of use:

© Institute of Mathematics AS CR, 2021

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

FUZZY DIFFERENTIAL SUBORDINATIONS CONNECTED WITH
THE LINEAR OPERATOR

SHEZA M. EL-DEEB, New Damietta, GEORGIA I. OROS, Oradea

Received November 15, 2019. Published online November 2, 2020.

Communicated by Dagmar Medková

Abstract. We obtain several fuzzy differential subordinations by using a linear operator $\mathcal{I}_{m,\gamma}^{n,\alpha} f(z) = z + \sum_{k=2}^{\infty} (1 + \gamma(k-1))^n m^\alpha (m+k)^{-\alpha} a_k z^k$. Using the linear operator $\mathcal{I}_{m,\gamma}^{n,\alpha}$, we also introduce a class of univalent analytic functions for which we give some properties.

Keywords: fuzzy differential subordination; fuzzy best dominant; linear operator

MSC 2020: 30C45

1. INTRODUCTION

Let $\mathcal{D} \subset \mathbb{C}$, $\mathcal{H}(\mathcal{D})$ be the class of holomorphic functions on \mathcal{D} and denote by $\mathcal{H}_n(\mathcal{D})$ the class of holomorphic and univalent functions on \mathcal{D} . In this paper, we denote by $\mathcal{H}(\mathbb{U})$ the class of holomorphic functions in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ with $\partial\mathbb{U} = \{z \in \mathbb{C} : |z| = 1\}$ the boundary of the unit disc. For $a \in \mathbb{C}$ and $k \in \mathbb{N}$, we denote

$$\begin{aligned} \mathcal{H}[a, k] &= \{f \in \mathcal{H}(\mathbb{U}) : f(z) = a + a_k z^k + a_{k+1} z^{k+1} + \dots, z \in \mathbb{U}\}, \\ \mathcal{A}_k &= \{f \in \mathcal{H}(\mathbb{U}) : f(z) = z + a_{k+1} z^{k+1} + \dots, z \in \mathbb{U}\} \text{ with } \mathcal{A}_1 = \mathcal{A}, \end{aligned}$$

and

$$\mathcal{S} = \{f \in \mathcal{A} : f \text{ is a univalent function in } \mathbb{U}\}.$$

Denote by

$$\mathcal{K} = \left\{ f \in \mathcal{A} : \Re \left(1 + \frac{z f''(z)}{f'(z)} \right) > 0, z \in \mathbb{U} \right\}$$

the class of convex functions in \mathbb{U} .

Definition 1.1 ([7] and [13]). Let f and g be analytic functions in \mathbb{U} . We say that f is *subordinate* to g , written $f \prec g$, if there exists a Schwarz function w , which is analytic in \mathbb{U} with $w(0) = 0$ and $|w(z)| < 1$ for all $z \in \mathbb{U}$, such that $f(z) = g(w(z))$. Furthermore, if the function g is univalent in \mathbb{U} , then we have the following equivalence:

$$f(z) \prec g(z) \Leftrightarrow f(0) = g(0) \text{ and } f(\mathbb{U}) \subset g(\mathbb{U}).$$

In order to introduce the notion of fuzzy differential subordination, we use the following definitions and propositions:

Definition 1.2 ([10]). Assume that \mathcal{Y} be a nonempty set. An application $\mathcal{F}: \mathcal{Y} \rightarrow [0, 1]$ is called *fuzzy subset*. A pair $(\mathcal{B}, \mathcal{F}_{\mathcal{B}})$, where $\mathcal{F}_{\mathcal{B}}: \mathcal{Y} \rightarrow [0, 1]$ and

$$(1.1) \quad \mathcal{B} = \{x \in \mathcal{Y}: 0 < \mathcal{F}_{\mathcal{B}}(x) \leq 1\} = \text{sup}(\mathcal{B}, \mathcal{F}_{\mathcal{B}}),$$

is called *fuzzy set*. The function $\mathcal{F}_{\mathcal{B}}$ is called *membership function* of the fuzzy set $(\mathcal{B}, \mathcal{F}_{\mathcal{B}})$.

Proposition 1.1 ([14]).

- (i) If $(\mathcal{B}, \mathcal{F}_{\mathcal{B}}) = (\mathcal{U}, \mathcal{F}_{\mathcal{U}})$, then we have $\mathcal{B} = \mathcal{U}$, where $\mathcal{B} = \text{sup}(\mathcal{B}, \mathcal{F}_{\mathcal{B}})$ and $\mathcal{U} = \text{sup}(\mathcal{U}, \mathcal{F}_{\mathcal{U}})$.
- (ii) If $(\mathcal{B}, \mathcal{F}_{\mathcal{B}}) \subseteq (\mathcal{U}, \mathcal{F}_{\mathcal{U}})$, then we have $\mathcal{B} \subseteq \mathcal{U}$, where $\mathcal{B} = \text{sup}(\mathcal{B}, \mathcal{F}_{\mathcal{B}})$ and $\mathcal{U} = \text{sup}(\mathcal{U}, \mathcal{F}_{\mathcal{U}})$.

Let $f, g \in \mathcal{H}(\mathcal{D})$. We denote

$$(1.2) \quad f(\mathcal{D}) = \{f(z): 0 < \mathcal{F}_{f(\mathcal{D})}f(z) \leq 1, z \in \mathcal{D}\} = \text{sup}(f(\mathcal{D}), \mathcal{F}_{f(\mathcal{D})})$$

and

$$(1.3) \quad g(\mathcal{D}) = \{g(z): 0 < \mathcal{F}_{g(\mathcal{D})}g(z) \leq 1, z \in \mathcal{D}\} = \text{sup}(g(\mathcal{D}), \mathcal{F}_{g(\mathcal{D})}).$$

Definition 1.3 ([14]). Let $z_0 \in \mathcal{D}$ and $f, g \in \mathcal{H}(\mathcal{D})$. The function f is said to be *fuzzy subordinate* to g , written $f \prec_{\mathcal{F}} g$ or $f(z) \prec_{\mathcal{F}} g(z)$, when following conditions are satisfied:

- (i) $f(z_0) = g(z_0)$,
- (ii) $\mathcal{F}_{f(\mathcal{D})}f(z) \leq \mathcal{F}_{g(\mathcal{D})}g(z), z \in \mathcal{D}$.

Proposition 1.2 ([14]). Assume that $z_0 \in \mathcal{D}$ and $f, g \in \mathcal{H}(\mathcal{D})$. If $f(z) \prec_{\mathcal{F}} g(z)$, $z \in \mathcal{D}$, then

- (i) $f(z_0) = g(z_0)$,
- (ii) $f(\mathcal{D}) \subseteq g(\mathcal{D}), \mathcal{F}_{f(\mathcal{D})}f(z) \leq \mathcal{F}_{g(\mathcal{D})}g(z), z \in \mathcal{D}$, where $f(\mathcal{D})$ and $g(\mathcal{D})$ are defined by (1.2) and (1.3), respectively.

Definition 1.4 ([16]). Assume that $\Phi: \mathbb{C}^3 \times \mathbb{U} \rightarrow \mathbb{C}$ and $h \in \mathcal{S}$ with $\Phi(a, 0, 0; 0) = h(0) = a$. If p is analytic in \mathbb{U} with $p(0) = a$ and satisfies the second order fuzzy differential subordination

$$(1.4) \quad \mathcal{F}_{\Phi(\mathbb{C}^3 \times \mathbb{U})} \Phi(p(z), zp'(z), z^2p''(z); z) \leq \mathcal{F}_{h(\mathbb{U})} h(z),$$

i.e. $\Phi(p(z), zp'(z), z^2p''(z); z) \prec_{\mathcal{F}} h(z), \quad z \in \mathbb{U},$

then p is called a *fuzzy solution* of the fuzzy differential subordination. The univalent function q is called a *fuzzy dominant* of the fuzzy solutions for the fuzzy differential subordination if

$$\mathcal{F}_{p(\mathbb{U})} p(z) \leq \overline{\mathcal{F}_{q(\mathbb{U})} q(z)}, \quad \text{i.e. } p(z) \prec_{\mathcal{F}} q(z), \quad z \in \mathbb{U}$$

for all p satisfying (1.4). A fuzzy dominant \tilde{q} that satisfies

$$\mathcal{F}_{\tilde{q}(\mathbb{U})} \tilde{q}(z) \leq \mathcal{F}_{q(\mathbb{U})} q(z), \quad \text{i.e. } \tilde{q}(z) \prec_{\mathcal{F}} q(z), \quad z \in \mathbb{U}$$

for all fuzzy dominants q of (1.4) is said to be the *fuzzy best dominant* of (1.4).

The *integral operator* $\mathcal{K}_m^\alpha: \mathcal{A} \rightarrow \mathcal{A}$ is defined for $\alpha > 0$ and $m > -1$ as follows (see Komatu [12] with $p = 1$):

$$\mathcal{K}_m^\alpha f(z) = f(z) = z + \sum_{k=1}^{\infty} a_{k+1} z^{k+1}$$

and

$$(1.5) \quad \mathcal{K}_m^\alpha f(z) = \frac{(m+1)^\alpha}{\Gamma(\alpha)z^m} \int_0^z t^{m-1} \left(\log \frac{z}{t}\right)^{\alpha-1} f(t) dt.$$

It can be easily verified that

$$(1.6) \quad \mathcal{K}_m^\alpha f(z) = z + \sum_{k=1}^{\infty} \left(\frac{m+1}{m+k+1}\right)^\alpha a_{k+1} z^{k+1}.$$

El-Ashwah et al. (see [9] with $p = 1$) introduced the *linear multiplier operator* $\mathcal{I}_{m,\gamma}^{n,\alpha}: \mathcal{A} \rightarrow \mathcal{A}$ given as:

$$\begin{aligned} \mathcal{I}_{m,\gamma}^{0,0} f(z) &= f(z), \\ \mathcal{I}_{m,\gamma}^{1,\alpha} f(z) &= \mathcal{I}_{m,\gamma}^\alpha f(z) = (1-\gamma)\mathcal{K}_m^\alpha f(z) + \gamma z(\mathcal{K}_m^\alpha f(z))' \\ &= z + \sum_{k=1}^{\infty} (1+\gamma k) \left(\frac{m+1}{m+k+1}\right)^\alpha a_{k+1} z^{k+1}, \\ \mathcal{I}_{m,\gamma}^{2,\alpha} f(z) &= \mathcal{I}_{m,\gamma}^\alpha (\mathcal{I}_{m,\gamma}^{1,\alpha} f(z)) \\ &= z + \sum_{k=1}^{\infty} (1+\gamma k)^2 \left(\frac{m+1}{m+k+1}\right)^\alpha a_{k+1} z^{k+1}, \end{aligned}$$

in general,

$$(1.7) \quad \mathcal{I}_{m,\gamma}^{n,\alpha} f(z) = \mathcal{I}_{m,\gamma}^\alpha (\mathcal{I}_{m,\gamma}^{n-1,\alpha} f(z)) = z + \sum_{k=1}^{\infty} (1 + \gamma k)^n \left(\frac{m+1}{m+k+1} \right)^\alpha a_{k+1} z^{k+1},$$

$$\alpha \geq 0, m > -1, \gamma \geq 0, n \in \mathbb{N}_0.$$

It is easily verified from (1.7) that

$$(1.8) \quad z(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' = (m+1)\mathcal{I}_{m,\gamma}^{n,\alpha-1} f(z) - m\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)$$

and

$$(1.9) \quad \gamma z(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' = \mathcal{I}_{m,\gamma}^{n+1,\alpha} f(z) - (1-\gamma)\mathcal{I}_{m,\gamma}^{n,\alpha} f(z).$$

By specializing the parameters m, γ, α and n , we obtain the following operators:

$$\begin{aligned} \mathcal{I}_{c,\gamma}^{0,\delta} f(z) &= \mathcal{K}_{c,1}^\delta f(z) && \text{(see [11]);} \\ \mathcal{I}_{a-1,\gamma}^{0,\alpha} f(z) &= L_a^\alpha f(z) && \text{(see [4] and [5]);} \\ \mathcal{I}_{c,0}^{1,\alpha} f(z) &= \mathcal{P}_c^\alpha f(z) && \text{(see [12] and [17]);} \\ \mathcal{I}_{c,0}^{1,1} f(z) &= \mathcal{L}_c f(z) && \text{(see [6]);} \\ \mathcal{I}_{1,0}^{1,\alpha} f(z) &= \mathcal{I}^\alpha f(z) && \text{(see [8]);} \\ \mathcal{I}_{m,\gamma}^{n,0} f(z) &= \mathcal{D}_\gamma^n f(z) && \text{(see [1]);} \\ \mathcal{I}_{m,1}^{n,0} f(z) &= \mathcal{D}^n f(z) && \text{(see [18]).} \end{aligned}$$

Also, we note that

$$(1.10) \quad \mathcal{I}_{m,\gamma}^{-n,\alpha} f(z) = z + \sum_{k=1}^{\infty} \left(\frac{1}{1+\gamma k} \right)^n \left(\frac{m+1}{m+k+1} \right)^\alpha a_{k+1} z^{k+1}.$$

From (1.7) and (1.10) we observe that the operator $\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)$ is well defined from $n \in \mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$.

By using the linear operator $\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)$ defined by (1.7), we will derive several fuzzy differential subordinations for this class.

Definition 1.5. A function $f \in \mathcal{A}$ belongs to the class $\mathcal{M}_{m,\gamma}^F(n, \alpha, \eta)$ for all $\eta \in [0, 1)$, $n \in \mathbb{N}_0$, $m > -1$, $\gamma \geq 0$ and $\alpha \geq 0$ if it satisfies the inequality

$$F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' > \eta, \quad z \in \mathbb{U}.$$

2. PRELIMINARY

To prove our results, we need the following lemmas.

Lemma 2.1 ([13]). *Let $\psi \in \mathcal{A}$ and $\mathcal{G}(z) = z^{-1} \int_0^z \psi(t) dt$, $z \in \mathbb{U}$. If*

$$\Re(1 + z\psi''(z)/\psi'(z)) > -\frac{1}{2}, \quad z \in \mathbb{U},$$

then $\mathcal{G} \in \mathcal{K}$.

Lemma 2.2 ([15], Theorem 2.6). *Assume that h is a convex function with $h(0) = a$ and $\nu \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$ with $\Re(\nu) \geq 0$. If $p \in \mathcal{H}[a, n]$ with $p(0) = a$, $\Phi: \mathbb{C}^2 \times \mathbb{U} \rightarrow \mathbb{C}$, $\Phi(p(z), zp'(z); z) = p(z) + \nu^{-1}zp'(z)$ is analytic function in \mathbb{U} and*

$$\mathcal{F}_{\Phi(\mathbb{C}^2 \times \mathbb{U})}\left(p(z) + \frac{1}{\nu}zp'(z)\right) \leq \mathcal{F}_{h(\mathbb{U})}h(z), \quad \text{i.e. } p(z) + \frac{1}{\nu}zp'(z) \prec_{\mathcal{F}} h(z), \quad z \in \mathbb{U},$$

then

$$\mathcal{F}_{p(\mathbb{U})}p(z) \leq \mathcal{F}_{q(\mathbb{U})}q(z) \leq \mathcal{F}_{h(\mathbb{U})}h(z), \quad \text{i.e. } p(z) \prec_{\mathcal{F}} q(z), \quad z \in \mathbb{U},$$

where

$$q(z) = \frac{\nu}{nz^{\nu/n}} \int_0^z h(t)t^{-1+\nu/n} dt, \quad z \in \mathbb{U}.$$

The function q is convex and it is the fuzzy best dominant.

Lemma 2.3 ([15], Theorem 2.7). *Let g be a convex function in \mathbb{U} and let $\psi(z) = g(z) + n\gamma zg'(z)$, where $z \in \mathbb{U}$, $n \in \mathbb{N}$ and $\gamma > 0$. If*

$$p(z) = g(0) + p_n z^n + p_{n+1} z^{n+1} + \dots$$

is holomorphic in \mathbb{U} and

$$\mathcal{F}_{p(\mathbb{U})}(p(z) + \gamma zp'(z)) \leq \mathcal{F}_{\psi(\mathbb{U})}\psi(z), \quad \text{i.e. } p(z) + \gamma zp'(z) \prec_{\mathcal{F}} \psi(z), \quad z \in \mathbb{U},$$

then

$$\mathcal{F}_{p(\mathbb{U})}(p(z)) \leq \mathcal{F}_{g(\mathbb{U})}g(z), \quad \text{i.e. } p(z) \prec_{\mathcal{F}} g(z), \quad z \in \mathbb{U};$$

this result is sharp.

For the general theory of fuzzy differential subordination and its applications, we refer the reader to [2], [12]–[14], [16].

The objective of the present section is to obtain several fuzzy differential subordinations associated with the integral operator $\mathcal{I}_{m,\gamma}^{n,\alpha}$ by using the method of fuzzy differential subordination.

3. MAIN RESULTS

Assume that $\eta \in [0, 1)$, $n \in \mathbb{N}_0$, $m > 0$, $\alpha, \gamma \geq 0$ and $z \in \mathbb{U}$ are mentioned through this paper:

Theorem 3.1. *Let k be a convex function in \mathbb{U} and suppose that $h(z) = k(z) + zk'(z)/(\lambda + 2)$. If $f \in \mathcal{M}_{m,\gamma}^F(n, \alpha, \eta)$ and*

$$(3.1) \quad G(z) = I^\lambda f(z) = \frac{\lambda + 2}{z^{\lambda+1}} \int_0^z t^\lambda f(t) dt,$$

then

$$(3.2) \quad F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \leq F_{h(\mathbb{U})}h(z), \quad \text{i.e. } (\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \prec_{\mathcal{F}} h(z),$$

implies that

$$F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} G)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \leq F_{k(\mathbb{U})}k(z), \quad \text{i.e. } (\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \prec_{\mathcal{F}} k(z);$$

this result is sharp.

Proof. Since

$$z^{\lambda+1}G(z) = (\lambda + 2) \int_0^z t^\lambda f(t) dt,$$

by differentiating, we obtain

$$(\lambda + 1)G(z) + zG'(z) = (\lambda + 2)f(z)$$

and

$$(3.3) \quad (\lambda + 1)\mathcal{I}_{m,\gamma}^{n,\alpha}G(z) + z(\mathcal{I}_{m,\gamma}^{n,\alpha}G(z))' = (\lambda + 2)\mathcal{I}_{m,\gamma}^{n,\alpha}f(z),$$

and also by differentiating (3.3) we obtain

$$(3.4) \quad (\mathcal{I}_{m,\gamma}^{n,\alpha}G(z))' + \frac{1}{(\lambda + 2)}z(\mathcal{I}_{m,\gamma}^{n,\alpha}G(z))'' = (\mathcal{I}_{m,\gamma}^{n,\alpha}f(z))'.$$

By using (3.4), the fuzzy differential subordination (3.2) becomes

$$(3.5) \quad F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}((\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' + \frac{1}{(\lambda + 2)}z(\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))'') \leq F_{h(\mathbb{U})}\left(k(z) + \frac{1}{(\lambda + 2)}zk'(z)\right).$$

We denote

$$(3.6) \quad q(z) = (\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \rightarrow q \in \mathcal{H}[1, n].$$

From (3.6) in (3.5) we have

$$(3.7) \quad F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})} \left(q(z) + \frac{1}{(\lambda+2)} zq'(z) \right) \leq F_{h(\mathbb{U})} \left(k(z) + \frac{1}{(\lambda+2)} zk'(z) \right),$$

by applying Lemma 2.3, we have

$$F_{q(\mathbb{U})} q(z) \leq F_{k(\mathbb{U})} k(z), \quad \text{i.e. } F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))'(\mathbb{U})} (\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \leq F_{k(\mathbb{U})} k(z),$$

then $(\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \prec_{\mathcal{F}} k(z)$, k is the fuzzy best dominant. \square

Theorem 3.2. Consider $h(z) = (1 + (2\eta - 1)z)/(1 + z)$, $\eta \in [0, 1)$, $\lambda > 0$ and I^λ is given by (3.1). Then

$$(3.8) \quad I^\lambda(\mathcal{M}_{m,\gamma}^F(n, \alpha, \eta)) \subset \mathcal{M}_{m,\gamma}^F(n, \alpha, \zeta),$$

where

$$(3.9) \quad \zeta = 2\eta - 1 + (\lambda + 2)(2 - 2\eta) \int_0^1 \frac{t^{\lambda+2}}{t+1} dt.$$

Proof. The function h is convex and using the same technique as in the proof of Theorem 3.1, we obtain from the hypothesis of Theorem 3.2 that

$$F_{q(\mathbb{U})} \left(q(z) + \frac{1}{(\lambda+2)} zq'(z) \right) \leq F_{h(\mathbb{U})} h(z),$$

where $q(z)$ is defined in (3.6). By using Lemma 2.2, we obtain that

$$F_{q(\mathbb{U})} q(z) \leq F_{k(\mathbb{U})} k(z) \leq F_{h(\mathbb{U})} h(z)$$

implies that

$$F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} G)'(\mathbb{U})} (\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \leq F_{k(\mathbb{U})} k(z) \leq F_{h(\mathbb{U})} h(z),$$

where

$$k(z) = \frac{\lambda+2}{z^{\lambda+2}} \int_0^z t^{\lambda+1} \frac{1+(2\eta-1)t}{1+t} dt = (2\eta-1) + \frac{(\lambda+2)(2-2\eta)}{z^{\lambda+2}} \int_0^z \frac{t^{\lambda+1}}{1+t} dt.$$

Since k is convex and $k(\mathbb{U})$ is symmetric with respect to the real axis, we conclude

$$(3.10) \quad F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} G)'(\mathbb{U})} (\mathcal{I}_{m,\gamma}^{n,\alpha} G(z))' \geq \min_{|z|=1} F_{k(\mathbb{U})} k(z) = F_{k(\mathbb{U})} k(1),$$

and $\zeta = k(1) = 2\eta - 1 + (\lambda + 2)(2 - 2\eta) \int_0^1 \frac{t^{\lambda+2}}{t+1} dt$. From (3.10), we conclude that we have the inclusion in relation (3.8) and hence the proof of the theorem is complete. \square

Theorem 3.3. Assume that k is a convex function in \mathbb{U} , $k(0) = 1$, and let $h(z) = k(z) + zk'(z)$. If $f \in \mathcal{A}$ and satisfies the fuzzy differential subordination

$$(3.11) \quad F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \leq F_{h(\mathbb{U})}h(z), \quad \text{i.e. } (\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \prec_{\mathcal{F}} h(z),$$

then

$$(3.12) \quad F_{\mathcal{I}_{m,\gamma}^{n,\alpha} f(\mathbb{U})} \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \leq F_{k(\mathbb{U})}k(z), \quad \text{i.e. } \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \prec_{\mathcal{F}} k(z),$$

and this result is sharp.

Proof. By denoting

$$\begin{aligned} q(z) &= \frac{1}{z} \mathcal{I}_{m,\gamma}^{n,\alpha} f(z) = \frac{1}{z} \left(z + \sum_{k=2}^{\infty} (1 + \gamma(k-1))^n \left(\frac{m}{m+k} \right)^{\alpha} a_k z^k \right) \\ &= 1 + \sum_{k=2}^{\infty} (1 + \gamma(k-1))^n \left(\frac{m}{m+k} \right)^{\alpha} a_k z^{k-1}, \end{aligned}$$

we obtain that $q(z) + zq'(z) = (\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))'$. Then

$$\begin{aligned} F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' &\leq F_{h(\mathbb{U})}h(z) \\ &\rightarrow F_{q(\mathbb{U})}(q(z) + zq'(z)) \leq F_{h(\mathbb{U})}h(z) = F_{k(\mathbb{U})}(k(z) + zk'(z)). \end{aligned}$$

Applying Lemma 2.3, we have

$$F_{q(\mathbb{U})}q(z) \leq F_{k(\mathbb{U})}k(z) \rightarrow F_{\mathcal{I}_{m,\gamma}^{n,\alpha} f(\mathbb{U})} \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \leq F_{k(\mathbb{U})}k(z),$$

we get

$$\frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \prec_{\mathcal{F}} k(z),$$

and this result is sharp. □

Theorem 3.4. For $h \in \mathcal{H}(\mathbb{U})$, $h(0) = 1$, which satisfies $\Re(1 + zh''(z)/h'(z)) > -\frac{1}{2}$, if $f \in \mathcal{A}$ and verifies the fuzzy differential subordination

$$(3.13) \quad F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \leq F_{h(\mathbb{U})}h(z), \quad \text{i.e. } (\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \prec_{\mathcal{F}} h(z),$$

then

$$(3.14) \quad F_{\mathcal{I}_{m,\gamma}^{n,\alpha} f(\mathbb{U})} \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \leq F_{k(\mathbb{U})}k(z), \quad \text{i.e. } \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \prec_{\mathcal{F}} k(z),$$

where

$$k(z) = \frac{1}{z} \int_0^z h(t) dt.$$

The function k is convex and it is the fuzzy best dominant.

Proof. Let

$$q(z) = \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} = 1 + \sum_{k=2}^{\infty} (1 + \gamma(k-1))^n \left(\frac{m}{m+k}\right)^\alpha a_k z^{k-1}, \quad q \in \mathcal{H}[1, 1],$$

where $\Re(1 + zh''(z)/h'(z)) > -\frac{1}{2}$. From Lemma 2.1, we have that

$$k(z) = \frac{1}{z} \int_0^z h(t) dt$$

is a convex function which satisfies the fuzzy differential subordination (3.13). Since

$$k(z) + zk'(z) = h(z)$$

is the fuzzy best dominant, we have $q(z) + zq'(z) = (\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))'$, then (3.13) becomes

$$F_{q(\mathbb{U})}(q(z) + zq'(z)) \leq F_{h(\mathbb{U})}h(z).$$

By applying Lemma 2.3, we have

$$F_{q(\mathbb{U})}q(z) \leq F_{k(\mathbb{U})}k(z) \rightarrow F_{\mathcal{I}_{m,\gamma}^{n,\alpha} f(\mathbb{U})} \frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \leq F_{k(\mathbb{U})}k(z),$$

and we obtain that

$$\frac{\mathcal{I}_{m,\gamma}^{n,\alpha} f(z)}{z} \prec_{\mathcal{F}} k(z).$$

□

Putting $h(z) = (1 + (2\beta - 1)z)/(1 + z)$ in Theorem 3.4, we obtain the following corollary:

Corollary 3.1. *Let $h = (1 + (2\beta - 1)z)/(1 + z)$ be a convex function in \mathbb{U} with $h(0) = 1$, $0 \leq \beta < 1$. If $f \in \mathcal{A}$ and verifies the fuzzy differential subordination*

$$F_{(\mathcal{I}_{m,\gamma}^{n,\alpha} f)'(\mathbb{U})}(\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \leq F_{h(\mathbb{U})}h(z), \quad \text{i.e. } (\mathcal{I}_{m,\gamma}^{n,\alpha} f(z))' \prec_{\mathcal{F}} h(z),$$

then

$$k(z) = 2\beta - 1 + \frac{2(1 - \beta)}{z} \ln(1 + z),$$

and the function k is convex and it is the fuzzy best dominant.

4. CONCLUSION

All the above results give us information about fuzzy differential subordinations for a linear operator $\mathcal{I}_{m,\gamma}^{n,\alpha}$. We give some properties for the class $\mathcal{M}_{m,\gamma}^F(n, \alpha, \eta)$ of univalent analytic functions.

References

- [1] *F. M. Al-Oboudi*: On univalent functions defined by a generalized Sălăgean operator. *Int. J. Math. Math. Sci.* *25-27* (2004), 1429–1436. [zbl](#) [MR](#) [doi](#)
- [2] *A. Alb Lupaş*: On special fuzzy differential subordinations using convolution product of Sălăgean operator and Ruscheweyh derivative. *J. Comput. Anal. Appl.* *15* (2013), 1484–1489. [zbl](#) [MR](#)
- [3] *A. Alb Lupaş, G. Oros*: On special fuzzy differential subordinations using Sălăgean and Ruscheweyh operators. *Appl. Math. Comput.* *261* (2015), 119–127. [zbl](#) [MR](#) [doi](#)
- [4] *M. K. Aouf*: Some inclusion relationships associated with the Komatu integral operator. *Math. Comput. Modelling* *50* (2009), 1360–1366. [zbl](#) [MR](#) [doi](#)
- [5] *M. K. Aouf*: The Komatu integral operator and strongly close-to-convex functions. *Bull. Math. Anal. Appl.* *3* (2011), 209–219. [zbl](#) [MR](#)
- [6] *S. D. Bernardi*: Convex and starlike univalent functions. *Trans. Am. Math. Soc.* *135* (1969), 429–446. [zbl](#) [MR](#) [doi](#)
- [7] *T. Bulboacă*: *Differential Subordinations and Superordinations: Recent Results*. House of Scientific Book Publ., Cluj-Napoca, 2005.
- [8] *A. Ebadian, S. Najafzadeh*: Uniformly starlike and convex univalent functions by using certain integral operators. *Acta Univ. Apulensis, Math. Inform.* *20* (2009), 17–23. [zbl](#) [MR](#)
- [9] *R. M. El-Ashwah, M. K. Aouf, S. M. El-Deeb*: Differential subordination for certain subclasses of p -valent functions associated with generalized linear operator. *J. Math.* *2013* (2013), Article ID 692045, 8 pages. [zbl](#) [MR](#) [doi](#)
- [10] *S. G. Gal, A. I. Ban*: *Elemente de Matematica Fuzzy*. University of Oradea, Oradea, 1996. (In Romanian.)
- [11] *S. M. Khairnar, M. More*: On a subclass of multivalent β -uniformly starlike and convex functions defined by a linear operator. *IAENG, Int. J. Appl. Math.* *39* (2009), 175–183. [zbl](#) [MR](#)
- [12] *Y. Komatu*: On analytic prolongation of a family of integral operators. *Math., Rev. Anal. Numér. Théor. Approximation, Math.* *32(55)* (1990), 141–145. [zbl](#) [MR](#)
- [13] *S. S. Miller, P. T. Mocanu*: *Differential Subordination: Theory and Applications*. Pure and Applied Mathematics, Marcel Dekker 225. Marcel Dekker, New York, 2000. [zbl](#) [MR](#) [doi](#)
- [14] *G. I. Oros, G. Oros*: The notation of subordination in fuzzy sets theory. *Gen. Math.* *19* (2011), 97–103. [zbl](#) [MR](#)
- [15] *G. I. Oros, G. Oros*: Dominants and best dominants in fuzzy differential subordinations. *Stud. Univ. Babeş-Bolyai, Math.* *57* (2012), 239–248. [zbl](#) [MR](#)
- [16] *G. I. Oros, G. Oros*: Fuzzy differential subordination. *Acta Univ. Apulensis, Math. Inform.* *30* (2012), 55–64. [zbl](#) [MR](#)
- [17] *R. K. Raina, I. B. Bapna*: On the starlikeness and convexity of a certain integral operator. *Southeast Asian Bull. Math.* *33* (2009), 101–108. [zbl](#) [MR](#)
- [18] *G. S. Sălăgean*: Subclasses of univalent functions. *Complex Analysis – Fifth Romanian-Finnish Seminar. Lecture Notes in Mathematics 1013*. Springer, Berlin, 1983, pp. 362–372. [zbl](#) [MR](#) [doi](#)

Authors' addresses: *Sheza Mohammed El-Deeb*, Department of Mathematics, Faculty of Science, Damietta University, New Damietta 34517, Egypt and Department of Mathematics, College of Science and Arts in Badaya, Qassim University, Buraidah 51452, Saudi Arabia, e-mail: shezaeldeeb@yahoo.com; *Georgia Irina Oros*, Department of Mathematics, University of Oradea, Str. Universitatii, No. 1, 410087 Oradea, Romania, e-mail: georgia_oros_ro@yahoo.co.uk.