

Genaro Hernandez-Mada; Humberto Abraham Martinez-Gil

Orbit-cone correspondence for the proalgebraic completion of normal toric varieties

Archivum Mathematicum, Vol. 61 (2025), No. 1, 1–8

Persistent URL: <http://dml.cz/dmlcz/152914>

Terms of use:

© Masaryk University, 2025

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ORBIT-CONE CORRESPONDENCE FOR THE PROALGEBRAIC COMPLETION OF NORMAL TORIC VARIETIES

GENARO HERNANDEZ-MADA AND HUMBERTO ABRAHAM MARTINEZ-GIL

ABSTRACT. We prove that there is an orbit-cone correspondence for the proalgebraic completion of normal toric varieties, which is analogous to the classical orbit-cone correspondence for toric varieties.

1. INTRODUCTION

Let N be a finite rank free abelian group (also called a lattice) and $\mathcal{F}$ a fan in $N_{\mathbb{R}} = N \otimes_{\mathbb{Z}} \mathbb{R}$. It is well-known that there is a toric variety $X^{\mathcal{F}}$ associated to $\mathcal{F}$. Moreover a classical result states that if T_N is the torus of $X^{\mathcal{F}}$ then there is a bijective correspondence between the T_N -orbits in $X^{\mathcal{F}}$ and the cones σ in $\mathcal{F}$. (See for example [2], Theorem 3.2.6).

Now consider $(I, \leq)$ a directed poset and a projective system of complex algebraic varieties

$$f_{ij}: V_j \rightarrow V_i, \quad i, j \in I, \quad i \leq j.$$

This system is called **proalgebraic variety**, and as pointed out in [1] this concept plays an important role in dynamical systems. In the cases that we will study, we can take projective limit and obtain **laminations**, (also called **solenoidal manifolds** in [6]). These have been used, for example in [4] and [6], to obtain results in conformal dynamics.

In this paper we consider a projective system of toric varieties. Namely, given a normal toric variety X , for each $m, n \in \mathbb{N}$ such that $n \mid m$, we can define

$$p_{n,m}: \mathbb{C} \rightarrow \mathbb{C}$$

by $z \mapsto z^{m/n}$, and these induce

$$q_{n,m}: X \rightarrow X,$$

which define a projective system. The projective limit, denoted by $X_{\mathbb{Q}}$ is called the **proalgebraic completion** of X . The first part of [1] is devoted to the study of the proalgebraic completion of toric varieties. More particularly, in Theorem 1 they

2020 *Mathematics Subject Classification*: primary 14M25; secondary 13B35, 57R30, 11R56.

Key words and phrases: toric varieties, orbit-cone correspondence, proalgebraic completion, algebraic solenoid.

Received September 24, 2024. Editor J. Slovák.

DOI: 10.5817/AM2025-1-1

give a description of its vector bundle category. For a more general treatment of proalgebraic completions, we refer to [3].

The main result of this paper is an analogous result to the one described in the first paragraph. Namely, we will prove that there is a *proalgebraic torus* acting on the proalgebraic completion of the toric variety $X^{\mathcal{F}}$ (defined by the fan $\mathcal{F}$), and there is a bijective correspondence of its orbits with the cones of $\mathcal{F}$. This correspondence satisfies similar properties to the classical one. See Theorem 3 for the precise statement and Theorem 1 for the comparison with the classical result.

Now let us give an outline of this paper. In Section 2 we recall some concepts regarding toric varieties, as well as the precise statement of the classical theorem that inspires this work (Theorem 1). In section 3 we give the basic definitions regarding proalgebraic completions of normal toric varieties. Namely, we define the algebraic solenoid, which allows to define the proalgebraic torus and finally the proalgebraic completion of a normal toric variety. For this we define it first for an affine toric variety and we use this for the general case. Finally in Section 4 we prove our main result.

2. PRELIMINARIES ON TORIC VARIETIES

Given a lattice N with dual $M = \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$ and a rational strongly convex polyhedral cone $\sigma \subset N \otimes_{\mathbb{Z}} \mathbb{R}$, we have the classical definition as in [5] of the affine normal toric variety

$$X^{\sigma} = \text{Hom}_{sg}(S^{\sigma}, \mathbb{C}),$$

where $S^{\sigma} \subset M$ is the semigroup associated to σ . In particular, each point of X^{σ} is a homomorphism of semigroups $S^{\sigma} \rightarrow \mathbb{C}$. In this setting, the torus of X^{σ} is the topological group

$$T_N = \text{Hom}_{\mathbb{Z}}(M, \mathbb{C}^*).$$

If $\mathcal{F}$ is a fan that lies in the vector space $N \otimes_{\mathbb{Z}} \mathbb{R}$, then for every $\sigma_1, \sigma_2 \in \mathcal{F}$ and $\tau = \sigma_1 \cap \sigma_2$ there are natural injective morphisms

$$\text{Hom}_{sg}(S^{\sigma_1}, \mathbb{C}) \hookleftarrow \text{Hom}_{sg}(S^{\tau}, \mathbb{C}) \hookrightarrow \text{Hom}_{sg}(S^{\sigma_2}, \mathbb{C}).$$

These define an equivalence relation so the affine toric varieties X^{σ} , $\sigma \in \mathcal{F}$ glue together to get the normal toric variety

$$X^{\mathcal{F}} = \bigsqcup_{\sigma \in \mathcal{F}} \text{Hom}_{sg}(S^{\sigma}, \mathbb{C}) / \sim,$$

with torus T_N . Recall that the action of T_N on $X^{\mathcal{F}}$ is given by the map

$$(1) \quad T_N \times X^{\mathcal{F}} \rightarrow X^{\mathcal{F}}, \quad \tilde{\gamma} \cdot [\gamma] = [\tilde{\gamma}|_{S^{\sigma}} \gamma],$$

where $\gamma: S^{\sigma} \rightarrow \mathbb{C}$, $\tilde{\gamma}: M \rightarrow \mathbb{C}^*$.

An important result in toric geometry is that the T_N -orbits of a toric variety

$X^{\mathcal{F}}$ are characterized by the **special points**. Namely, by the semigroup homomorphism (point in $X^{\sigma} = \text{Hom}_{sg}(S^{\sigma}, \mathbb{C})$) defined for every $\sigma \in \mathcal{F}$ by

$$m \in S^{\sigma} \mapsto \begin{cases} 1 & \text{if } m \in S^{\sigma} \cap \sigma^{\perp} = \sigma^{\perp} \cap M, \\ 0 & \text{otherwise,} \end{cases}$$

where $\sigma^{\perp}$ is the orthogonal complement of σ on the dual vector space of $N \otimes_{\mathbb{Z}} \mathbb{R}$.

Theorem 1. *Let $X^{\mathcal{F}}$ be the toric variety associated to the fan $\mathcal{F}$ on $N \otimes_{\mathbb{Z}} \mathbb{R}$. Then*

(a) *There is a bijective correspondence*

$$\begin{aligned} \{\text{Cones } \sigma \in \mathcal{F}\} &\longleftrightarrow \{T_N\text{-orbits on } X^{\mathcal{F}}\} \\ \sigma &\longleftrightarrow T_N \cdot \gamma^{\sigma}. \end{aligned}$$

(b) *If $\dim N \otimes_{\mathbb{Z}} \mathbb{R} = r$, then for every cone $\sigma \in \mathcal{F}$ the orbit $T_N \cdot \gamma^{\sigma}$ is an algebraic torus of dimension $r - \dim \sigma$.*

(c) *For every $\sigma \in \mathcal{F}$ the affine toric variety X^{σ} is the union of orbits*

$$X^{\sigma} = \bigcup_{\tau \text{ face of } \sigma} T_N \cdot \gamma^{\tau}.$$

The proof is in [2] Theorem 3.2.6. Our main result is a proalgebraic version of this theorem. To get there, we will use the notion of proalgebraic completion of a toric variety from [1].

3. PROALGEBRAIC COMPLETION OF NORMAL TORIC VARIETIES

For every $n, m \in \mathbb{N}$ such that $n \mid m$ (n divides m) there is a finite covering map

$$(2) \quad p_{n,m}: \mathbb{C}^* \rightarrow \mathbb{C}^*, \quad z \mapsto z^{m/n}.$$

This determines a projective system $\{\mathbb{C}^*, p_{n,m}\}_{n|m}$ of covering spaces whose projective limit is the **algebraic solenoid** $\mathbb{C}_{\mathbb{Q}}^*$ which is a topological group with the initial topology defined by the canonical projections

$$\pi_n: \mathbb{C}_{\mathbb{Q}}^* \rightarrow \mathbb{C}^*, \quad (z_j)_{j \in \mathbb{N}} \mapsto z_n.$$

Let $\widehat{\mathbb{Z}}$ be the profinite completion of the integers. Then the algebraic solenoid is a 2-dimensional **lamination** in the sense of [4] with **transversal space** $\widehat{\mathbb{Z}}$. Furthermore, the **global leaves** of the lamination are just copies of $\mathbb{C}$, and more precisely all the global leaves are obtained by translations of the image of the injective and continuous map

$$\nu: \mathbb{C} \rightarrow \mathbb{C}_{\mathbb{Q}}^*, \quad z \mapsto (e^{i(z/j)})_{j \in \mathbb{N}}$$

by elements of $\widehat{\mathbb{Z}}$. This algebraic solenoid is thoroughly studied in [1].

Definition 1. Let N be a lattice with dual $M = \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$. The group

$$(T_N)_{\mathbb{Q}} = \text{Hom}_{\mathbb{Z}}(M, \mathbb{C}_{\mathbb{Q}}^*)$$

is called **proalgebraic torus**.

Note that if $\text{rank } M = r$, then we have a natural group isomorphisms

$$N \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^* \cong (T_N)_{\mathbb{Q}} \cong (\mathbb{C}_{\mathbb{Q}}^*)^r.$$

Hence the proalgebraic torus $(T_N)_{\mathbb{Q}}$ is also a lamination ($2r$ -dimensional) with transversal space $\widehat{\mathbb{Z}}^r$. In this case, the global leaves are copies of $\mathbb{C}^r$, and all of them are obtained via translations by elements of $\widehat{\mathbb{Z}}^r$. The maps in 2 used to define the algebraic solenoid extend naturally to the whole complex plane $\mathbb{C}$, so we can consider the limit

$$\mathbb{C}_{\mathbb{Q}} = \{(z_j)_{j \in \mathbb{N}} \mid z_j \in \mathbb{C} \text{ for all } j \in \mathbb{N}, p_{n,m}(z_m) = z_n \text{ whenever } n \mid m\}.$$

Definition 2. Let σ be a rational strongly convex polyhedral cone with associated semigroup S^σ . The set

$$X_{\mathbb{Q}}^\sigma = \text{Hom}_{sg}(S^\sigma, \mathbb{C}_{\mathbb{Q}})$$

is **the proalgebraic completion** of the affine toric variety $X^\sigma = \text{Hom}_{sg}(S^\sigma, \mathbb{C})$.

It is clear that an element γ in $X_{\mathbb{Q}}^\sigma$ induces a collection $\{\gamma_j: S^\sigma \rightarrow \mathbb{C}\}_{j \in \mathbb{N}}$ of semigroup homomorphisms satisfying $(\gamma_m(s))^{m/n} = \gamma_n(s)$ for all $s \in S^\sigma$ whenever $n \mid m$, and there is a bijective correspondence

$$X_{\mathbb{Q}}^\sigma \rightarrow \varprojlim_{q_{n,m}} \text{Hom}_{sg}(S^\sigma, \mathbb{C}), \quad \gamma \mapsto (\gamma_j)_{j \in \mathbb{N}},$$

where

$$q_{n,m}: \text{Hom}_{\mathbb{Z}}(S^\sigma, \mathbb{C}) \rightarrow \text{Hom}_{\mathbb{Z}}(S^\sigma, \mathbb{C}), \quad \gamma \mapsto p_{n,m} \circ \gamma$$

is an algebraic map. Under this correspondence we can endow the proalgebraic completion $X_{\mathbb{Q}}^\sigma$ with a topology.

More generally, let $\mathcal{F}$ be a fan and consider the collection of proalgebraic completions

$$\{X_{\mathbb{Q}}^\sigma\}_{\sigma \in \mathcal{F}} = \{\text{Hom}_{sg}(S^\sigma, \mathbb{C}_{\mathbb{Q}})\}_{\sigma \in \mathcal{F}}.$$

As in the classical case, if $\sigma_1, \sigma_2 \in \mathcal{F}$, and τ is a face of $\sigma_1 \cap \sigma_2$, we have natural injective maps

$$(3) \quad X_{\mathbb{Q}}^{\sigma_1} = \text{Hom}_{sg}(S^{\sigma_1}, \mathbb{C}_{\mathbb{Q}}) \hookleftarrow \text{Hom}_{sg}(S^\tau, \mathbb{C}_{\mathbb{Q}}) \hookrightarrow \text{Hom}_{sg}(S^{\sigma_2}, \mathbb{C}_{\mathbb{Q}}) = X_{\mathbb{Q}}^{\sigma_2}.$$

These maps allow to define an equivalence relation on

$$\bigsqcup_{\sigma \in \mathcal{F}} \text{Hom}_{sg}(S^\sigma, \mathbb{C}_{\mathbb{Q}})$$

as follows: $(\gamma_1: S^{\sigma_1} \rightarrow \mathbb{C}_{\mathbb{Q}}) \sim (\gamma_2: S^{\sigma_2} \rightarrow \mathbb{C}_{\mathbb{Q}})$ if and only if there exists a face τ of $\sigma_1 \cap \sigma_2$ and a homomorphism $\gamma: S^\tau \rightarrow \mathbb{C}_{\mathbb{Q}}$ such that $\gamma|_{S^{\sigma_1}} = \gamma_1, \gamma|_{S^{\sigma_2}} = \gamma_2$.

Definition 3. Let N be a lattice and let $\mathcal{F}$ be a fan in $N \otimes_{\mathbb{Z}} \mathbb{R}$. We define **the proalgebraic completion** $X_{\mathbb{Q}}^{\mathcal{F}}$ of the toric variety $X^{\mathcal{F}}$ as the quotient space

$$X_{\mathbb{Q}}^{\mathcal{F}} := \bigsqcup_{\sigma \in \mathcal{F}} \text{Hom}_{sg}(S^\sigma, \mathbb{C}_{\mathbb{Q}}) / \sim.$$

As in the affine case, the proalgebraic completion $X_{\mathbb{Q}}^{\mathcal{F}}$ turns out to be a projective limit. More precisely we have the following (see [1] Section 4):

Theorem 2. *The proalgebraic completion $X_{\mathbb{Q}}^{\mathcal{F}}$ of a toric variety $X^{\mathcal{F}}$ is homomorphic to the projective limit over all its toric covers.*

With the previous result it is clear that there is a canonical inclusion

$$(T_N)_{\mathbb{Q}} \subset X_{\mathbb{Q}}^{\mathcal{F}}$$

induced by the inclusion $T_N \subset X^{\mathcal{F}}$. Moreover, the action described in (1) induces the action

$$(4) \quad (T_N)_{\mathbb{Q}} \times X_{\mathbb{Q}}^{\mathcal{F}} \rightarrow X_{\mathbb{Q}}^{\mathcal{F}}, \quad \tilde{\gamma} \cdot [\gamma] = [\tilde{\gamma}|_{S^{\sigma}} \cdot \gamma],$$

where $\gamma: S^{\sigma} \rightarrow \mathbb{C}_{\mathbb{Q}}$, $\tilde{\gamma}: M \rightarrow \mathbb{C}_{\mathbb{Q}}^*$.

Remark 1. If X is a toric variety defined by a fan $\mathcal{F}$ in $N \otimes_{\mathbb{Z}} \mathbb{R}$, for every $m \in \mathbb{N}$ we take $X_m = X$, and then for every $n \mid m$ we have a map $N \rightarrow N$ which is multiplication by the integer $\frac{m}{n}$. Since this map takes cones of $\mathcal{F}$ into cones of $\mathcal{F}$, it defines a map of toric varieties $X_m \rightarrow X_n$, and thus a projective system where all the maps involved are algebraic. Moreover, these maps are affine, so we get a scheme structure on $X_{\mathbb{Q}}$.

Remark 2. Note that if X is singular, it is not a manifold, and perhaps a more suitable name for the proalgebraic completion would be solenoidal variety, instead of solenoidal manifold.

4. THE ORBIT-CONE CORRESPONDENCE

In order to study the $(T_N)_{\mathbb{Q}}$ -orbits of the proalgebraic completion $X_{\mathbb{Q}}^{\mathcal{F}}$ we have the following definition which is the proalgebraic version of the special points in $X^{\mathcal{F}}$.

Definition 4. Let σ be a rational strongly convex polyhedral cone in $N \otimes_{\mathbb{Z}} \mathbb{R}$. We define the **special point** (or distinguished point) in $X_{\mathbb{Q}}^{\sigma} = \text{Hom}_{sg}(S^{\sigma}, \mathbb{C}_{\mathbb{Q}}^*)$ as the semigroup homomorphism

$$\gamma_{\mathbb{Q}}^{\sigma}(m) = \begin{cases} (1_j)_j & \text{if } m \in S^{\sigma} \cap \sigma^{\perp} = \sigma^{\perp} \cap M, \\ (0_j)_j & \text{otherwise,} \end{cases}$$

for $m \in S^{\sigma}$.

As we shall see in the following Lemma, the $(T_N)_{\mathbb{Q}}$ -orbits in $X_{\mathbb{Q}}^{\mathcal{F}}$ corresponding to the special points are proalgebraic torus themselves.

Lemma 1. *Let σ be a rational strongly convex polyhedral cone in $N \otimes_{\mathbb{Z}} \mathbb{R}$ and consider the lattice $N(\sigma) = N/N_{\sigma}$, where N_{σ} is the sublattice generated by the elements in $\sigma \cap N$. Then there is a bijection*

$$(T_N)_{\mathbb{Q}} \cdot \gamma_{\mathbb{Q}}^{\sigma} \simeq (T_{N(\sigma)})_{\mathbb{Q}} = \text{Hom}_{\mathbb{Z}}(\sigma^{\perp} \cap M, \mathbb{C}_{\mathbb{Q}}^*).$$

Proof. We define the map

$$(5) \quad (T_N)_{\mathbb{Q}} \cdot \gamma_{\mathbb{Q}}^{\sigma} \rightarrow (T_{N(\sigma)})_{\mathbb{Q}}, \quad \gamma \cdot \gamma_{\mathbb{Q}}^{\sigma} \mapsto (\gamma \cdot \gamma_{\mathbb{Q}}^{\sigma})|_{\sigma^{\perp} \cap M},$$

which is clearly injective by definition of $\gamma_{\mathbb{Q}}^{\sigma}$. To see that this correspondence is surjective, consider the short exact sequence

$$0 \rightarrow N_{\sigma} \rightarrow N \rightarrow N(\sigma) \rightarrow 0.$$

If we tensor with $\mathbb{C}_{\mathbb{Q}}^*$ then we obtain a surjective homomorphism

$$N \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^* \rightarrow N(\sigma) \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^* \rightarrow 0,$$

and hence a transitive action of $N \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^*$ on $N(\sigma) \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^*$.

Under the natural isomorphisms

$$N \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^* \cong \text{Hom}_{\mathbb{Z}}(M, \mathbb{C}_{\mathbb{Q}}^*) = (T_N)_{\mathbb{Q}}$$

and

$$N(\sigma) \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^* \cong \text{Hom}_{\mathbb{Z}}(\sigma^{\perp} \cap M, \mathbb{C}_{\mathbb{Q}}^*) = (T_{N(\sigma)})_{\mathbb{Q}},$$

the action of $N \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^*$ on $N(\sigma) \otimes_{\mathbb{Z}} \mathbb{C}_{\mathbb{Q}}^*$ corresponds with the action

$$(T_N)_{\mathbb{Q}} \times (T_{N(\sigma)})_{\mathbb{Q}} \rightarrow (T_{N(\sigma)})_{\mathbb{Q}}$$

given by $\gamma \cdot \alpha = \gamma|_{\sigma^{\perp} \cap M} \alpha$.

Note that by restricting the special point $\gamma_{\mathbb{Q}}^{\sigma}$ to the group $\sigma^{\perp} \cap M$ we get an element in $(T_{N(\sigma)})_{\mathbb{Q}}$. Since the previous action is transitive, the map (5) is surjective. $\square$

Remark 3. The group $\text{Hom}_{\mathbb{Z}}(\sigma^{\perp} \cap M, \mathbb{C}_{\mathbb{Q}}^*)$ from the previous lemma can be described as the following set of semigroup homomorphisms:

$$\{\gamma: S^{\sigma} \rightarrow \mathbb{C}_{\mathbb{Q}} \mid \gamma(m) \neq (0_j)_j \Leftrightarrow m \in \sigma^{\perp} \cap M\}.$$

Indeed, any element of this set can be restricted to obtain an element in

$$\text{Hom}_{\mathbb{Z}}(\sigma^{\perp} \cap M, \mathbb{C}_{\mathbb{Q}}^*).$$

Conversely, extending by zero, any element on the group $\text{Hom}_{\mathbb{Z}}(\sigma^{\perp} \cap M, \mathbb{C}_{\mathbb{Q}}^*)$ can be seen as an element in the set above.

Theorem 3 (The Orbit-Cone correspondence). *Let $\mathcal{F}$ be a fan in $N \otimes_{\mathbb{Z}} \mathbb{R}$ and let $X_{\mathbb{Q}}^{\mathcal{F}}$ be the proalgebraic completion of the normal toric variety $X^{\mathcal{F}}$. Then*

(a) *There is a bijective correspondence*

$$\begin{aligned} \{\text{Cones } \sigma \in \mathcal{F}\} &\longleftrightarrow \{(T_N)_{\mathbb{Q}}\text{-orbits in } X_{\mathbb{Q}}^{\mathcal{F}}\} \\ \sigma &\longleftrightarrow (T_N)_{\mathbb{Q}} \cdot \gamma_{\mathbb{Q}}^{\sigma}. \end{aligned}$$

(b) *If $\dim N \otimes_{\mathbb{Z}} \mathbb{R} = r$, then for every cone $\sigma \in \mathcal{F}$, the orbit $(T_N)_{\mathbb{Q}} \cdot \gamma_{\mathbb{Q}}^{\sigma}$ is a lamination of dimension $2(r - \dim \sigma)$.*

(c) *For every $\sigma \in \mathcal{F}$ the completion $X_{\mathbb{Q}}^{\sigma}$ is the union of orbits*

$$X_{\mathbb{Q}}^{\sigma} = \bigcup_{\tau \text{ face of } \sigma} (T_N)_{\mathbb{Q}} \cdot \gamma_{\mathbb{Q}}^{\tau}.$$

Proof. For surjectivity in (a), let O be a $(T_N)_\mathbb{Q}$ -orbit of $X_\mathbb{Q}^\mathcal{F}$. Since

$$X_\mathbb{Q}^\mathcal{F} = \bigcup_{\sigma \in \mathcal{F}} X_\mathbb{Q}^\sigma,$$

if $\sigma_1, \sigma_2 \in \mathcal{F}$, the elements of $X_\mathbb{Q}^{\sigma_1}$ that are related to the elements of $X_\mathbb{Q}^{\sigma_2}$ are precisely the elements of $X_\mathbb{Q}^{\sigma_1 \cap \sigma_2}$, i.e.,

$$X_\mathbb{Q}^{\sigma_1} \cap X_\mathbb{Q}^{\sigma_2} = X_\mathbb{Q}^{\sigma_1 \cap \sigma_2}.$$

Moreover each $X_\mathbb{Q}^\sigma$ is invariant under the action (4) of the torus $(T_N)_\mathbb{Q}$. Hence there exists a unique minimal cone $\sigma \in \mathcal{F}$ such that $O \subseteq X_\mathbb{Q}^\sigma$. We shall prove that $O = (T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^\sigma$. Indeed, let $\gamma \in O$ and consider the set

$$S = \{m \in S^\sigma \mid \gamma(m) \neq (0_j)_j\}.$$

For every $j \in \mathbb{N}$, let $\pi_j: \mathbb{C}_\mathbb{Q} \rightarrow \mathbb{C}$ be the canonical projection which is in particular a semigroup homomorphism, and note that

$$S = \{m \in S^\sigma \mid \gamma(m) \neq (0_j)_j\} = \{m \in S^\sigma \mid (\pi_j \circ \gamma)(m) \neq 0\}$$

for every $j \in \mathbb{N}$. It follows that $S = \Gamma \cap M$ for some face Γ of $\sigma^\vee$, and consequently there exists a face τ of σ such that $\Gamma = \sigma^\vee \cap \tau^\perp$.

Now the equality

$$S = \Gamma \cap M = \sigma^\vee \cap \tau^\perp \cap M$$

implies $\gamma \in X_\mathbb{Q}^\tau$, and then $\tau = \sigma$ because of the minimality of σ . By Remark 3,

$$S = \sigma^\perp \cap M$$

implies that $\gamma \in (T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^\sigma$ and therefore $O = (T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^\sigma$. This proves surjectivity.

For injectivity, let $\sigma_1 \neq \sigma_2$ be cones in $\mathcal{F}$. If $\dim \sigma_1 \neq \dim \sigma_2$, it follows from Lemma 1 that the orbits are not equal. If $\dim \sigma_1 = \dim \sigma_2$, an easy computation shows that

$$(T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^{\sigma_1}, (T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^{\sigma_2} \not\subseteq X_\mathbb{Q}^{\sigma_1 \cap \sigma_2},$$

and hence $(T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^{\sigma_1} \neq (T_N)_\mathbb{Q} \cdot \gamma_\mathbb{Q}^{\sigma_2}$.

Part (b) follows from Lemma 1 and the fact that any proalgebraic torus is a lamination; part (c) follows directly from part (a). $\square$

REFERENCES

- [1] Burgos, J.M., Verjovsky, A., *Adelic Toric Varieties and Adelic Loop Groups*, [arXiv:2001.07997v2](#). Preprint 2021.
- [2] Cox, D., Little, J., Schenck, H., *Toric Varieties*, Grad. Stud. Math., vol. 124, AMS, 2011.
- [3] Grothendieck, A., Raynaud, M., *Revêtement Étales et Groupe Fondamental (SGA1)*, Lecture Notes in Math., Springer, Berlin Heidelberg New York, 1971.
- [4] Lyubich, M., Minsky, Y., *Laminations in holomorphic dynamics*, J. Differential Geom. **47** (1997), 17–94.
- [5] Oda, T., *Convex bodies and algebraic geometry: An introduction to the theory of toric varieties*, Springer, 1988.
- [6] Sullivan, D., *Solenoidal manifolds*, J. Singul. **9** (2014), 203–205.

DEPARTAMENTO DE MATEMÁTICAS,
UNIVERSIDAD DE SONORA,
BLVD. LUIS ENCINAS S/N, HERMOSILLO,
MEXICO, C.P. 83000
E-mail: `genaro.hernandez@unison.mx` `humberto.martinezguni@gmail.com`