

Jürgen Flachsmeyer

On Grothendieck spaces of type $C(K)$

In: Zdeněk Frolík (ed.): Abstracta. 5th Winter School on Abstract Analysis.
Czechoslovak Academy of Sciences, Praha, 1977. pp. 17–18.

Persistent URL: <http://dml.cz/dmlcz/701078>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic,
1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project
DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

FIFTH WINTER SCHOOL (1977)

ON GROTHENDIECK SPACES OF TYPE C(K)

By

Jürgen FLACHSMEYER

In his fundamental work on weakly compact operators Grothendieck presented the following theorem (see Canad. J. Math. 5 (1953)):

For a Banach space E the following properties are equivalent:

- (1) Every continuous linear map $u: E \rightarrow Y$ from E into some separable Banach space is weakly compact, i.e. u transmits the unit ball into some relative weakly compact set.
- (2) Weak- \star -convergence and weak convergence for sequences coincide in the dual E' .

Banach spaces with these properties (1) and (2) are called Grothendieck spaces (see for example J. Diestel: Grothendieck spaces and vector measures. In: Vector and operator valued measures and applications. Ed. by D.H. Tucker, H.B. Maynard, Acad. Press. Inc. 1973, 97-108.)

The following problem (problem 3 in Diestel's paper is) unsolved: Characterize those compact Hausdorff spaces K for which the Banach space $C(K)$ of all continuous real-valued functions on K is a Grothendieck space.

What is known about this problem?

Let K be a compact Hausdorff space. We will write $K \in G$ iff $C(K)$ is Grothendieck.

Grothendieck (1953): (i) K Stonian (=extremally disconnected) \Rightarrow

Ando (1961): (ii) K \mathcal{C} -extremally disconnected $\Rightarrow K \in G$.

Semadeni (1964): by another approach received (ii).

Seeever (1968): (iii) K an F -space $\Rightarrow K \in G$.

H. Schaefer (1971) also proved (ii). Of course, (iii) \Rightarrow (ii) \Rightarrow (i).

By the Riesz representation theorem the dual $C'(K)$ can be identified with the space $M(K)$ of bounded signed Radon measures on K .

Using this approach $K \in G$ gets equivalent to the following.

For every sequence (μ_n) of bounded Radon measures holds:

$$\mu_n(f) \rightarrow 0 \quad \forall f \in C(K) \Rightarrow \mu_n(g) \rightarrow 0 \quad \forall \text{ bounded Borel functions } g.$$

Thus, a necessary condition for $K \in G$ is that K must be sequentially discrete.

The lecture now explains the following result:

For every infinite compact F -space K the Alexandrov-double $K \hat{\otimes} K$ is never an F -space.

For every $K \in G$ the Alexandrov-double $K \hat{\otimes} K$ belongs to G . Thus a good deal of non- F -spaces are in G .

(Remark: The extension of the class G in a suitable way to non-compact spaces was treated in a thesis (Greifswald 1976) by Nguyen Doan Tien).