

Ryszard Grzaślewicz

A universal convex set in Euclidean space

In: Zdeněk Frolík (ed.): Abstracta. 8th Winter School on Abstract Analysis.
Czechoslovak Academy of Sciences, Praha, 1980. pp. 90–93.

Persistent URL: <http://dml.cz/dmlcz/701184>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic,
1980

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project
DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

A universal convex set in Euclidean space

Ryszard Grząślewicz

Professor C. Ryll-Nardzewski has raised the question whether there exists a compact convex set Q in \mathbb{R}^3 such that every compact convex set with non-empty interior in \mathbb{R}^2 is affine isomorphic to some intersection of Q with a plane.

In this note we present an example of a compact convex set Q in \mathbb{R}^{n+2} ($n \geq 1$) such that every closed convex subset of the unit ball B of \mathbb{R}^n is an intersection of Q with some k -dimensional affine subspace of \mathbb{R}^{n+2} .

Let 2^B denote the space of all closed non-empty subsets of B endowed with the Hausdorff distance

$$\text{dist}(A_1, A_2) = \max \left(\sup_{x \in A_1} d(x, A_2), \sup_{y \in A_2} d(y, A_1) \right)$$

where d stands for the Euclidean metric $d(x, y) = \|x - y\| = \sqrt{\langle x - y, x - y \rangle}$ in \mathbb{R}^n . It is well known that 2^B is compact. It is also easy to see that if $\text{dist}(A_n, A_0) \rightarrow 0$ and $d(x_n, x_0) \rightarrow 0$ as $n \rightarrow \infty$ with $x_n \in A_n \in 2^B$, then $x_0 \in A_0$.

Lemma. The set \mathcal{C} of all convex sets in 2^B is a locally arcwise connected metric continuum.

Proof. Let a sequence A_n of elements in \mathcal{C} converge to $A_0 \in 2^B$ and suppose $x \in A_0$. Then clearly there exists a sequence (x_k) with $x_k \in A_k$ converging to x . This implies that if $x, y \in A_0$ then $\lambda x + (1 - \lambda)y \in A_0$ for every $0 \leq \lambda \leq 1$, so A_0 is convex. Thus \mathcal{C} is a closed subset of 2^B , so compact.

Now we prove that \mathcal{C} is locally arcwise connected. It is sufficient to show that for every different $A_0, A_1 \in \mathcal{C}$ there

exists an arc A_0A_1 with diameter $\leq \text{dist}(A_0, A_1)$ (see [1], p. 242). We denote $A_t = tA_1 + (1-t)A_0 = \{ty + (1-t)x : x \in A_0, y \in A_1\} \in \mathcal{C}$.

Let $x \in A_0, y \in A_1$ and let $x_0 \in A_0, y_0 \in A_1$ be such that $d(x, y_0) \leq \text{dist}(A_0, A_1)$ and $d(y, x_0) \leq \text{dist}(A_0, A_1)$. For $0 \leq t < s \leq 1$ we have

$$\begin{aligned} d(sy + (1-s)x, A_t) &\leq d(sy + (1-s)x, ty + (1-t) \left[\frac{1-s}{1-t} x + \frac{s-t}{1-t} x_0 \right]) = \\ &= \|(s-t)(y-x_0)\| \leq |s-t| \text{dist}(A_0, A_1) \text{ and, analogously,} \\ d(ty + (1-t)x, A_s) &\leq |s-t| \text{dist}(A_0, A_1). \end{aligned}$$

Thus for $t, s \in [0, 1]$ we have

$$\text{dist}(A_t, A_s) \leq |s-t| \text{dist}(A_0, A_1).$$

Let $x_1, x_2 \in A_0$ and $y_1, y_2 \in A_1$ be such that

$$\begin{aligned} \sup_{y \in A_1} d(y, A_0) &= d(y_1, A_0) = d(y_1, x_1) \text{ and } \sup_{x \in A_0} d(x, A_1) = d(x_2, A_1) = \\ &= d(x_2, y_2). \end{aligned}$$

$$\text{Then } \sup_{x \in A_0, y \in A_1} d(ty + (1-t)x, A_0) \geq d(ty_1 + (1-t)x_1, A_0) = td(x_1, y_1).$$

For any $y \in A_1$ we have $\|(\lambda y + (1-\lambda)y_2) - x_2\| \geq \|y_2 - x_2\|$ for every $\lambda \in [0, 1]$, so $\langle y - y_2, y_2 - x_2 \rangle \geq 0$. For any $x \in A_0$

there exists $y_3 \in A_1$ such that $d(x, y_3) \leq d(x_2, y_2)$, then

$$\begin{aligned} \|y_3 - y_2 + y_2 - x_2 + x_2 - x\|^2 &= \|y_3 - x\|^2 \leq \|y_2 - x_2\|^2, \text{ so } \|y_3 - y_2 + x_2 - x\|^2 + \\ &+ 2\langle y_3 - y_2, y_2 - x_2 \rangle \leq 2\langle y_2 - x_2, x - x_2 \rangle. \text{ Because} \end{aligned}$$

$\langle y_3 - y_2, y_2 - x_2 \rangle \geq 0$ we have $\langle y_2 - x_2, x - x_2 \rangle \geq 0$. This implies

$$\text{that } \sup_{z \in A_0} d(z, A_t) \geq d(x_2, A_t) = \inf_{x \in A_0, y \in A_1} \|ty + (1-t)x - x_2\| =$$

$$= \inf_{x \in A_0, y \in A_1} \|t(y - y_2) + (1-t)(x - x_2) + t(y_2 - x_2)\| \geq t\|y_2 - x_2\| \text{ and}$$

$$\text{dist}(A_0, A_t) \geq \max(td(x_1, y_1), td(x_2, y_2)) = t \text{dist}(A_0, A_1), \text{ so}$$

$$\text{dist}(A_0, A_t) = t \text{dist}(A_0, A_1) \text{ and, analogously, } \text{dist}(A_t, A_1) =$$

$$= (1-t) \text{dist}(A_0, A_1). \text{ Therefore for any } s, t \in [0, 1] \text{ we obtain}$$

$$\text{dist}(A_s, A_t) = |s-t| \text{dist}(A_0, A_1),$$

so the arc $A_0 A_1 = \{A_t : 0 \leq t \leq 1\}$ has $\text{diameter} \leq \text{dist}(A_0, A_1)$.

Theorem. For every $n \geq 1$ there exists a compact convex set Q in \mathbb{R}^{n+2} such that every closed subset of B_n can be obtained as an intersection of Q with some k -dimensional affine subspace of \mathbb{R}^{n+2} .

Proof. By the Lemma and the Peano Theorem ([1], p. 246) it follows that there exists a continuous function ψ from the interval $[0, 1]$ onto \mathcal{C} . For $t \in [0, 1]$ we define

$$C_t = \psi(t) \times \{(\cos t, \sin t)\} \subset \mathbb{R}^{n+2}$$

and put

$$Q = \text{conv} \bigcup_{t \in [0, 1]} C_t.$$

The set Q is compact. Indeed, let $x_k = (x_k^1, \dots, x_k^n, \cos t_k, \sin t_k) \in Q$. Because of $\|x_k\| \leq \sqrt{2}$, there exists a subsequence $x_{k'}$ of x_k converging to some $x_0 = (x_0^1, \dots, x_0^n, \cos t_0, \sin t_0) \in \mathbb{R}^{n+2}$. Obviously $t_{k'} \rightarrow t_0$ and $y_{k'} = (x_{k'}^1, \dots, x_{k'}^n) \in \mathbb{R}^n$ converges to $y_0 = (x_0^1, \dots, x_0^n) \in \mathbb{R}^n$. We have $y_{k'} \in \psi(t_{k'})$ and $\text{dist}(\psi(t_{k'}), \psi(t_0)) \rightarrow 0$. By the remark preceding Lemma this implies that $y_0 \in \psi(t_0)$, so $x_0 \in Q$.

Since ψ is an onto mapping, for every convex subset D of B_n there exists $t \in [0, 1]$ such that $\psi(t) = D$ and for the k -dimensional affine subspace H_t of \mathbb{R}^{n+2} defined as $H_t = \mathbb{R}^n \times \{(\cos t, \sin t)\}$, we have

$$Q \cap H_t = D \times \{(\cos t, \sin t)\}.$$

Indeed, let $x \in Q \cap H_t$, then there exist elements $x_1 \in C_{t_1}$ and real numbers $\alpha_1, i=1, \dots, m$ such that $\sum \alpha_i = 1$ and $x = \sum \alpha_i x_i$. In particular $\sum \alpha_i (\cos t_1, \sin t_1) = (\cos t, \sin t)$. By the strict convexity of the unit disc in \mathbb{R}^2 this implies $(\cos t_1, \sin t_1) = (\cos t, \sin t)$, i.e.

$t_1 = t$ for $i=1, \dots, m$. Thus $x_1 \in C_t$, so $x \in D \times (\cos t, \sin t)$. Since the reverse inclusion is obvious, the proof is complete.

Let us observe that by an easy application of the Peano theorem together with some of the above arguments (for $n = 2$) the set

$$P = \bigcup_{t \in [0,1]} \{(x_1, x_2, t) : (x_1, x_2) \in \psi(t)\} \subset \mathbb{R}^3$$

satisfies the condition: Every closed convex set in \mathbb{R}^2 with diameter ≤ 1 can be obtained as the intersection of P with some plane (note that P is not convex).

We still do not know whether there exists a compact convex set in \mathbb{R}^3 with the above property.

References

- [1] R. Engelking: Outline of General Topology. PWN, Warszawa 1968