

Gerald W. Johnson

The use of mixed norms: Two examples

In: Zdeněk Frolík (ed.): Abstracta. 8th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1980. pp. 104–107.

Persistent URL: <http://dml.cz/dmlcz/701187>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1980

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

The Use of Mixed Norms: Two Examples

by

G.W. Johnson

In a paper on "the Feynman integral" G.W. Johnson and D.L. Skong [JS1] found a certain "mixed norm" useful. Once oriented in this direction they have found mixed norms useful in several places in their work on the Feynman integral and related problems [JS2-JS6]. In a paper on thin sets in harmonic analysis, specifically, on "p-Sidon sets", G.W. Johnson and G.S. Woodward [JW] proved a certain mixed-norm inequality which was crucial for the results of that paper. In the two examples which will be discussed, mixed norms were not involved in the statement of the problem but they were involved in the solution.

My experience with the use of mixed norms has led me to the strong conviction that the mixed-norm point of view could be used to advantage in many more situations if more mathematicians thought in these terms. In some cases it seems to enable one to solve problems that could not otherwise be solved.

G.W. Johnson and L.V. Petersen [P and JP] have attempted to give the beginnings of a general framework for considering mixed norm results. W.A.J. Luxemburg [L] has independently developed much the same theory; indeed a more satisfactory form of the key theorem will appear in [L].

1st Example: Let $C_0[0, t]$ denote the real-valued continuous functions X on $[0, t]$ such that $X(0) = 0$. Given a function $F : C_0[0, t] \rightarrow \mathbb{C}$ (the complex numbers) one can define $J(F)$, the analytic operator-valued Feynman integral of F [CS].

If it exists, $J(F)$ is a bounded linear operator on $L_2(\mathbb{R})$.

Let $0 < t_1 < \dots < t_n \leq t$ ($n \geq 3$) and let $f : \mathbb{R}^n \rightarrow \mathbb{C}$. Let $F(X) = f(X(t_1), \dots, X(t_n))$ for X in $C_0[0, t]$. Johnson and Skoug sought some L_p condition on f insuring the existence of $J(F)$ but instead found

Theorem: [JS1] There exists f in $\bigcap_{1 \leq p \leq \infty} L_p(\mathbb{R}^n)$ such that

$J(F)$ fails to exist.

On the other hand, they proved,

Theorem: [JS1] If $\|f\|_2 \|\dots\|_2 < \infty$, then $J(F)$ exists where

$$\|f\|_2 \|\dots\|_2 = \left\{ \int_{\mathbb{R}} \left[\int_{\mathbb{R}} \dots \int_{\mathbb{R}} \dots \int_{\mathbb{R}} |f(u_1, \dots, u_n)|^2 du_1 \right]^{1/2} \cdot du_2 \dots du_{n-1} \right\}^{1/2}.$$

2nd Example: Let $1 \leq p < 2$. A subset E of the integers is said to be p -Sidon ($E \in S_p$) if there exists a constant C such that for every finite subset F of E and any subset $\{c_n : n \in F\}$ of complex numbers indexed by F we have

$$\left(\sum_{n \in F} |c_n|^p \right)^{1/p} \leq C \left\| \sum_{n \in F} c_n e^{-int} \right\|_{\infty}.$$

(See [ER, LR, JW].) If $1 \leq p_1 < p_2 < 2$, clearly $S_{p_1} \subset S_{p_2}$.

In the deepest theorem in the paper [ER] it is shown that

Theorem: [ER] If $1 \leq p_1 < 4/3 \leq p_2 < 2$, then $S_{p_1} \not\subset S_{p_2}$.

A key element in the argument was a mixed norm inequality of Littlewood [Li]:

$$\sum_{i=1}^{\infty} \left(\sum_{j=1}^{\infty} |a_{ij}|^2 \right)^{1/2} + \sum_{j=1}^{\infty} \left(\sum_{i=1}^{\infty} |a_{ij}|^2 \right)^{1/2} \geq \left(\sum_{i,j} |a_{ij}|^{4/3} \right)^{3/4}.$$

[JW] extended the result of [ER]. One of the two substantial difficulties was finding and proving the following exten-

sion of Littlewood's inequality.

$$\sum_{k=1}^n \sum_{i_k} \left(\sum_{(i_k)} |a_{i_1, \dots, i_n}|^2 \right)^{1/2} \geq \left(\sum_{i_1, \dots, i_n} |a_{i_1, \dots, i_n}|^{\frac{2n}{n+1}} \right)^{\frac{n+1}{2n}}.$$

$[(i_k)]$ indicates summation over all the indices except the k^{th} .

Theorem: [JW] If $1 \leq p_1 < 4/3 \leq p_2 < 6/4 \leq \dots \leq p_{n-1} < \frac{2n}{n+1} \leq p_n < \dots$

then $S_{p_1} \subsetneq S_{p_2} \subsetneq \dots \subsetneq S_{p_{n-1}} \subsetneq S_{p_n} \subsetneq \dots$.

Recently R. Blei [B] has shown that $S_{p_1} \subsetneq S_{p_2}$ whenever

$1 \leq p_1 < p_2 < 2$. His argument involves a further extension and refinement of the old mixed norm inequality of Littlewood.

Bibliography

- [B] R. Blei: Fractional Cartesian product of sets, preprint
- [CS] R.H. Cameron and D.A. Storvick: An operator-valued function space integral and a related integral equation, J.Math. and Mech. 18 (1968), 517-552
- [ER] R.E. Edwards and K.A. Ross: p -Sidon sets, J.Func.Anal, 15 (1974), 404-427
- [JS1] G.W. Johnson and D.L. Skoug: Feynman integrals of non-factorable finite-dimensional functionals, Pacific J. of Math. 45 (1973), 257-267
- [JS2] ————— : Cameron and Storvick's function space integral for certain Banach spaces of functionals, J.London Math.Soc. 9 (1974), 103-117
- [JS3] ————— : Cameron and Storvick's function space integral for a Banach space of functionals generated by finite-dimensional functionals, Annali di Matematica Pura Ed Applicata 104 (1975), 67-83
- [JS4] ————— : The Cameron-Storvick function space integral: The L_1 theory, J.Math.Anal. and Appl. 50 (1975), 647-67

- [JS5] ——— : The Cameron-Storvick function space integral:
An $L(L_p, L_{p'})$ theory, Nagoya Math. J. 60 (1976), 93-137
- [JS6] ——— : An L_p analytic Fourier-Feynman transform,
Michigan J. of Math. (1979)
- [JP] G.W. Johnson and L.V. Petersen: On product Banach function
spaces
- [JW] G.W. Johnson and G.S. Woodward: On p -Sidon sets, Indiana
Math. J. 24 (1974), 161-167
- [Li] J.E. Littlewood: On bounded bilinear forms in an infinite
number of variable, Quarterly J. of Math. 1 (1930),
164-174
- [LR] Lopez and K.A. Ross: Sidon Sets (1975), Marcel Delcer
- [L] W.A.J. Luxemburg: manuscript in preparation
- [P] L.V. Peterson: Thesis, U. of Nebraska, 1976