

Marek Wilhelm

Open relations

In: Zdeněk Frolík (ed.): Abstracta. 8th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1980. pp. 199–202.

Persistent URL: <http://dml.cz/dmlcz/701208>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1980

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Open relations

M. Wilhelm

The letters X and Y will stand for topological spaces, G and H for topological groups, E and F for topological vector spaces over \mathbb{R} or \mathbb{C} . The letter Γ will stand for a relation in $X \times Y$, or in $G \times H$, or in $E \times F$. Given $A \subseteq X$, $\Gamma(A)$ denotes the image set $\{y \in Y : (x, y) \in \Gamma \text{ for some } x \in A\}$. Γ is said to be open if the image of each open set is open; Γ is open at a point $y \in Y$ if for each open set U in X , $y \in \Gamma(U)$ implies $y \in \text{Int } \Gamma(U)$. Γ is said to be nearly open if the image of each open set U is nearly open, i.e. $\Gamma(U) \subseteq \text{Int } \overline{\Gamma(U)}$; Γ is nearly open at $y \in Y$ if for each open subset U of X , $y \in \Gamma(U)$ implies $y \in \text{Int } \overline{\Gamma(U)}$. The set of all points at which Γ is nearly open will be denoted by $\text{NO}(\Gamma)$.

Theorem 1. Let Γ be a surjective relation.

- (i) If X is second-countable and Y is second category, then $\text{NO}(\Gamma)$ is non-empty.
- (ii) If X is second-countable and Y is Baire, then $\text{NO}(\Gamma)$ is dense in Y .
- (iii) If G is separable or Lindelöf, H is second category, and Γ is a subgroup of $G \times H$, then Γ is nearly open.
- (iv) If F is second category and Γ is a vector subspace of $E \times F$, then Γ is nearly open.
- (v) If E and F are locally convex, F is barrelled, and Γ is a vector subspace of $E \times F$, then Γ is nearly open.

Consider the following condition strictly related to Banach's Open Mapping and Closed Graph Theorems:

- (B) If Γ is nearly open, then Γ is open.

Theorem 2. If X is locally compact and Γ is a closed subset of $X \times Y$, then (B) holds.

Γ is said to be separating if for each two points $x_1 \in X$ there exist open sets U_i in X containing x_i ($i = 1, 2$) such that $\overline{\Gamma(U_1)} \cap \Gamma(U_2) = \emptyset$. This implies that Γ is an injective relation which is a closed subset of $X \times \Gamma(X)$, and that $\Gamma^*(X) \in T_2$.

Theorem 3 [10]. If X is Čech-complete and Γ is separating and surjective, then (B) holds.

Theorem 3 partially supports Conjecture from [5].

Corollary [2]. If X is Čech-complete, $Y \in T_2$ and f is a bijection of X onto Y , then (B) holds (for $\Gamma = f$) provided either Y is Čech-complete or f is continuous.

The following two examples answer some questions from [6].

Example 1 [2]. There exists a separable complete metric space X and a continuous nearly open surjection $f : X \rightarrow [0, 1]$ which is not open.

Example 2 [9]. There exists a separable complete metric space X , a separable second category metric space Y and a nearly continuous nearly open bijection f of X onto Y having a closed graph, which is neither continuous nor open.

Theorem 4 [8, 10]. If G is Čech-complete and Γ is a closed subgroup of $G \times H$, then (B) holds.

Some partial results in the direction of Theorem 4 were previously obtained e.g. in [4, 3, 1] (the last under the assumption that Γ is a continuous function). For locally convex spaces the result is in [7] (proof based on duality theory). Čech-completeness cannot be replaced with completeness in the left

(or two-sided) uniformity.

In the following X and Y stand for uniform spaces, and $U(V)$ stand for members of the uniformities on X (resp. Y). Γ is said to be uniformly open if $\forall U \exists V \forall x, \Gamma(U(x)) \supset V(\Gamma(x))$; Γ is uniformly nearly open if $\forall U \exists V \forall x, \overline{\Gamma(U(x))} \supset V(\Gamma(x))$.

Consider the condition

(B') If Γ is uniformly nearly open, then Γ is uniformly open.

Theorem 5 [10]. If X is uniformly Čech-complete and Γ is a closed subset of $X \times Y$, then (B') holds.

For complete metric space the result is in [4]. Theorem 5 answers some question from [6].

The notion of uniform Čech-completeness ($U\check{C}C$) is studied in [10]; X is $U\check{C}C$ if there exists a countable complete family of uniform covers of X . If X is $U\check{C}C$, then X is complete and (topologically) Čech-complete and paracompact. If X is uniformly locally compact, then X is $U\check{C}C$. If a uniform space X is metrizable, then X is $U\check{C}C$ iff X is complete. If a topological space X is Čech-complete and paracompact, then X is $U\check{C}C$ with respect to the finest uniformity on X . X is $U\check{C}C$ iff there exists a perfect uniformly continuous mapping of X onto a complete metric space. A closed subspace of a $U\check{C}C$ space is $U\check{C}C$. Countable product of $U\check{C}C$ spaces is $U\check{C}C$. A topological group G is $U\check{C}C$ with respect to the two-sided uniformity on G iff G is Čech-complete; G is $U\check{C}C$ with respect to the left uniformity on G iff G is Čech-complete and left-complete. A topological vector space E (over a number field) which is $U\check{C}C$ (with respect to the translation-invariant uniformity on E) admits a complete metric.

References

- [1] L.G.Brown, Topologically complete groups, Proc.Amer.Math. Soc. 35 (1972), 593-600.
- [2] T.Byczkowski and R.Pol, On the Closed Graph and Open Mapping Theorems, Bull.Acad.Polon.Sci. 24 (1976), 723-726.
- [3] T.Husain, Introduction to Topological Groups, Philadelphia 1966.
- [4] J.L.Kelley, General Topology, Princeton 1955.
- [5] P.Mah and S.A.Naimpally, Open and uniformly open relations, Proc.Amer.Math.Soc. 66 (1977), 159-166.
- [6] B.J.Pettis, Some topological questions related to Open Mapping and Closed Graph Theorems, Studies in Topology, New York 1975, 451-456.
- [7] V.Pták, On the closed graph theorem, Czech.Math.Journ. 9 (1959), 69-72,
- [8] M.Wilhelm, Relations among some closed graph and open mapping theorems, Colloq.Math. 42 (1979) (to appear).
- [9] M.Wilhelm, On a question of B.J.Pettis, Bull.Acad.Polon. Sci. (to appear).
- [10] M.Wilhelm, Criteria of openness for relations, Fund.Math. (to appear).