

Ivan Kolář; Jan Slovák

On the geometric functors on manifolds

In: Jarolím Bureš and Vladimír Souček (eds.): Proceedings of the Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1989. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 21. pp. [223]–233.

Persistent URL: <http://dml.cz/dmlcz/701443>

Terms of use:

© Circolo Matematico di Palermo, 1989

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE GEOMETRIC FUNCTORS ON MANIFOLDS

Ivan Kolář, Jan Slovák

A. Nijenhuis pointed out that the classical bundles of geometric objects can be viewed as certain functors transforming manifolds and their local diffeomorphisms into fibred manifolds and their morphisms, [10]. Some of those functors are defined on the whole category \underline{Mf} of all manifolds and all smooth maps. However, one can observe that several general aspects of the theory of the bundle functors on the whole category \underline{Mf} are rather different from the case of the classical bundles of geometric objects. Recently it has been deduced, [1], [5], [7], that the product-preserving bundle functors on \underline{Mf} coincide with the functors defined by means of the finite-dimensional local algebras by A. Weil, [15]. The main aim of our present paper are some general properties of the non-product-preserving bundle functors on \underline{Mf} . We deduce that the fibres of the bundle functors with the so-called point property, [6], are diffeomorphic to numerical spaces. We show that the product-preserving functors are fully characterized by the corresponding condition on dimensions. We prove that every bundle functor on \underline{Mf} satisfies the so-called prolongation axiom by J. Pradines, [13]. Finally we deduce that the bundle functors without the point property can be interpreted as certain parametrized systems of functors with the point property. - We assume all manifolds and maps to be infinitely differentiable and all manifolds to be paracompact.

1. ORDER OF A BUNDLE FUNCTOR

Let \underline{Mf} be the category of all manifolds and all maps, \underline{FM} be the category of all fibred manifolds and their morphisms and $B: \underline{FM} \rightarrow \underline{Mf}$ be the base functor. Given a functor $F: \underline{Mf} \rightarrow \underline{FM}$ satisfy-

This paper is in final form and no version of it will be submitted for publication elsewhere.

ing $B \circ F = \text{id}_{\underline{Mf}}$, we denote by $p_M: FM \rightarrow M$ its value on a manifold M and by $F_x f: F_x M \rightarrow F_{f(x)} N$ the restriction of its value $Ff: FM \rightarrow FN$ on $f: M \rightarrow N$ to the fibres of FM over x and of FN over $f(x)$, $x \in M$.

Definition 1. A bundle functor on \underline{Mf} is a functor $F: \underline{Mf} \rightarrow \underline{FM}$ satisfying $B \circ F = \text{id}_{\underline{Mf}}$ and the localization condition: if $i: U \hookrightarrow M$ is the inclusion of an open subset, then $FU = p_M^{-1}(U)$ and Fi is the inclusion $p_M^{-1}(U) \hookrightarrow FM$.

If we replace the category \underline{Mf} by the category \underline{Mf}_m of all m -dimensional manifolds and their local diffeomorphisms, we obtain the classical concept of a natural bundle in dimension m by Nijenhuis, [10], and Palais-Terng, [11]. Hence the restriction F_m of a bundle functor F on \underline{Mf} to \underline{Mf}_m is a natural bundle in dimension m .

Let M, N, P be manifolds. A parametrized system of smooth maps $f_p: M \rightarrow N$, $p \in P$ is said to be smoothly parametrized, if the resulting map $f: M \times P \rightarrow N$ is smooth. The following result was deduced in a more general context in [14], but we find it useful to present a direct proof here.

Proposition 1. Every bundle functor $F: \underline{Mf} \rightarrow \underline{FM}$ satisfies the regularity condition: if $f: M \times P \rightarrow N$ is a smoothly parametrized family, then the family $\tilde{F}f: FM \times P \rightarrow FN$ defined by $(\tilde{F}f)_p = F(f_p)$ is also smoothly parametrized.

Proof. Since smoothness is local property, it suffices to discuss the case $f: R^m \times R^k \rightarrow N$. Denote by $i: R^m \rightarrow R^m \times R^k$ the injection $x \mapsto (x, 0)$ and by t_p the translation on $R^m \times R^k$ transforming the origin into $(0, p)$, $p \in R^k$. A deep analytical result by Epstein and Thurston reads that Ft_p , $p \in R^k$, is a smoothly parametrized family, [2]. We have $f_p = f \circ t_p \circ i$, so that $Ff_p = Ff \circ Ft_p \circ Fi$ is a smoothly parametrized family as well, QED.

According to Palais-Terng, [11], every natural bundle F_m has a finite order $r(m)$, i.e. $j^{r(m)} f(x) = j^{r(m)} g(x)$ implies $F_x f = F_x g$ for any two local diffeomorphisms of m -dimensional manifolds. We deduce a similar result for arbitrary maps. Write $m = \dim M$, $n = \dim N$ and $r(m, n) = r(\max(m, n))$.

Proposition 2. For any maps $f, g: M \rightarrow N$, $j^{r(m, n)} f(x) = j^{r(m, n)} g(x)$ implies $F_x f = F_x g$.

Proof. By locality, it suffices to consider two maps $f, g: R^m \rightarrow R^n$.

We have to discuss three cases.

I. Let any two maps $f, g: R^m \rightarrow R^m$ satisfy $j^r f(x) = j^r g(x)$ with $r = r(m)$. Consider one-parameter families $f_t = f + t \text{id}_{R^m}$, $g_t = g + t \text{id}_{R^m}$, $t \in R$. Since their Jacobians at x are certain non-zero

polynomials in t , f_t and g_t are local diffeomorphisms in a neighbourhood of x except a finite number values of t . Since $j^r f_t(x) = j^r g_t(x)$ for all t , the classical result, [11], implies $F_x f_t = F_x g_t$ except a finite number values of t . Then the regularity condition yields $F_x f_0 = F_x g_0$.

II. Let $m = n+k$, $k > 0$, and $f, g: R^{n+k} \rightarrow R^n$ satisfy $j^r f(x) = j^r g(x)$ with $r = r(m)$. Consider $\bar{f} = (f, \text{pr}_2)$, $\bar{g} = (g, \text{pr}_2): R^m \rightarrow R^m$, where $\text{pr}_2: R^n \times R^k \rightarrow R^k$ is the second product projection. Obviously, it holds $j^r \bar{f}(x) = j^r \bar{g}(x)$. Since $f = \text{pr}_1 \circ \bar{f}$, $g = \text{pr}_1 \circ \bar{g}$, functoriality and I. imply $F_x f = F_{\bar{f}(x)} \text{pr}_1 \circ F_x \bar{f} = F_{\bar{g}(x)} \text{pr}_1 \circ F_x \bar{g} = F_x g$.

III. Let $m+k = n, k > 0$ and $f, g: R^m \rightarrow R^{m+k}$ satisfy $j^r f(x) = j^r g(x)$ with $r = r(n)$. Consider $\bar{f} = f \circ \text{pr}_1$, $\bar{g} = g \circ \text{pr}_1: R^n \rightarrow R^n$, where $\text{pr}_1: R^m \times R^k \rightarrow R^m$ is the first product projection. We have $j^r \bar{f}(y) = j^r \bar{g}(y)$ for every y satisfying $\text{pr}_1(y) = x$. Since $f = \bar{f} \circ i$, $g = \bar{g} \circ i$, functoriality and I. yield $F_x f = F_{i(x)} \bar{f} \circ F_x i = F_{i(x)} \bar{g} \circ F_x i = F_x g$, QED.

Remark 1. If F is a product-preserving functor, it is a Weil functor, [1], [5], [7]. If r is the order of the corresponding local algebra, it holds $r(m) = r$ for all dimensions m . On the other hand, Mikulski has constructed a non-product-preserving functor of infinite order, i.e. with an unbounded sequence of $r(m)$, [8]:

2. FUNCTORS WITH THE POINT PROPERTY

Let pt denote a one-point manifold.

Definition 2, [6]. A bundle functor $F: \underline{Mf} \rightarrow \underline{FM}$ is said to have the point property, if $F(pt) = pt$.

Obviously, every product-preserving functor has the point property. An example of a non-product-preserving functor with the point property is the r -th order tangent functor in the sense of

F. W. Pohl, [12], [6].

An interesting feature of the bundle functors with the point property is the existence of natural canonical sections $c_M: M \rightarrow FM$ defined by $c_M(x) = Fi_x(pt)$, where $i_x: pt \rightarrow M$ is the injection $i_x(pt) = x$ of pt into $x \in M$. The regularity condition of Proposition 1 implies that c_M are smooth maps. Naturality of those maps means $c_N \circ f = Ff \circ c_M$ for all maps $f: M \rightarrow N$, which follows directly from the definition.

Proposition 3. If F is a bundle functor with the point property, then every fibre $F_x M$ is diffeomorphic to a numerical space $R^{k(m)}$, $m = \dim M$.

The proof is based on a lemma from differential topology.

Lemma. Let S be a paracompact m -dimensional manifold and $s \in S$ be a point. Let h_t be a smoothly parametrized system of maps, $t \in R$, $h_1 = id_S$, $h_0(S) = \{s\}$ and let h_t be diffeomorphisms for all $t \neq 0$. Then S is diffeomorphic to R^m .

Proof. We first recall a well known fact that if $S = \bigcup_{k=0}^{\infty} S_k$, where

S_k are open submanifolds diffeomorphic to R^m and $S_k \subset S_{k+1}$ for all

k , then S is diffeomorphic to R^m , see [3], Chapter 1, § 2. We are

going to construct a sequence with the latter property. Choose an increasing sequence of relatively compact open submanifolds

$K_n \subset K_{n+1} \subset S$, $S = \bigcup_{k=1}^{\infty} K_n$ and take a relatively compact neighbourhood

U of the point s diffeomorphic to R^m . Put $S_0 = U$. Since S_0 is relatively compact, there exist an integer n_1 with $K_{n_1} \supset S_0$ and a

$t_1 > 0$ with $h_{t_1}(K_{n_1}) \subset U$. Then we define $S_1 = (h_{t_1})^{-1}(U)$. We have

$S_1 \supset K_{n_1} \supset S_0$, S_1 is relatively compact and diffeomorphic to R^m .

Iterating this procedure, we construct S_k and n_k satisfying

$S_k \supset K_{n_k} \supset S_{k-1}$, $n_k > n_{k-1}$, QED.

Proof of Proposition 3. It suffices to deduce that $F_0 R^m$ is diffeomorphic to $R^{k(m)}$. Write $S = F_0 R^m$, $s = c_{R^m}(0)$ with $0 \in R^m$ and let

$g_t: R^m \rightarrow R^m$ be the system of homotheties $g_t(x) = tx$, $t \in R$. Since

$g_t(0) = 0$ for all t and g_0 coincides with the composition

$R^m \rightarrow pt \rightarrow \{0\}$, the smoothly parametrized system $h_t = Fg_t|S$ satisfies all assumptions of the Lemma, QED.

3. PRODUCTS AND DIMENSIONS

Let F be an arbitrary bundle functor on \underline{Mf} .

Proposition 4. If $f: M \rightarrow N$ is an immersion, then $Ff: FM \rightarrow FN$ is an immersion as well.

Proof. It suffices to discuss an immersion in its local canonical form $i: R^m \rightarrow R^n = R^{m+k}$, $x \mapsto (x, 0)$. Since the canonical projection $p: R^n = R^{m+k} \rightarrow R^m$ satisfies $p \circ i = id_{R^m}$, we have $Fp \circ Fi = id_{FR^m}$.

Hence $TFp \circ T_y Fi = id$ for all $y \in FR^m$, which implies that Fi is an immersion.

Remark 2. It is well known that connected paracompact smooth manifolds are exactly smooth neighbourhood retracts in numerical spaces, see [3], Chapter 1, § 1. A direct consequence is that a smooth map $f: M \rightarrow N$ is an embedding if and only if there are an open submanifold $U \subset N$ with $f(M) \subset U$ and a map $g: U \rightarrow M$ satisfying $g \circ f = id_M$. This implies that any bundle functor F on \underline{Mf} transforms embeddings into embeddings. Indeed, if $f: M \rightarrow N$ is an embedding and $g: U \rightarrow M$ is the above map, then $Fg \circ Ff = id_{FM}$ and FU is an open submanifold in FN , so that Ff is an embedding as well.

For technical reasons, we study the bundle functors with the point property in the rest of this section, but we shall show in the next section that similar results hold for arbitrary bundle functors on \underline{Mf} . Consider a product of two manifolds $M \xleftarrow{p} M \times N \xrightarrow{q} N$. The induced maps $FM \xleftarrow{Fp} F(M \times N) \xrightarrow{Fq} FN$ determine a canonical map $\pi: F(M \times N) \rightarrow FM \times FN$.

Proposition 5. If F has the point property then $\pi: F(M \times N) \rightarrow FM \times FN$ is a surjective submersion.

Proof. It suffices to discuss the case $M = R^m$, $N = R^n$. Write $O_1 = c_{R^m}(0) \in FR^m$, $O_2 = c_{R^n}(0) \in FR^n$, $O_3 = c_{R^{m+n}}(0) \in FR^{m+n}$. Let $i: R^m \rightarrow R^{m+n}$, $x \mapsto (x, 0)$ and $j: R^n \rightarrow R^{m+n}$, $y \mapsto (0, y)$ be the canonical injections. In the tangent space $T_{O_3} F(R^{m+n})$ we have two sub-

spaces $V = \text{TFi}(T_{O_1} \mathbb{R}^m)$ and $W = \text{TFj}(T_{O_2} \mathbb{R}^n)$. We are going to deduce

$V \cap W = 0$. Let $A \in V \cap W$, $A = \text{TFi}(B) = \text{TFj}(C)$, $B \in T_{O_1} \mathbb{R}^m$, $C \in T_{O_2} \mathbb{R}^n$.

On one hand, $p \circ i = \text{id}_{\mathbb{R}^m}$ implies $\text{TFp}(A) = \text{TFp}(\text{TFi}(B)) = B$. On the

other hand, $p \circ j$ is the constant map of \mathbb{R}^n into the zero point of \mathbb{R}^m . The latter map can be factorized as $\mathbb{R}^n \rightarrow \text{pt} \xrightarrow{i_0} \mathbb{R}^m$. Hence

$\text{Fp} \circ \text{Fj}$ is the constant map of \mathbb{R}^n into O_1 . This implies $0 = \text{TFp} \circ \text{TFj}(C) = \text{TFp}(A) = B$. Hence $V \cap W = 0$, so that π is a submersion at $O_3 \in \mathbb{F}\mathbb{R}^{m+n}$ and consequently on a neighbourhood $U \subset \mathbb{F}\mathbb{R}^{m+n}$ of O_3 .

Since all homotheties g_t on \mathbb{R}^m , \mathbb{R}^n and \mathbb{R}^{m+n} commute with the product projections and π is induced by Fp and Fq , the images $\text{F}g_t$ commute with π as well. The family $\text{F}g_t$ is smoothly parametrized

and $\text{F}g_0(\mathbb{F}\mathbb{R}^{m+n}) = \{O_3\}$, so that every point of $\mathbb{F}\mathbb{R}^{m+n}$ can be mapped into U by a suitable $\text{F}g_t$, $t > 0$. Taking into account that $\text{F}g_t$, $t > 0$, are diffeomorphisms, we see that π is a submersion. Therefore the image $\pi(\mathbb{F}\mathbb{R}^{m+n})$ is an open neighbourhood of $(O_1, O_2) \in \mathbb{F}\mathbb{R}^m \times \mathbb{F}\mathbb{R}^n$.

Similarly as above, every point of $\mathbb{F}\mathbb{R}^m \times \mathbb{F}\mathbb{R}^n$ can be mapped into

$\pi(\mathbb{F}\mathbb{R}^{m+n})$ by a suitable $\text{F}g_t$, $t > 0$. This implies that π is surjective, QED.

Remark 3. It is easy to check that Proposition 5 can be extended to an arbitrary finite product of manifolds,

Proposition 6. If F has the point property and $f: M \rightarrow N$ is a submersion, then $\text{F}f: \text{F}M \rightarrow \text{F}N$ is also a submersion.

Proof. It suffices to discuss a submersion in its local canonical form $p: \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}^n$. Then $\text{F}p = \text{pr}_1 \circ \pi$ is a composition of two submersions $\pi: \mathbb{F}(\mathbb{R}^n \times \mathbb{R}^k) \rightarrow \mathbb{F}\mathbb{R}^n \times \mathbb{F}\mathbb{R}^k$ and $\text{pr}_1: \mathbb{F}\mathbb{R}^n \times \mathbb{F}\mathbb{R}^k \rightarrow \mathbb{F}\mathbb{R}^n$, QED.

Proposition 7. If F has the point property, then $k(m+n) \geq k(m)+k(n)$. The equality holds if and only if F preserves products in dimensions m and n .

Proof. By Proposition 5, $\pi: \mathbb{F}(\mathbb{R}^m \times \mathbb{R}^n) \rightarrow \mathbb{F}\mathbb{R}^m \times \mathbb{F}\mathbb{R}^n$ is a submersion, which implies $k(m+n) \geq k(m)+k(n)$. If the equality holds, π is a local diffeomorphism at each point, so that π is a covering. But

$FR^m = R^m \times R^{k(m)}$ by Proposition 3, so that $FR^m \times FR^n$ is simply connected and π must be a global diffeomorphism. Dealing with arbitrary manifolds M, N , we obtain the result by the localization property of the bundle functors and by a standard diagram chasing, QED.

Corollary 1. A bundle functor $F: \underline{Mf} \rightarrow \underline{FM}$ preserves products if and only if $F(pt) = pt$ and $k(m) = mk(1)$ for all integers m .

Remark 4. An interesting consequence of our results is that a bundle functor F on \underline{Mf} with the point property can transform principal fibre bundles into principal fibre bundles if and only if F preserves products. More precisely, for any principal fibre bundle (P, p, M, G) we are looking for a natural principal fibre bundle structure on $Fp: FP \rightarrow FM$ with respect to a natural group structure on FG . This can be defined by prolongating all maps in question if F preserves products and this is impossible if F does not preserve products for the dimension reasons of Proposition 7. (We remark that there is a natural group structure on FG even for the non-product-preserving functor of the r -th order tangent vectors, [6].)

4. FUNCTORS WITHOUT THE POINT PROPERTY

Consider an arbitrary bundle functor $F: \underline{Mf} \rightarrow \underline{FM}$. Hence $Q = F(pt)$ is a manifold and the unique map $q_M: M \rightarrow pt$ induces $Fq_M: FM \rightarrow Q$. Similarly to § 2, every point $a \in Q$ determines canonical natural sections $c(a)_M: M \rightarrow FM$ defined by $c(a)_M(x) = Fi_x(a)$, where i_x is the injection of pt into $x \in M$. Let $G: \underline{Mf} \rightarrow \underline{FM}$ be the bundle functor defined by $GM = M \times Q$ and $Gf = f \times id_Q$ for all manifolds and maps.

Proposition 8. The maps $\sigma_M(x, a) = c(a)_M(x)$, $x \in M$, $a \in Q$, and $\zeta_M(z) = (p_M(z), Fq_M(z))$, $z \in FM$, define natural transformations $\sigma: G \rightarrow F$ and $\zeta: F \rightarrow G$ satisfying $\zeta \circ \sigma = id$. Moreover, σ_M is an embedding and ζ_M is a surjective submersion for every manifold M .

Proof. The proof of the first sentence is straightforward. By Remark 2, the equality $\zeta_M \circ \sigma_M = id_{M \times Q}$ implies that σ_M is an embedding. The latter equality also implies that ζ_M is surjective and has the maximal rank on a neighbourhood U of the image

$\mathcal{G}_M(M \times Q)$. It suffices to prove that every \mathcal{F}_{R^m} is a submersion.

Consider the homotheties $g_t(x) = tx$ on R^m , Then Fg_t is a smoothly parametrized family with $Fg_1 = \text{id}$ and $Fg_0(FR^m) = F\iota_0 \circ Fq_{R^m}(FR^m) \subset$

$\mathcal{G}_M(R^m \times Q)$. Hence every point of FR^m is mapped into U by some Fg_t , $t > 0$ and consequently \mathcal{F}_{R^m} has maximal rank everywhere, QED.

Corollary 2. For every $a \in Q$, the rule $F_a M = (Fq_M)^{-1}(a)$, $F_a f = Ff|_{F_a M}$ determines a bundle functor with the point property.

Corollary 3. Every bundle functor $F: \underline{Mf} \rightarrow \underline{FM}$ transforms submersions into submersions.

Proof. By Proposition 8, every induced map $Ff: FM \rightarrow FN$ is a base-preserving morphism of fibred manifold $Fq_M: FM \rightarrow Q$ into $Fq_N: FN \rightarrow Q$. If $f: M \rightarrow N$ is a submersion, then every $F_a f: F_a M \rightarrow F_a N$ is a submersion by Proposition 6. Hence Ff must be also a submersion, QED.

The following corollary was deduced by quite different methods by Mikulski, [9].

Corollary 4. Let $F: \underline{Mf} \rightarrow \underline{FM}$ be a bundle functor with compact fibres. Then F is naturally equivalent to a trivial bundle functor of order 0.

Proof. If the standard fibres of F are compact, then all functors F_a of Corollary 2 coincide with the identity functor on \underline{Mf} by Proposition 3. Hence the natural transformations \mathcal{G} and \mathcal{F} of Proposition 8 are natural equivalences, QED.

Remark 5. We remark that Proposition 5 does not hold for general bundle functors. However, using Propositions 7 and 8 and Corollary 2, we can deduce the inequality $k(m+n) \geq k(m) + k(n) - \dim Q$. Another simple consequence of Corollary 2 is that the equality holds if and only if all functors F_a , $a \in Q$, preserve products in dimensions m and n .

Let $f: Y \rightarrow X$ be a submersion and $FX \oplus Y$ be the pullback of Y with respect to $p_X: FX \rightarrow X$. Since $Ff: FY \rightarrow FX$ and $p_Y: FY \rightarrow Y$ satisfy $p_X \circ Ff = f \circ p_Y$, we have an induced pullback map $\mu: FY \rightarrow FX \oplus Y$.

Proposition 9. For every submersion $f: Y \rightarrow X$, the pullback map

$\mu: FY \rightarrow FX \oplus Y$ is also a submersion.

Proof. Taking into account the universal property of pullbacks, the fibration $\mathcal{P}_M: FM \rightarrow M \times Q$ from Proposition 8, the functors F_a , $a \in Q$ defined in Corollary 2 and standard diagram chasings, we may restrict ourselves to the functors with the point property. It suffices to discuss a submersion in its local canonical form $f: R^{m+n} \rightarrow R^m$, $f(x,y) = x$. Let $s: R^m \rightarrow R^{m+n}$ be the sections $x \mapsto (x,y)$, $y \in R^n$. Denoting by $FR^m \oplus R^{m+n}$ the Whitney sum over R^m , we define a map $t: FR^m \oplus R^{m+n} \rightarrow FR^{m+n}$ by $t(z, (x,y)) = Fs_y(z)$. Then t is smooth and $t(c_{R^m}(x), (x,y)) = c_{R^{m+n}}(x,y)$, so that we have constructed a smooth

section of μ through the values $c_{R^{m+n}}(x,y)$. Therefore μ is of

maximal rank on a neighbourhood of $c_{R^{m+n}}(R^{m+n})$ and the proof is

completed using the family of all homotheties on R^{m+n} in the same way as in the proof of Proposition 8, QED.

Remark 6. We can reformulate Proposition 9 by saying that every bundle functor on \underline{Mf} satisfies the so-called prolongation axiom introduced by Pradines, [13], in a more general situation.

The simplest example of a bundle functor without the point property can be constructed as follows. We take any bundle functor G with the point property and any manifold Q and we define $FM = GM \times Q$, $Ff = Gf \times id_Q$. We present an example showing that not all bundle functors on \underline{Mf} are of this type, not even if Q is connected. The basic idea of our example is that some of the individual "fibre components" F_a of our functor F coincide with the functor T_1^2 of 1-dimensional velocities of the second order and the other ones are the Whitney sum $T \oplus T$ of the tangent functor with itself in dependence on the zero values of a smooth function on Q . However, to give an exact proof of it, we shall use a formal procedure by Janyška, [4].

Example. Let L^2 be the category, the objects of which are the non-negative integers, the morphisms $L^2(m,n) = J_0^2(R^m, R^n)_0$ are the second order jets of R^m into R^n with source 0 and target 0 and the composition in L^2 is the composition of jets. Consider any manifold Q and define $S_n = Q \times R^n \times R^n$ for $n = 0, 1, \dots$. By [4], every action of L^2 on the system $S = \{S_0, S_1, \dots\}$ determines a second order bundle functor on \underline{Mf} . Let a_{1j}^p, a_{1j}^q be the canonical coordi-

ates on $L^2(m,n)$ and y^i, z^i be the canonical coordinates on $R^n \times R^n$. Take any smooth function $f: Q \rightarrow R$ and define

$$(a_{i,j}^p, a_{i,j}^p)(q, y^i, z^i) = (q, a_{i,j}^p y^i, f(q) a_{i,j}^p y^i y^j + a_{i,j}^p z^i)$$

$q \in Q$. One verifies easily that this really is an action of L^2 on S . Let F be the bundle functor defined by this action, so that $F(\text{pt}) = Q$. Obviously, if $f(q) = 0$, then the functor F_q in the sense of Corollary 2 is the Whitney sum $T \oplus T$. If $f(q) \neq 0$, then F_q is naturally equivalent to the functor T_1^2 mentioned above. This can be easily deduced by the Janyška's method: the maps $R^{2n} \rightarrow R^{2n}, y^i \mapsto y^i, z^i \mapsto f(q)z^i$ are L^2 -covariant and invertible, so that they determine a natural equivalence of T_1^2 into F_q .

REFERENCES

- [1] ECK D. J. "Product preserving functors on smooth manifolds", J. Pure Appl. Algebra 42 (1986), 133-140.
- [2] EPSTEIN D. B. A., THURSTON W. P. "Transformation groups and natural bundles", Proc. London Math. Soc. (3) 38 (1979), 219-236.
- [3] HIRSCH M. W. "Differential topology", Springer-Verlag 1976.
- [4] JANYŠKA J. "Geometric properties of prolongation functors", Časopis pěst. mat. 110 (1985), 77-86.
- [5] KAINZ G., MICHOR P. W. "Natural transformations in differential geometry", Czechoslovak Math. J. 37 (1987), 584-607.
- [6] KOLÁŘ I. "Functorial prolongations of Lie groups and their actions", Časopis pěst. mat. 108 (1983), 289-293.
- [7] LUCIANO O. O. "Categories of multiplicative functors and Morimoto's conjecture", Preprint N° 46, Institut Fourier, Laboratoire de Mathématiques, Grenoble, 1986.
- [8] MIKULSKI W. M. "There exists a prolongation functor of infinite order", to appear in Časopis pěst. mat.
- [9] MIKULSKI W. M. "Continuity of projections of natural homomorphisms", preprint.
- [10] NIJENHUIS A. "Natural bundles and their general properties", Differential Geometry, Kinokuniya, Tokyo 1972, 317-334.
- [11] PALAIS R. S., TERNG C. L. "Natural bundles have finite order", Topology 16 (1977), 271-277.
- [12] POHL F. W. "Differential geometry of higher order", Topology

- 1 (1962), 169-211.
- [13] PRADINES J. "Fibrés vectoriels double symétriques et jets holonomes d'ordre 2", CRAS Paris 264 (1967), A245-248.
- [14] SLOVÁK J. "Smooth structures on fibre jet spaces", Czechoslovak Math. J. 36 (1986), 358-375.
- [15] WEIL A. "Théorie des points proches sur les variétés différentiables", Colloque du CNRS, Strasbourg 1953, 111-117.

INSTITUT OF MATHEMATICS OF THE ČSAV,
BRANCH BRNO
MENDELOVO NÁM. 1
CS-66282 BRNO