

Nadine Acker

The maximal function of a complex measure

In: Jarolím Bureš and Vladimír Souček (eds.): Proceedings of the Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1991. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 26. pp. [41]–46.

Persistent URL: <http://dml.cz/dmlcz/701478>

Terms of use:

© Circolo Matematico di Palermo, 1991

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE MAXIMAL FUNCTION OF A COMPLEX MEASURE

Nadine Van Acker

ABSTRACT. The Marcinkiewicz Interpolation Theorem is proved in the setting of the unit sphere in euclidean space of arbitrary dimension. This leads to a key result in the study of the maximal function of a complex measure and, in particular, of an integrable function on this unit sphere.

1. INTRODUCTION

In the theory of the H^p spaces of monogenic functions [2], which is developed in the framework of Clifford analysis [1], an important role is played by the results concerning the boundary behaviour of Poisson integrals of complex measures and, in particular, of integrable functions on the unit sphere.

Let P denote the Poisson kernel in \mathring{B}_m , the unit ball in \mathbb{R}^m ; it is given by

$$P(x,y) = \frac{1-r^2}{|x-y|^m}$$

where $r^2 = |x|^2$ and y is on the unit sphere S^{m-1} .

Definition 1.1.

If μ is a complex measure on S^{m-1} , then its Poisson integral $\mathcal{P}[\mu]$ is defined as

$$\mathcal{P}[\mu](x) = \int_{S^{m-1}} P(x,y) d\mu(y), \quad x \in \mathring{B}_m.$$

In the special case where the measure μ is derived from an L_1 -function \tilde{f} , this definition rewrites as

$$\mathcal{P}[\tilde{f}](x) = \int_{S^{m-1}} P(x,y) \tilde{f}(y) d\sigma(y), \quad x \in \mathring{B}_m,$$

σ being the normalized Lebesgue measure on S^{m-1} , $\sigma(S^{m-1}) = 1$.

The main result about the boundary behaviour of Poisson integrals is the

so called Koranyi theorem, relating the maximum value of the continuous function $\mathcal{P}[\mu]$ in a cone D_α , to the value of the maximal function of μ at the top of the cone. Let us first introduce these notions explicitly before stating the theorem.

Definition 1.2.

Let $\xi \in S^{m-1}$ and $c_\alpha > 1$. The conical region D_α with top ξ is defined by

$$D_\alpha(\xi) = \{y \in \mathbb{R}^m : |\xi - y| < c_\alpha(1 - |y|)\}$$

Definition 1.3.

If μ is a complex measure on S^{m-1} , then its maximal function

$M\mu : S^{m-1} \rightarrow [0, +\infty]$ is given by :

$$M\mu(\xi) = \sup_{\psi > 0} \frac{|\mu|(bk(\xi, \psi))}{\sigma(bk(\xi, \psi))}$$

where $|\mu|$ is the total variation of μ and $bk(\xi, \psi)$ is a sphere segment centered at $\xi \in S^{m-1}$ with half a solid angle ψ .

Theorem 1.4. (Koranyi)

Given a complex measure μ on S^{m-1} , for each conical region D_α there exists a constant A_α such that on S^{m-1} :

$$\sup_{x \in D_\alpha(\xi)} |\mathcal{P}[\mu](x)| \leq A_\alpha M\mu(\xi) .$$

Proof : see [2].

The aim is to study, in the third section, this maximal function of a complex measure, especially when that measure is derived from an integrable function on S^{m-1} . The key result on the maximal function is strongly related to the so called Marcinkiewicz Interpolation Theorem, which we treat of first in the second section.

2. THE MARCINKIEWICZ INTERPOLATION THEOREM.

Let Σ denote a positive measure on S^{m-1} . We consider an operator T , acting on functions $\tilde{f} \in L_1(S^{m-1})$, and mapping them into Σ -measurable functions $T\tilde{f} : S^{m-1} \rightarrow [0, +\infty]$. Moreover we assume T to be subadditive :

$$T(\tilde{f} + \tilde{g}) \leq T\tilde{f} + T\tilde{g} \quad \text{for all } \tilde{f}, \tilde{g} \in L_1(S^{m-1}) .$$

Next we define numbers c_r , $1 \leq r \leq \infty$, as to be the smallest constants for which the estimates

$$\Sigma(\{T\tilde{f} > t\}) \leq c_r t^{-r} \int_{S^{m-1}} |\tilde{f}|^r d\Sigma \quad \text{and} \quad \|T\tilde{f}\|_\infty \leq c_\infty \|\tilde{f}\|_\infty$$

hold over the whole of S^{m-1} and for all $t, 0 < t < \infty$. Notice that the constants c_r might be ∞ . Finally we introduce the notation $K(a,b,c,\dots)$ for a constant K , depending upon the parameters a,b,c,\dots , and which is finite whenever all the parameters are finite.

Theorem 2.1.

For all $\tilde{f} \in L_p(S^{m-1})$ and $1 < p < r \leq \infty$,

$$\int_{S^{m-1}} (T\tilde{f})^p d\Sigma \leq K_p(c_1, c_r) \int_{S^{m-1}} |\tilde{f}|^p d\Sigma.$$

Proof :

Let $F : S^{m-1} \rightarrow [0, +\infty]$ be a Σ -measurable function. It is then easily seen that

$$\int_{S^{m-1}} F(\eta)^p d\Sigma(\eta) = \int_0^\infty pt^{p-1} \Sigma(\{F(\eta) > t\}) dt.$$

Given $t > 0$, consider the decomposition of $\tilde{f} \in L_1(S^{m-1})$:

$$\tilde{f} = \tilde{g}_t + \tilde{h}_t \quad \text{where} \quad \tilde{g}_t(\xi) = \begin{cases} 0 & \text{if } |\tilde{f}(\xi)| < t \\ \tilde{f}(\xi) & \text{if } |\tilde{f}(\xi)| \geq t \end{cases}$$

and where $\tilde{h}_t = \tilde{f} - \tilde{g}_t$. We define

$$G(t) = \Sigma(\{T\tilde{g}_t > \frac{t}{2}\}) \quad \text{and} \quad H(t) = \Sigma(\{T\tilde{h}_t > \frac{t}{2}\}).$$

Let $G_1(t)$ denote $\frac{2c_1}{t} \int_{|\tilde{f}| \geq t} |\tilde{f}| d\Sigma$, and, for given $r < \infty$,

$$H_1(t) = \frac{2^r c_r}{t^r} \int_{|\tilde{f}| < t} |\tilde{f}|^r d\Sigma.$$

It is then clear that

$$G(t) \leq c_1 \frac{2}{t} \int_{S^{m-1}} |\tilde{g}_t| d\Sigma \leq \frac{2c_1}{t} \int_{|\tilde{f}| \geq t} |\tilde{f}| d\Sigma = G_1(t),$$

and similarly, $H(t) \leq H_1(t)$.

Now applying Fubini's Theorem on

$$\int_0^\infty pt^{p-1} G_1(t) dt = 2c_1 p \int_0^\infty \int_{|\tilde{f}| \geq t} t^{p-2} |\tilde{f}| d\Sigma dt$$

we find that

$$\int_0^\infty pt^{p-1} G_1(t) dt = \frac{2c_1 p}{p-1} \int_{S^{m-1}} |\tilde{f}|^p d\Sigma.$$

Similarly, one gets

$$\begin{aligned} \int_0^{\infty} pt^{p-1} H_1(t) dt &= 2^r c_r p \int_{S^{m-1}} |\tilde{f}(\xi)|^r d\Sigma(\xi) \int_{|\tilde{f}(\xi)|}^{\infty} t^{p-r-1} dt \\ &= \frac{2^r c_r p}{r-p} \int_{S^{m-1}} |\tilde{f}|^p d\Sigma \end{aligned}$$

since $p < r$ by assumption.

In view of $T\tilde{f} \leq T\tilde{g}_t + T\tilde{h}_t$, it follows that

$$\Sigma(\{T\tilde{f} > t\}) \leq \Sigma(\{T\tilde{g}_t > \frac{t}{2}\}) + \Sigma(\{T\tilde{h}_t > \frac{t}{2}\}) \leq G_1(t) + H_1(t).$$

Combining the obtained results we find that

$$\begin{aligned} \int_{S^{m-1}} (T\tilde{f})^p d\Sigma &= \int_0^{\infty} pt^{p-1} \Sigma(\{T\tilde{f} > t\}) dt \leq \int_0^{\infty} pt^{p-1} G_1(t) dt + \int_0^{\infty} pt^{p-1} H_1(t) dt \\ &\leq K_p(c_1, c_r) \int_{S^{m-1}} |\tilde{f}|^p d\Sigma \end{aligned}$$

whence the desired result for $r < +\infty$.

If $r = +\infty$, we may assume without loss of generality that $c_{\infty} \leq \frac{1}{2}$ (if not, consider $\frac{T}{2c_{\infty}}$).

It is then easily seen that

$$\|\tilde{h}_t\|_{\infty} = \sup_{\xi \in S^{m-1}} |\tilde{h}_t(\xi)| = \sup_{|f| < t} |\tilde{f}(\xi)| < t$$

and hence $\|T\tilde{h}_t\|_{\infty} \leq c_{\infty} \|\tilde{h}_t\|_{\infty} \leq \frac{t}{2}$.

As $T\tilde{h}_t \leq \frac{t}{2}$ a.e. on S^{m-1} , it follows that $H(t) = \Sigma(\{T\tilde{h}_t > \frac{t}{2}\}) = 0$ leading to $\Sigma(\{T\tilde{f} > t\}) = G(t) \leq G_1(t)$. Combining the obtained results, the desired result indeed follows since

$$\begin{aligned} \int_{S^{m-1}} (T\tilde{f})^p d\Sigma &= \int_0^{\infty} pt^{p-1} \Sigma(\{T\tilde{f} > t\}) dt \\ &\leq \int_0^{\infty} pt^{p-1} G_1(t) dt \\ &= \frac{2pc_1}{p-1} \int_{S^{m-1}} |\tilde{f}|^p d\Sigma \\ &= K_p(c_1, c_{\infty}) \int_{S^{m-1}} |\tilde{f}|^p d\Sigma. \end{aligned}$$

3. THE MAXIMAL FUNCTION OF A COMPLEX MEASURE.

The notion of the maximal function of a complex measure μ on S^{m-1} was already introduced in the first section (definition 1.3). The following two propositions may be proved along classical lines.

Proposition 3.1.

The maximal function $M\mu$ of the complex measure μ on S^{m-1} is lower semi-continuous.

Proposition 3.2.

There exists a constant c_1 such that for all complex measures μ on S^{m-1} and all $\lambda > 0$:

$$\sigma(\{M\mu > \lambda\}) \leq c_1 \frac{\|\mu\|}{\lambda} .$$

Now consider the special case where the complex measure μ is derived from an L_1 -function : $\mu = \tilde{f}d\sigma$, $\tilde{f} \in L_1(S^{m-1})$. Its maximal function also reads

$$M\tilde{f}(\xi) = \sup_{\psi > 0} \frac{1}{\omega(\xi, \psi)} \int_{bk(\xi, \psi)} |\tilde{f}(\eta)| d\sigma(\eta)$$

where $\omega(\xi, \psi) = \sigma(bk(\xi, \psi))$.

From proposition 3.2 it now follows readily that

Corollary 3.3.

For all $\tilde{f} \in L_1(S^{m-1})$ and all $\lambda > 0$: $\sigma(\{M\tilde{f} > \lambda\}) \leq \frac{2m}{\lambda} \|\tilde{f}\|_1$.

The following proposition is also easily verified :

Proposition 3.4

The operator $M : \tilde{f} \in L_1(S^{m-1}) \mapsto M\tilde{f}$ is subadditive and moreover satisfies

$$M\tilde{f}(\xi) \leq \|\tilde{f}\|_\infty , \quad \xi \in S^{m-1} .$$

Combining the above results with the Marcinkiewicz Interpolation Theorem of the second section, we finally arrive at the main result

Theorem 3.5.

Let $\tilde{f} \in L_p(S^{m-1})$, $1 < p < \infty$, then there exists a constant $A(p)$ such that

$$\int_{S^{m-1}} |M\tilde{f}|^p d\sigma \leq A(p) \int_{S^{m-1}} |\tilde{f}|^p d\sigma .$$

REFERENCES

- [1] BRACKX F., DELANGHE R., SOMMEN F. "Clifford Analysis", Pitman, London, 1982.
- [2] VAN ACKER N. " H^p Spaces of monogenic functions", Proceedings of the Second Workshop on Clifford Algebras and their Applications in Mathematical Physics, Montpellier, 1989.

NADINE VAN ACKER
UNIVERSITY OF GHENT
SEMINAR OF MATHEMATICAL ANALYSIS
SINT PIETERSNIEUWSTRAAT 39
B-9000 GENT
BELGIUM