

Antonella Cabras; Ivan Kolář

On the second order absolute differentiation

In: Jan Slovák and Martin Čadek (eds.): Proceedings of the 18th Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1999. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 59. pp. 123–133.

Persistent URL: <http://dml.cz/dmlcz/701631>

Terms of use:

© Circolo Matematico di Palermo, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE SECOND ORDER ABSOLUTE DIFFERENTIATION

ANTONELLA CABRAS, IVAN KOLÁŘ

ABSTRACT. First we compare two different approaches to the second order absolute differentiation on an arbitrary fibered manifold. Then we extend the second approach to connections on the functional bundle of all smooth maps between the fibers over the same base point of two fibered manifolds over the same base. (For the first approach, this problem was solved in [4].)

There are two different approaches to the second order absolute differentiation in the case of a principal or linear connection Γ . The first one constructs $\nabla_{\Gamma, \Lambda}^2$ by means of an auxiliary linear connection Λ on the base manifold, [17], which is related to the ideas of tensor calculus. The second one applies another geometric idea by C. Ehresmann, [6], and constructs ∇_{Γ}^2 by means of Γ only. In Section 1 we recall the first construction in the case of a connection Γ on an arbitrary fibered manifold $\pi : Y \rightarrow M$, which has been developed recently in [1]. In Section 2 we generalize Ehresmann's approach to connections on a finite-dimensional groupoid to the groupoid $\mathcal{G}Y$ of all diffeomorphisms between the individual fibers of Y . We use systematically the structure of a smooth space in the sense of Frölicher on $\mathcal{G}Y$. The groupoid approach clarifies directly that the values of ∇_{Γ}^2 are semiholonomic 2-jets. But it is remarkable that it also interprets some prolongation procedures for connections on Y from a new point of view. In Section 3 we present a construction of ∇_{Γ}^2 by using second tangent bundles, which we need for a generalization in Section 6. Then we comment on some differences between $\nabla_{\Gamma, \Lambda}^2$ and ∇_{Γ}^2 .

The second part of the present paper is devoted to a functional version of the second order absolute differentiation. Consider two locally trivial fibered manifolds $p_1 : Y_1 \rightarrow M$, $p_2 : Y_2 \rightarrow M$ over the same base and the bundle of all fiber maps

$$(1) \quad \mathcal{F}(Y_1, Y_2) = \bigcup_{x \in M} C^\infty(Y_{1x}, Y_{2x}),$$

1991 *Mathematics Subject Classification.* 53C05, 58A20.

Key words and phrases. Second order absolute differentiation, bundle of smooth maps, connection on a functional bundle, semiholonomic 2-jet.

This work has been performed during the visit of I. Kolář at Dipartimento di Matematica Applicata "G. Sansone", Università di Firenze, supported by G.N.S.A.G.A. of C.N.R. The second author was also supported by a grant of the GA ČR No 201/96/0079.

This paper is in final form and no version of it will be published elsewhere.

which is a smooth space in the sense of Frölicher, [2]. The first approach to the second order absolute differentiation on $\mathcal{F}(Y_1, Y_2)$ was studied in [4], so that we go directly to the second one. In Section 5 we define the absolute differential $\nabla_\Gamma f$ of any smooth map f of a manifold N into $\mathcal{F}(Y_1, Y_2)$ with respect to a connection Γ on $\mathcal{F}(Y_1, Y_2)$. Then we construct $\nabla_\Gamma^2 f$ by using the machinery of second tangent bundles. In Section 7 we deduce for a finite order connection Γ on $\mathcal{F}(Y_1, Y_2)$ and a section $s : M \rightarrow \mathcal{F}(Y_1, Y_2)$ that the deviation of semiholonomic 2-jet $\nabla_\Gamma^2 s(x)$ coincides up to the sign with the curvature of Γ at $s(x)$.

If we deal with finite dimensional manifolds and maps between them, we always assume they are of class C^∞ , i.e. smooth in the classical sense. On the other hand, the concept of smoothness in the infinite dimension is due to Frölicher, [7], see also [3].

1. An auxiliary linear connection on the base. On an arbitrary fibered manifold $\pi : Y \rightarrow M$, a connection can be defined as a section $\Gamma : Y \rightarrow J^1 Y$, see e.g. [13]. We denote by $v_\Gamma : TY \rightarrow VY$ its vertical projection. If $s : M \rightarrow Y$ is a section, we define its absolute differential by

$$(2) \quad \nabla_\Gamma s = v_\Gamma \circ Ts,$$

i.e. we construct the vertical projection of the tangent map of s . Hence $\nabla_\Gamma s$ is a section $M \rightarrow VY \otimes T^*M$. Let x^i, y^p be some local fiber coordinates on Y and let Γ be expressed by

$$(3) \quad dy^p = F_i^p(x, y) dx^i$$

and s by $y^p = s^p(x)$. Then the coordinate form of (2) is

$$(4) \quad \frac{\partial s^p}{\partial x^i} - F_i^p(x, s(x)).$$

There is a canonical isomorphism $i_Y : V(J^1 Y \rightarrow M) \rightarrow J^1(VY \rightarrow M)$, [13], p. 255. If we compose the vertical tangent map $V\Gamma : VY \rightarrow VJ^1 Y$ with i_Y , we obtain a connection $\mathcal{V}\Gamma := i_Y \circ V\Gamma$ on $VY \rightarrow M$, which is called the vertical prolongation of Γ . If Y^p are the additional coordinates on VY , then the equations of $\mathcal{V}\Gamma$ are (3) and

$$(5) \quad dY^p = \frac{\partial F_i^p}{\partial y^q} Y^q dx^i.$$

Let Λ be a linear connection on TM and Λ^* be the dual connection on T^*M . Since $\mathcal{V}\Gamma$ is semilinear, we can construct the tensor product $\mathcal{V}\Gamma \otimes \Lambda^*$, which is a connection on $VY \otimes T^*M$, [1], [13]. If Y_i^p are the tensor coordinates on $VY \otimes T^*M$ and $\Lambda_{jk}^i(x)$ are the Christoffel symbols of Λ , then the coordinate expression of $\mathcal{V}\Gamma \otimes \Lambda^*$ is (3) and

$$(6) \quad dY_i^p = \left(\frac{\partial F_j^p}{\partial y^q} Y_i^q - \Lambda_{ij}^k Y_k^p \right) dx^j.$$

Hence we can construct the second order absolute differential

$$(7) \quad \nabla_{\Gamma, \Lambda}^2 s = \nabla_{\mathcal{V}\Gamma \otimes \Lambda^*} (\nabla_{\Gamma} s).$$

If we have a vector bundle $\pi : E \rightarrow M$, then $\mathcal{V}E \approx E \times_M E$. If Γ is a linear connection on E , then $\mathcal{V}\Gamma$ coincides with the product $\Gamma \times \Gamma$. For every section $s : M \rightarrow E$, we have $pr_2 \circ \nabla_{\Gamma} s : M \rightarrow E \otimes T^*M$ and $\nabla_{\Gamma, \Lambda}^2 s$ is identified with $(\nabla_{\Gamma} s, \nabla_{\Gamma \otimes \Lambda^*} (\nabla_{\Gamma} s))$. This is the classical tensor approach, [16]. In particular, if E is a tensor power of TM and T^*M and Γ is the corresponding tensor power of a linear connection Λ on TM and of its dual Λ^* , we take Λ again for the auxiliary linear connection on TM . Then we obtain the classical procedures of tensor calculus.

2. The groupoid approach. Let $\pi : Y \rightarrow M$ be a locally trivial fibered manifold. We write $IsoC^\infty(Y_x, Y_y)$ for the set of all diffeomorphisms of Y_x into Y_y .

Definition 1. The set

$$(8) \quad \mathcal{G}Y = \bigcup_{(x,y) \in M \times M} IsoC^\infty(Y_x, Y_y)$$

is called the groupoid of all diffeomorphisms of the fibers of Y or the groupoid of Y .

We are going to show that $\mathcal{G}Y$ is a smooth space in the sense of Frölicher.

In general, let $p_1 : Y_1 \rightarrow M_1$ and $p_2 : Y_2 \rightarrow M_2$ be two locally trivial fibered manifolds. Then we define

$$(9) \quad \mathcal{F}ib(Y_1, Y_2) = \bigcup_{(x,y) \in M_1 \times M_2} C^\infty(Y_{1x}, Y_{2y})$$

We denote by $p : \mathcal{F}ib(Y_1, Y_2) \rightarrow M_1 \times M_2$ the canonical projection. Consider the product projections $pr_1 : M_1 \times M_2 \rightarrow M_1$, $pr_2 : M_1 \times M_2 \rightarrow M_2$ and construct the pullbacks $\bar{Y}_1 = pr_1^* Y_1$, $\bar{Y}_2 = pr_2^* Y_2$, which are fibered manifolds over $M_1 \times M_2$. Then we have defined $\mathcal{F}(\bar{Y}_1, \bar{Y}_2)$ in the sense of (1). By the definition of pullback,

$$(10) \quad \mathcal{F}ib(Y_1, Y_2) = \mathcal{F}(\bar{Y}_1, \bar{Y}_2).$$

This introduces the structure of a smooth space on $\mathcal{F}ib(Y_1, Y_2)$, [2]. In other words, for every manifold N a map $f : N \rightarrow \mathcal{F}ib(Y_1, Y_2)$ is smooth, if the base map $p \circ f : N \rightarrow M_1 \times M_2$ is of class C^∞ and the induced map

$$(11) \quad \tilde{f} : f_1^* Y_1 \rightarrow Y_2,$$

$\tilde{f}(u, y_1) = f(u)(y_1)$, $f_1 = pr_1 \circ p \circ f$, $u \in N$, $y_1 \in Y_1$, $f_1(u) = p_1(y_1)$, is of class C^∞ . Moreover, τ -jets of N into $\mathcal{F}ib(Y_1, Y_2)$ are introduced by

$$J^\alpha(N, \mathcal{F}ib(Y_1, Y_2)) := J^\alpha(N, \mathcal{F}(\bar{Y}_1, \bar{Y}_2)),$$

where the right-hand side was defined in [4]. In the case of product bundles $Y_1 = M_1 \times Q_1$, $Y_2 = M_2 \times Q_2$, we have

$$(12) \quad J^\alpha(N, \mathcal{F}ib(Y_1, Y_2)) = J^\alpha(N, M_1 \times M_2) \times_N C^\infty(Q_1, J^\alpha(N, Q_2)),$$

where the subscript α indicates that we consider the maps into the fibers of the jet projection $\alpha : J^\alpha(N, Q_2) \rightarrow N$.

The inclusion $\mathcal{G}Y \subset \mathcal{F}ib(Y, Y)$ defines the structure of smooth space on $\mathcal{G}Y$. We shall write $a = pr_1 \circ p : \mathcal{G}Y \rightarrow M$, $b = pr_2 \circ p : \mathcal{G}Y \rightarrow M$. The following definition extends an idea by Ehresmann, [6], to the infinite dimensional space $\mathcal{G}Y$.

Definition 2. An element of connection on $\mathcal{G}Y$ at $x \in M$ is a 1-jet at x of a smooth map $\sigma : U \rightarrow \mathcal{G}Y$ of a neighbourhood U of x satisfying

$$(13) \quad a\sigma(u) = x, \quad b\sigma(u) = u \quad \text{for all } u \in U \quad \text{and} \quad \sigma(x) = \text{id}_{Y_x}.$$

The set of all elements of connection on $\mathcal{G}Y$ will be denoted by $Q\mathcal{G}Y$. Since $Q\mathcal{G}Y \subset J^1(M, \mathcal{G}Y)$, it is a smooth space as well. The source jet map is a projection $Q\mathcal{G}Y \rightarrow M$.

Every $A \in Q_x\mathcal{G}Y$, $A = j_x^1\sigma(u)$, defines a section $\tilde{A} : Y_x \rightarrow J_x^1Y$, $\tilde{A}(y) = j_x^1\sigma(u)(y)$. Conversely, Proposition 5 of [19] implies that for every section $B : Y_x \rightarrow J_x^1Y$ there exists a neighbourhood U of $x \in M$ and a map $\sigma : U \rightarrow \text{Fib}(Y, Y)$ satisfying (13) such that $B = j_x^1\sigma$. Since $\sigma(x) = \text{id}_{Y_x}$, the map $\sigma : U \times Y_x \rightarrow Y$ is local diffeomorphism in a neighbourhood of $\{x\} \times Y_x$. In this local sense, connections on Y correspond to smooth sections $\Gamma : M \rightarrow Q\mathcal{G}Y$.

Given a section $s : M \rightarrow Y$ and an element of connection $A = j_x^1\sigma(u) \in Q_x\mathcal{G}Y$, we define the absolute differential $\nabla_A s$ by

$$(14) \quad \nabla_A s = j_x^1(\sigma^{-1}(u)(s(u))) \in J_x^1(M, Y_x),$$

where $\sigma^{-1}(u)$ denotes the inverse diffeomorphism, so that $\sigma^{-1}(u)(s(u))$ is a local map $M \rightarrow Y_x$. We are going to show that (14) coincides with (2) for $\tilde{A} = \Gamma|_{Y_x}$. In some local coordinates x^i, y^p , let $\tilde{y}^p = f^p(u, y)$ be the coordinate expression of $\sigma(u)$. Then $\tilde{A} : Y_x \rightarrow J_x^1Y$ is given by

$$(15) \quad F_i^p(x, y) = \frac{\partial f^p(x, y)}{\partial x^i}.$$

If $y^p = s^p(u)$ is the coordinate form of s , $\sigma^{-1}(u)(s(u))$ is expressed by $\tilde{f}^p(u, s(u))$, where $\tilde{f}^p(u, y)$ is the inverse diffeomorphism of $\sigma(u)$. Then the coordinate form of $j_x^1\sigma^{-1}(u)(s(u))$ is

$$(16) \quad \frac{\partial \tilde{f}^p(x, y)}{\partial x^i} + \frac{\partial \tilde{f}^p(x, y)}{\partial y^q} \frac{\partial s^q}{\partial x^i}.$$

But $\sigma(x) = \text{id}_{Y_x}$ implies $\partial \tilde{f}^p(x, y)/\partial y^q = \delta_q^p$. Differentiating $\sigma^{-1}(u) \circ \sigma(u) = \text{id}_{Y_x}$, we obtain $\partial \tilde{f}^p(x, y)/\partial x^i = -\partial f^p(x, y)/\partial x^i = -F_i^p(x, y)$. Hence (16) coincides with (4).

Using this point of view, we interpret $\nabla_{\Gamma} s$ as a section of the union

$$(17) \quad J^1(M, Y, \pi) := \bigcup_{x \in M} J^1(M, Y_x),$$

which is a fibered manifold over M . Every diffeomorphism $\varphi : Y_x \rightarrow Y_y$ is extended into a map $J^1(\text{id}_M, \varphi) : J^1(M, Y_x) \rightarrow J^1(M, Y_y)$. This defines an injection $\mathcal{G}Y \hookrightarrow \mathcal{G}(J^1(M, Y, \pi))$. Hence every element of connection $A = j_x^1\sigma(u)$ on $\mathcal{G}Y$ is extended into an element of connection A_1 on $\mathcal{G}(J^1(M, Y, \pi))$ defined by

$$(18) \quad A_1 = j_x^1 J^1(\text{id}_M, \sigma(u)).$$

The correctness of this definition follows from the coordinate expressions. The local coordinates x^i, y^p on Y induce jet coordinates v^i, y^p, y_i^p on each $J^1(M, Y_x)$. If $\sigma(u)$ is expressed by

$$(19) \quad \bar{x}^i = u^i, \quad \bar{y}^p = f^p(u, y),$$

then the additional coordinate expression of $J^1(\text{id}_M, \sigma(u))$ is

$$(20) \quad \bar{v}^i = v^i, \quad \bar{y}_i^p = \frac{\partial f^p(u, y)}{\partial y^q} y_i^q.$$

Hence A_1 is of the form

$$(21) \quad dv^i = 0, \quad dy^p = F_i^p(x, y) dx^i, \quad dy_i^p = \frac{\partial F_j^p}{\partial y^q} y_i^q dx^j.$$

Thus, every connection Γ on Y is canonically extended into a connection Γ_1 on $J^1(M, Y, \pi)$. Since $\nabla_{\Gamma} s$ is a section of $J^1(M, Y, \pi)$, every

$$\nabla_{\Gamma_1} (\nabla_{\Gamma} s)(x) = j_x^1 J^1(\text{id}_M, \sigma(u))^{-1} (\nabla_{\Gamma} s(u))$$

is a semiholonomic 2-jet of M into Y_x .

Definition 3. The map

$$(22) \quad \nabla_{\Gamma}^2 s := \nabla_{\Gamma_1} (\nabla_{\Gamma} s) : M \rightarrow \bigcup_{x \in M} J_x^2(M, Y_x)$$

is called the second absolute differential of s with respect to Γ .

Proposition 1. *The coordinate form of $\nabla_{\Gamma}^2 s$ is (4) and*

$$(23) \quad \frac{\partial^2 s}{\partial x^i \partial x^j} - \frac{\partial F_i^p}{\partial x^j} - \frac{\partial F_i^p}{\partial y^q} \frac{\partial s^q}{\partial x^i} - \frac{\partial F_j^p}{\partial y^q} \frac{\partial s^q}{\partial x^i} + \frac{\partial F_j^p}{\partial y^q} F_i^q.$$

Proof. This follows directly from (4) and (21).

We remark that the idea of extending the groupoid $\mathcal{G}Y$ can be applied for prolongating connections in many similar cases. For example, every diffeomorphism $\varphi : Y_x \rightarrow Y_y$ induces the tangent map $T\varphi : V_x Y \rightarrow V_y Y$. Hence every element of connection $A = j_x^1 \sigma(u)$ on $\mathcal{G}Y$ defines an element of connection $\mathcal{V}A = j_x^1 T(\sigma(u))$ on the groupoid $\mathcal{G}VY$ of the vertical tangent bundle. For a connection Γ on Y , $\mathcal{V}\Gamma$ coincides with the vertical prolongation from Section 1.

3. The use of second tangent bundles. For every vector bundle $p : E \rightarrow M$, there are two vector bundle structures $\pi_E : TE \rightarrow E$ and $Tp : TE \rightarrow TM$ on TE . Moreover, we have an injection $i : E \rightarrow TE$ which identifies E_x with the tangent space $T_{0_x}(E_x)$ of the fiber E_x at its zero vector 0_x . In other words, $i(E)$ is the common kernel of both projection π_E and Tp . Using the terminology of J. Pradines, [18], [15], we say that $i(E) =: HE \subset TE$ is the *heart of E*. Clearly, if $q : D \rightarrow N$ is another vector bundle and $f : E \rightarrow D$ is a linear morphism, then Tf is a linear morphism of both vector bundle structures $\pi_E \rightarrow \pi_D$ and $Tp \rightarrow Tq$. We shall also say that Tf is linear in both directions. Moreover, $Tf(HE) \subset HD$ and the restriction $Hf : HE \rightarrow HD$ of Tf coincides with f .

Every non-holonomic 2-jet $X \in \tilde{J}_x^2(M, N)_y$ is of the form $j_x^1 \sigma$, where $\sigma : M \rightarrow J^1(M, N)$ is a section of the source projection $\alpha : J^1(M, N) \rightarrow M$, [5]. Every $\sigma(u) \in J_u^1(M, N)$ is identified with a linear map $\mu(\sigma(u)) : T_u M \rightarrow TN$, so that X defines a map

$$(24) \quad \mu X : TT_x M \rightarrow TT_y N, \quad \mu X = T_x \mu(\sigma(u)).$$

Consider the projections $\pi_{TM} : TTM \rightarrow TM$ and $T\pi_M : TTM \rightarrow TM$.

Lemma 1. (J. Pradines, [18]) *A map $A : TT_x M \rightarrow TT_y N$ represents a non-holonomic 2-jet $X \in \tilde{J}_x^2(M, N)_y$, i.e. $A = \mu X$, iff all following conditions are fulfilled:*

- (i) *A is π_T -projectable over a linear map $A_1 : T_x M \rightarrow T_y N$ and $T\pi$ -projectable over a linear map $A_2 : T_x M \rightarrow T_y N$,*
- (ii) *A is a linear morphism with respect to both vector bundle structures π_T and $T\pi$,*
- (iii) *the heart restriction $A_0 : H_x TM \rightarrow H_y TN$ coincides with A_1 .*

Moreover, X is semiholonomic, iff $A_1 = A_2$.

Proof. If $f^p(u)$, $f_i^p(u)$ is the coordinate expression of σ , then $\mu\sigma(u)$ is of the form $y^p = f^p(u)$, $Y^p = f_i^p(u)X^i$. For $T_x \mu\sigma(u)$ we find

$$(25) \quad Y^p = f_i^p(x)X^i, \quad dy^p = \frac{\partial f^p(x)}{\partial x^i} dx^i, \quad dY^p = \frac{\partial f_i^p(x)}{\partial x^j} X^i dx^j + f_i^p(x) dX^i.$$

This is the coordinate form of our claim. □

The absolute differentiation of sections of a fibered manifold $\pi : Y \rightarrow M$ can be extended to any map $f : N \rightarrow Y$. We define

$$(26) \quad \nabla_\Gamma f = v_\Gamma \circ Tf,$$

so that $\nabla_\Gamma f$ is a \mathcal{VB} -morphism $TN \rightarrow VY$ over $f : N \rightarrow Y$. If u^s are some local coordinates on N , U^s are the additional coordinates on TN ,

$$(27) \quad x^i = f^i(u), \quad y^p = f^p(u)$$

is the coordinate expression of f and Γ is given by (3), then the coordinate form of $\nabla_{\Gamma}f$ is (27) and

$$(28) \quad Y^p = \left(\frac{\partial f^p}{\partial u^s} - F_i^p(f^j(u), f^a(u)) \frac{\partial f^i}{\partial u^s} \right) U^s.$$

$\nabla_{\Gamma}f$ is a map with values in VY , so that we can construct its absolute differential with respect to any connection Δ on $VY \rightarrow M$. Since $\varrho : VY \rightarrow Y$ is a vector bundle, J^1VY is a vector bundle over J^1Y . A connection $\Delta : VY \rightarrow J^1VY$ is called semilinear, if it is projectable, i.e. there exists a connection $\Delta_0 : Y \rightarrow J^1Y$ satisfying $\Delta_0 \circ \varrho = (J^1\varrho) \circ \Delta$, and Δ is a \mathcal{VB} -morphism $VY \rightarrow J^1VY$ over Δ_0 . Clearly, both $T\varrho : TVY \rightarrow TY$ and $V\varrho : VVY \rightarrow VY$ are vector bundles. If Δ is a semilinear connection, its vertical projection $v_{\Delta} : TVY \rightarrow VVY$ is a \mathcal{VB} -morphism $T\varrho \rightarrow V\varrho$ over $v_{\Delta_0} : TY \rightarrow VY$.

Proposition 2. *Let Γ be a connection on $\pi : Y \rightarrow M$, Δ be a semilinear connection on $VY \rightarrow M$ and $f : N \rightarrow Y$ be a map. Then*

$$(29) \quad \nabla_{\Delta}(\nabla_{\Gamma}f)(u) : TT_uN \rightarrow VV_{f(u)}Y$$

corresponds to a non-holonomic 2-jet of $\tilde{J}_u^2(N, Y_{\pi(f(u))})$. If $\Delta_0 = \Gamma$, then each jet (29) is semiholonomic.

Proof. Since $\nabla_{\Gamma}f : TN \rightarrow VY$ is a linear morphism, $T\nabla_{\Gamma}f : TTN \rightarrow TVY$ is linear in both directions. Since Δ is semilinear, its vertical projection v_{Δ} is linear in both directions. Hence $v_{\Delta} \circ T\nabla_{\Gamma}f$ is linear in both directions over $v_{\Delta_0} \circ Tf$ and $v_{\Gamma} \circ Tf$. The heart map is $v_{\Gamma} \circ Tf$. Then our claim follows from Lemma 1. \square

Proposition 3. *If we take $\Delta = \mathcal{V}\Gamma$, then for every section $s : M \rightarrow Y$ we have*

$$(30) \quad \nabla_{\mathcal{V}\Gamma}\nabla_{\Gamma}s(x) = \mu(\nabla_{\Gamma}^2s(x)).$$

Proof. By (28), the coordinate form of $\nabla_{\Gamma}s$ is $y^p = s^p(x)$ and

$$(31) \quad Y^p = \left(\frac{\partial f^p}{\partial x^i} - F_i^p(x, s(x)) \right) X^i.$$

Using (5), we find $\nabla_{\mathcal{V}\Gamma}\nabla_{\Gamma}s$ in the form corresponding to (23). \square

4. Remarks. The groupoid approach to connections was invented by C. Ehresmann for Lie groupoids, which correspond to the classical principal fiber bundles, [6]. Every principal fiber bundle $\pi : P \rightarrow M$ with structure group G determines the associated groupoid PP^{-1} which can be defined as the factor space $P \times P / \sim$ with respect to the equivalence relation $(u, v) \sim (ug, vg)$, $u, v \in P$, $g \in G$. Writing uv^{-1} for such an equivalence class, we have two projections $a, b : PP^{-1} \rightarrow M$, $a(uv^{-1}) = \pi v$, $b(uv^{-1}) = \pi u$. The formula

$$(uv^{-1})(vw^{-1}) = uw^{-1}$$

defines a partial composition law in PP^{-1} and $e_x = uu^{-1}$ is its unit for every $x = \pi u \in M$. By definition, a Lie groupoid Φ over M is isomorphic to PP^{-1} for a principal bundle $P \rightarrow M$. If E is a fiber bundle associated with P with standard fiber S , every $v \in P_x$ determines the "frame map" $q_v : S \rightarrow E_x$, [13]. Then $q_u \circ q_v^{-1} : E_x \rightarrow E_y$, $u \in P_y$, depends on uv^{-1} only. This defines a map $PP^{-1} \rightarrow \mathcal{G}E$, which is called the action of PP^{-1} on E .

An element of connection on a Lie groupoid Φ at $x \in M$ is 1-jet of a local map $\sigma : U \rightarrow \Phi$ of a neighbourhood U of $x \in M$ satisfying $a\sigma(u) = x$, $b\sigma(u) = u$, $s(x) = e_x$, [6]. The space of all elements of connection on Φ is a fibered manifold $Q\Phi \rightarrow M$. A connection on Φ is a section $\Gamma : M \rightarrow Q\Phi$. If Φ acts on a fibered manifold $E \rightarrow M$, Γ induces a connection $\Gamma_E : E \rightarrow J^1E$, $\Gamma_E(y) = j_x^1\sigma(u)(y)$, provided $\Gamma(x) = j_x^1\sigma(u)$. In particular, $\Phi = PP^{-1}$ acts canonically on P and the connection Γ_P is principal. One verifies easily that the rule $\Gamma \rightarrow \Gamma_P$ establishes a bijection between connections on PP^{-1} and principal connections on P . For Φ acting on E , the absolute differentiation of sections of E with respect to a connection on Φ was introduced by Ehresmann, [6]. The principal bundle form of this operation was studied in [10]. Section 2 of the present paper represents a generalization of these ideas to the infinite dimensional groupoid $\mathcal{G}Y$.

We have already remarked in Section 1 that the first approach to the iterated absolute differentiation is related with the classical ideas of tensor calculus. On the other hand, the second approach is of different geometric character and its interesting applications can be found, e.g. in the theory of submanifolds of a space with Cartan connection, [9]. The connection in question determines the geometry of every submanifold N and the use of the contact elements generated by N , [13] (which are called jets of the submanifold N by some authors), eliminates any role of a linear connection on N . For example, the higher order torsions of N can be introduced in the framework of the second approach, [9].

5. Maps to the functional bundle. Consider two locally trivial fibered manifolds $p_1 : Y_1 \rightarrow M$, $p_2 : Y_2 \rightarrow M$ and the functional bundle (1). Write $p : \mathcal{F}(Y_1, Y_2) \rightarrow M$ for the canonical projection. The set $\mathcal{F}(Y_1, Y_2)$ is a smooth space in the sense of Frölicher, [2]. A connection Γ on $\mathcal{F}(Y_1, Y_2)$ is a smooth section $\Gamma : \mathcal{F}(Y_1, Y_2) \rightarrow J^1\mathcal{F}(Y_1, Y_2)$, [2]. For every smooth map $f : N \rightarrow \mathcal{F}(Y_1, Y_2)$, we can construct the tangent map $Tf : TN \rightarrow T\mathcal{F}(Y_1, Y_2)$. Using the vertical projection $v_\Gamma : T\mathcal{F}(Y_1, Y_2) \rightarrow V\mathcal{F}(Y_1, Y_2)$ of Γ , we define the absolute differential

$$(32) \quad \nabla_\Gamma f = v_\Gamma \circ Tf : TN \rightarrow V\mathcal{F}(Y_1, Y_2).$$

By linearity, (32) can be considered as a map $N \rightarrow V\mathcal{F}(Y_1, Y_2) \otimes T^*N$.

We remark that the construction of $\nabla_\Gamma f$ can be reduced to the absolute differentiation of a section of an induced bundle with respect to the induced connection analogously to the classical case of fibered manifolds. In general, if $g : N \rightarrow M$ is a map, we construct the induced bundles g^*Y_i , $i = 1, 2$,

$$g^*Y_i = \{(u, y_i) \in N \times Y_i, g(u) = p_i(y_i)\}$$

and define $g^*\mathcal{F}(Y_1, Y_2) = \mathcal{F}(g^*Y_1, g^*Y_2)$, which is a smooth space over N . If $s : M \rightarrow \mathcal{F}(Y_1, Y_2)$ is a smooth section, the formula $(g^*s)(u) = s(g(u))$ defines the induced

section $g^*s : N \rightarrow g^*\mathcal{F}(Y_1, Y_2)$. The rule $j_x^1s \mapsto j_u^1g^*s$ defines a map $J_x^1\mathcal{F}(Y_1, Y_2) \rightarrow J_u^1(g^*\mathcal{F}(Y_1, Y_2))$, $x = g(u)$. In this way, every connection $\Gamma : \mathcal{F}(Y_1, Y_2) \rightarrow J^1\mathcal{F}(Y_1, Y_2)$ induces a connection $g^*\Gamma : g^*\mathcal{F}(Y_1, Y_2) \rightarrow J^1g^*\mathcal{F}(Y_1, Y_2)$. Every smooth map $f : N \rightarrow \mathcal{F}(Y_1, Y_2)$ with $g = p \circ f$ defines a section $g^*f : N \rightarrow g^*\mathcal{F}(Y_1, Y_2)$. Then we have an identification

$$(33) \quad \nabla_\Gamma f \approx \nabla_{g^*\Gamma} g^*f.$$

Since $\Gamma : \mathcal{F}(Y_1, Y_2) \rightarrow J^1\mathcal{F}(Y_1, Y_2)$ is a kind of differential operator, one can characterize an r -th order connection, $r \geq 1$, [2]. We recall that every $X \in J_x^1\mathcal{F}(Y_1, Y_2)_\varphi$ is identified with an affine bundle morphism $\tilde{X} : J_x^1Y_1 \rightarrow J_x^1Y_2$ over $\varphi : Y_{1x} \rightarrow Y_{2x}$, whose derived linear morphism is $T\psi \otimes \text{id}_{T^*M}$. We say that Γ is of order r , if the condition $j_y^r\varphi = j_y^r\psi$, $\varphi, \psi \in C^\infty(Y_{1x}, Y_{2x})$, $y \in Y_{1x}$ implies

$$(34) \quad \tilde{\Gamma}(\varphi)|(J^1Y_1)_y = \tilde{\Gamma}(\psi)|(J^1Y_1)_y,$$

i.e. the restriction of the associated maps $\tilde{\Gamma}(\varphi), \tilde{\Gamma}(\psi) : J_x^1Y_1 \rightarrow J_x^1Y_2$ to the fiber $(J^1Y_1)_y$ over y coincide.

Write $\mathcal{F}\mathcal{J}^r(Y_1, Y_2) = \bigcup_{x \in M} J^r(Y_{1x}, Y_{2x})$, which is a finite dimensional manifold.

If x^i, y^p or x^i, z^α are some local fiber coordinates on Y_1 or Y_2 , respectively, then the induced coordinates on $\mathcal{F}\mathcal{J}^r(Y_1, Y_2)$ are x^i, y^p, z^α , where α is a multiindex of the range equal to the range of y^p with $0 \leq |\alpha| \leq r$. Let $S(J^1Y_1, J^1Y_2)$ be the space of all affine maps $(J^1Y_1)_y \rightarrow (J^1Y_2)_x$ with the derived linear map of the form $B \otimes \text{id}_{T_x^*M}$, $B \in V_x Y_2 \otimes V_y^* Y_1$. An r -th order connection Γ determines the associated map $\mathcal{G} : \mathcal{F}\mathcal{J}^r(Y_1, Y_2) \rightarrow S(J^1Y_1, J^1Y_2)$ by (34). Its coordinate form is

$$(35) \quad z^\alpha = z_p^\alpha y_i^p + \Phi_i^\alpha(x^i, y^p, z_\alpha^\alpha), \quad 0 \leq |\alpha| \leq r.$$

We say that Φ_i^α is the coordinate expression of Γ . Analogously to [2], if $x^i = f^i(u)$, $z^\alpha = f^\alpha(u, y)$ is the coordinate form of $f : N \rightarrow \mathcal{F}(Y_1, Y_2)$, then the coordinate expression of $\nabla_\Gamma f$ is

$$(36) \quad \left(\frac{\partial f^\alpha(u, y)}{\partial u^s} - \Phi_i^\alpha(x^i(u), y^p, \partial_\alpha f^\alpha(u, y)) \frac{\partial f^i}{\partial u^s} \right) U^s.$$

6. The second order procedure. In the remaining two sections we assume Γ is a finite order connection. Its vertical prolongation $\mathcal{V}\Gamma : V\mathcal{F}(Y_1, Y_2) \rightarrow J^1V\mathcal{F}(Y_1, Y_2)$ is a semilinear connection, [4]. Thus, for every map $F : N \rightarrow \mathcal{F}(Y_1, Y_2)$ we construct the iterated absolute differential

$$(37) \quad \nabla_{\mathcal{V}\Gamma}(\nabla_\Gamma f) : TTN \rightarrow VV\mathcal{F}(Y_1, Y_2).$$

We are going to deduce that the value of (37) at each $u \in N$ corresponds to a semiholonomic 2-jet of N into $C^\infty(Y_{1x}, Y_{2x})$, $x = p(f(u))$.

The non-holonomic and semiholonomic 2-jets of N into any functional bundle $\mathcal{F}(Y_1, Y_2)$ ($C^\infty(Y_{1x}, Y_{2x})$ is the case of one-point base) can be introduced as a special case of the iterated 2-jets studied in [4]. In particular, for the product bundles $Y_1 = M \times Q_1, Y_2 = M \times Q_2$, we have $\mathcal{F}(Y_1, Y_2) = M \times C^\infty(Q_1, Q_2)$ and Section 6 of [4] gives the following identifications

$$(38) \quad \tilde{J}^2(N, M \times C^\infty(Q_1, Q_2)) = \tilde{J}^2(N, M) \times_N C^\infty_\alpha(Q_1, \tilde{J}^2(N, Q_2)),$$

$$(39) \quad \bar{J}^2(N, M \times C^\infty(Q_1, Q_2)) = \bar{J}^2(N, M) \times_N C^\infty_\alpha(Q_1, \bar{J}^2(N, Q_2)),$$

where the subscript α indicates that we consider the maps into the fibers of the jet prolongation $\alpha : \tilde{J}^2(N, Q) \rightarrow N$ or $\alpha : \bar{J}^2(N, Q) \rightarrow N$. On the other hand, as a special case of Proposition 1 of [12], we obtain another trivialization formula

$$(40) \quad TT(M \times C^\infty(Q_1, Q_2)) = TTM \times C^\infty(Q_1, TTQ_2).$$

Every element $X \in \tilde{J}^2_u(N, \mathcal{F}(Y_1, Y_2))_\psi$ is of the form $X = j^1_u \sigma(v)$. Each $\sigma(v) \in J^1_v(N, \mathcal{F}(Y_1, Y_2))$ is identified with a linear map $\mu\sigma(v) : T_v N \rightarrow T\mathcal{F}(Y_1, Y_2)$, so that X defines a map

$$(41) \quad \mu X : TT_u N \rightarrow TT_\psi \mathcal{F}(Y_1, Y_2), \quad \mu X = T_u \mu(\sigma(v)).$$

Proposition 4. *For every $u \in N$, there exists a unique element*

$$\nabla^2_\Gamma f(u) \in \bar{J}^2_u(N, C^\infty(Y_{1x}, Y_{2x}))_{f(u)},$$

$x = p(f(u))$, satisfying

$$(42) \quad \nabla_{\nu\Gamma} \nabla_\Gamma f(u) = \mu(\nabla^2_\Gamma f(u)).$$

Proof. In the same way as in the proof of Proposition 2 we deduce that $\nabla_{\nu\Gamma} \nabla_\Gamma f$ satisfies the functional modification of Lemma 1 with the semiholonomicity condition. Using the trivializations (39) and (40), we can apply Lemma 1 pointwise. □

7. Relations to the curvature. For a finite order connection Γ with the coordinate expression Φ_i^α from (35), the additional coordinate expression of $\mathcal{V}\Gamma$ is

$$(43) \quad \frac{\partial \Psi_i^\alpha}{\partial z^b} Z^b + \dots + \frac{\partial \Phi_i^\alpha}{\partial z^\alpha} Z^\alpha$$

with $Z^\alpha_b = dz^\alpha_b$, [4]. If $z^\alpha = f^\alpha(x, y)$ is the coordinate expression of a section $s : M \rightarrow \mathcal{F}(Y_1, Y_2)$, then we obtain the coordinate form $\nabla_\Gamma s$ as a special case of (36)

$$(44) \quad \left(\frac{\partial f^\alpha(x, y)}{\partial x^i} - \Phi_i^\alpha(x^i, y^p, \partial_\alpha f^\alpha(x, y)) \right) X^i =: f_i^\alpha X^i.$$

Hence the coordinate form of the “second order term” in $\nabla_{\nu\Gamma} \nabla_\Gamma f$ is

$$(45) \quad \left(\frac{\partial}{\partial x^j} (f_i^\alpha) - \frac{\partial \Phi_j^\alpha}{\partial z^b} f_i^p - \dots - \frac{\partial \Phi_j^\alpha}{\partial z^\alpha} \partial_\alpha (f_i^p) \right) X^i \otimes dx^j$$

Analogously to the formula (36) of [4], we find that the alternation in i and j of (45) is $-(CT)(s(x))$, where CT is the curvature of Γ , [2].

We recall that every semiholonomic 2-jet $X \in \bar{J}^2_u(N, \mathcal{F}(Y_1, Y_2))_\psi$ determines the deviation $\Delta X \in T_\psi \mathcal{F}(Y_1, Y_2) \otimes \Lambda^2 T_u^* N$, whose coordinate expression is just the alternation of the “second order” component of X , [2]. Hence we have proved

Proposition 5. *For every finite order connection Γ on $\mathcal{F}(Y_1, Y_2)$ and every section $s : M \rightarrow \mathcal{F}(Y_1, Y_2)$, we have*

$$(46) \quad \Delta(\nabla_{\Gamma}^2 s(x)) = -C\Gamma(s(x)), \quad x \in M.$$

REFERENCES

- [1] Cabras A., *The Ricci identity for general connections*, Proc. Conf. Diff. Geom. and Applications, 1995, Masaryk University, Brno, 1996, 121–126.
- [2] Cabras A., Kolář I., *Connections on some functional bundles*, Czechoslovak Math. J., **45**(1995), 529–548.
- [3] Cabras A., Kolář I., *The universal connection of an arbitrary system*, to appear in Annali di Matematica.
- [4] Cabras A., Kolář I., *On the iterated absolute differentiation on some functional bundles*, Archivum Math. (Brno) **1-2** (1997), 23–35.
- [5] Ehresmann C., *Extension du calcul des jets aux jets non holonomes*, CRAS Paris, **239**(1954), 1762–1764.
- [6] Ehresmann C., *Sur les connexions d'ordre supérieur*, Atti del V° Cong. dell' Unione Mat. Italiana, 1955, Roma Cremonese 1956, 344–346.
- [7] Frölicher A., *Smooth structures*, Category theory 1981, LNM 962, Springer-Verlag, 1982, 69–81.
- [8] Jadczyk A., Modugno M., *Galilei general relativistic quantum mechanics*, to appear.
- [9] Kolář I., *Higher order torsions of manifolds with connection*, Archivum Math.(Brno), **8**(1972), 149–156.
- [10] Kolář I., *On the absolute differentiation of geometric object fields*, Ann. Polon. Math. **27**(1973), 293–304.
- [11] Kolář I., *Higher order absolute differentiation with respect to generalized connections*, Differential Geometry, Banach Center Publications **12**(1984), 153–162.
- [12] Kolář I., *Recent results and new ideas on Weil bundles*, Proc. Conference Geom. Topol., Editura MIRTON, Timisoara 1996, 127–136.
- [13] Kolář I., Michor P. W., Slovák J., *Natural Operations in Differential Geometry*, Springer-Verlag, 1993.
- [14] Kolář I., Modugno M., *The Frölicher-Nijenhuis bracket on some functional spaces*, to appear in Ann. Polon. Math.
- [15] Libermann P., *Introduction to the theory of semi-holonomic jets*, Archivum Math. (Brno) **3** (1997), 173–189.
- [16] Modugno M., *Systems of vector valued forms on a fibered manifold and applications to gauge theories*, Springer-Verlag, 1987, 238–264.
- [17] Pohl F. W., *Connections in differential geometry of higher order*, Trans. Amer. Math. Soc., **125**(1966), 310–325.
- [18] Pradines J., *Représentation des jets non holonomes par des morphismes vectoriels doubles soudés*, CRAS Paris, série A **278**, 1974, 1523–1526.
- [19] Slovák J., *Smooth structures on fibre jet spaces*, Czechoslovak Math. J., **36**(1986), 358–375.

Antonella Cabras

Dipartimento di Matematica Applicata “G. Sansone”

Via S. Marta 3

50139 Firenze, ITALY

Ivan Kolář

Department of Algebra and Geometry

Faculty of Science, Masaryk University

Janáčkovo nám. 2a

662 95 Brno, CZECH REPUBLIC