

Lubomír Klapka

Integrals and variational multipliers of second-order ordinary differential equations

In: Jan Slovák and Martin Čadek (eds.): Proceedings of the 20th Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 2001. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 66. pp. 129–131.

Persistent URL: <http://dml.cz/dmlcz/701673>

Terms of use:

© Circolo Matematico di Palermo, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

INTEGRALS AND VARIATIONAL MULTIPLIERS OF SECOND-ORDER ORDINARY DIFFERENTIAL EQUATIONS

LUBOMÍR KLAPKA

ABSTRACT. In this paper the relation between integrals and variational multipliers of a system of second-order ordinary differential equations is studied. A simple necessary and sufficient local condition for the existence of a multiplier is given.

1. INTRODUCTION

There are a few different approaches to the inverse problem of the variational calculus for second-order differential equations (see, e.g. [1, 2, 6, 7]), but all of them use the notion of variational multipliers. The purpose of this paper is to study the local relation between variational multipliers and integrals of these differential equations. An appropriate mathematical apparatus was developed in [3, 4]. For simplicity, all mappings which are used in this paper are supposed to be infinitely differentiable and all expressions which are used here are local expressions only.

2. INTEGRALS AND MULTIPLIERS

We say that a given system of differential equations of the form

$$\ddot{x}^k = f^k(t, x, \dot{x}), \quad (1)$$

where $k = 1, 2, \dots, n$, has an *integral* F if and only if $F(t, x, \dot{x})$ is a function such that $dF/dt = 0$ on any solution of (1). In this case, $dF/dt \equiv (\partial F/\partial \dot{x}^k)(\ddot{x}^k - f^k)$. Integrals are said to be *independent* if they are independent as functions.

We say that (1) has a *variational multiplier* if and only if there exists a non-singular matrix $(g_{kl}(t, x, \dot{x}))$, such that for some function $L(t, x, \dot{x})$

$$g_{kl}(\ddot{x}^l - f^l) \equiv \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{x}^k} \right) - \frac{\partial L}{\partial x^k}. \quad (2)$$

2000 *Mathematics Subject Classification*: 34A34, 34A55, 37J05, 37J35, 70H03, 70H15, 93C15.

Key words and phrases: Inverse problem of variational calculus, second-order ordinary differential equations, variational multipliers, integrals of differential equations, canonical transformations.

The paper is in final form and no version of it will be submitted elsewhere.

Theorem. A system of differential equations (1) has a variational multiplier if and only if it has independent integrals $P_1, P_2, \dots, P_n, Q^1, Q^2, \dots, Q^n$ satisfying conditions

$$\frac{\partial P_\alpha}{\partial \dot{x}^k} \frac{\partial Q^\alpha}{\partial \dot{x}^i} - \frac{\partial P_\alpha}{\partial \dot{x}^i} \frac{\partial Q^\alpha}{\partial \dot{x}^k} = 0. \quad (3)$$

Proof. Suppose that P_α, Q^α are independent integrals of (1) satisfying the condition (3). This condition implies the existence of a function $S(t, x, \dot{x})$, such that

$$\frac{\partial S}{\partial \dot{x}^k} = -P_\alpha \frac{\partial Q^\alpha}{\partial \dot{x}^k}. \quad (4)$$

Then the function $L \equiv P_\alpha(dQ^\alpha/dt) + dS/dt$ does not depend on \dot{x}^k and so by straightforward calculation we have (2), where

$$g_{kl} \equiv \frac{\partial P_\alpha}{\partial \dot{x}^l} \frac{\partial Q^\alpha}{\partial x^k} - \frac{\partial P_\alpha}{\partial x^k} \frac{\partial Q^\alpha}{\partial \dot{x}^l}. \quad (5)$$

From (3) and (5) we get $dP_1 \wedge \dots \wedge dP_n \wedge dQ^1 \wedge \dots \wedge dQ^n = (-\det |g_{kl}|)^n dx^1 \wedge \dots \wedge dx^n \wedge d\dot{x}^1 \wedge \dots \wedge d\dot{x}^n$. Since the integrals are independent, the matrix (g_{kl}) is non-singular. Hence, the system (1) has a variational multiplier.

Conversely, suppose that a system of differential equations (1) has a variational multiplier. Then from (2) one gets $\partial^2 L / \partial \dot{x}^k \partial \dot{x}^l = g_{kl}$. Since (g_{kl}) is a non-singular matrix, we can use the Hamiltonian formalism. Denote the Hamilton function by H . There exists a canonical transformation $(p_k, x^k, t) \rightarrow (P_\alpha, Q^\alpha, t)$ reducing H to zero (see, e.g. [4, 5]). In this case P_α, Q^α are independent integrals of motion and $p_k dx^k - P_\alpha dQ^\alpha - H dt = dS$, where S is a corresponding generating function. If we return to Lagrangean coordinates x^k, \dot{x}^k, t we get, among others, the formula (4). The integrability condition of (4) with respect to S gives (3). This completes the proof.

3. REMARK

The above Theorem is a direct consequence of Theorem 1. in [3], p. 3785, as well as Proposition 9.2.11 in [4], p. 180. However, they lead to another proof, based on the concept of a Lepagean 2-form.

ACKNOWLEDGMENTS

The author is indebted to Olga Krupková for here attention to this work. The support by Grants CEZ:J10/98:192400002 and VS 96003 "Global Analysis" of the Czech Ministry of Education, Youth and Sports, and Grant No. 201/00/0724 of the Czech Grant Agency is gratefully acknowledged.

REFERENCES

- [1] I. Anderson, G. Thompson, *The inverse problem of the calculus of variations for ordinary differential equations*, *Memoirs Amer. Math. Soc.* **98** (1992) No. 473.
- [2] J. Douglas, *Solution of the inverse problem of the calculus of variations*, *Trans. Amer. Math. Soc.* **50** (1941) 71-128.
- [3] L. Klapka, *Integrals of motion and semi-regular Lepagean forms in higher-order mechanics*, *J. Phys. A: Math. Gen.* **16** (1983) 3783-3794.
- [4] O. Krupková, *The Geometry of Ordinary Variational Equations*, *Lecture Notes in Mathematics* **1678** (Springer, Berlin, 1997).

- [5] A. Lichnerowicz, *Geometry of the canonical transformations*, In: Dynamical systems and micro-physics, control theory and mechanics, A. Blaquiere, G. Leitmann (eds.), Proc. 3rd Int. Semin. Udine, Italy, Sept. 4 - 9, 1983 (Academic Press Inc., San Diego, California 1984), 287-327.
- [6] Z. Muzsnay, *Sur le problème inverse du calcul des variations*, Thesis (Univ. Paul Sabatier, Toulouse, 1997).
- [7] W. Sarlet, *Integrability aspects of the inverse problem of the calculus of variations*, In: Differential Geometry and Applications, I. Kolář, O. Kowalski, D. Krupka, J. Slovák (eds.), Proc. Conf. Brno, Czech Republic, Aug. 10 - 14, 1998 (Masaryk University, Brno, 1999) 563-574.

SILESIA UNIVERSITY AT OPAVA, MATHEMATICAL INSTITUTE
BEZRUČOVO NÁM. 13, 746 01 OPAVA 1, CZECH REPUBLIC
E-mail: Lubomir.Klapka@math.slu.cz