

Assem Tharwat; Karel Zimmermann

Some optimization problems on solubility sets of separable Max-Min equations and inequalities


Acta Universitatis Carolinae. Mathematica et Physica, Vol. 38 (1997), No. 2, 45--57

Persistent URL: <http://dml.cz/dmlcz/702044>

Terms of use:

© Univerzita Karlova v Praze, 1997

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Some Optimization Problems on Solubility Sets of Separable Max-Min Equations and Inequalities

A. THARWAT and K. ZIMMERMANN

Praha*)

Received 10. January 1997

1 Introduction

The aim of this paper is to suggest a direct Parametric method for solving some optimization problems on attainable sets of so called max-separable operators. Such problems in a less general form connected with the fuzzy set theory were considered e.g. in [1], [4]. The problem considered in this paper is presented independently of the fuzzy sets context as a non-linear nonconvex optimization problem. Parametric approach to its solution suggested is flexible enough to allow further extension and generalization, which are briefly discussed in the concluding sections.

2 Notations and Formulation of the Basic Problem

In this paper, we shall consider the following system of equations and inequalities

$$R_i(x) \equiv \max_{j \in N} (a_{ij} \wedge r_{ij}(x_j)) = b_i, \quad \forall i: i \in S$$

$$h_j \leq x_j \leq H_j, \quad \forall j: j \in N \quad (1)$$

where $N \equiv \{1, 2, \dots, n\}$, $S \equiv \{1, 2, \dots, m\}$, $x \equiv (x_1, \dots, x_n) \in R^n$, $b \equiv (b_1, \dots, b_m) \in R^m$, $h \equiv (h_1, \dots, h_n) \in R^n$, $H \equiv (H_1, \dots, H_n) \in R^n$, $a_{ij} \wedge r_{ij}(x_j) \equiv \min(a_{ij}, r_{ij}(x_j))$, $R(x) = (R_1(x), \dots, R_m(x))$, let us assume further that $r_{ij}: R \rightarrow R$ are given strictly increasing continuous functions $\forall i: i \in S, \forall j: j \in N$. Using the above vector notation we can reformulate the system (1) as follows:

$$R(x) = b, \quad h \leq x \leq H \quad (2)$$

*) Department of Operation Research, Faculty of Mathematics and Physics, Charles University, 118 00 Praha 1, Malostranské nám. 25, Czech Republic

Denote the set of all solutions of the system (1) (or (2)) by $M(b)$. Each component of $R : E^n \rightarrow E^m$ is a function depending on n variables; this function is expressed as the maximum n nondecreasing functions of one variable of the form $a_{ij} \wedge r_{ij}(x_j)$, so that these functions are separated by a max-operation. By similarity with the additive separability, we call this property of the functions $R_i(x)$ max-separability and $R(x)$ is called a max-separable operator.

The vector b in the system (2) can be understood as a vector, which is attained by the left hand side $R(x)$ when an appropriate $x \in M(b)$ is chosen. Therefore those b 's, for which $M(b) \neq \emptyset$, are called attainable elements and the set

$$A \equiv \{b \mid M(b) \neq \emptyset\} \quad (3)$$

is called the attainable set.

If an element $\hat{b} \in A$, then there exists a solution of the system (2) with $b = \hat{b}$ which can be obtained using some of the methods described in the literature (see e.g. [2], [3]). If $\hat{b} \notin A$, we want to find an approximate solution of the system (2) with the right hand side \hat{b} . For this purpose, we look for an element $b^{opt} \in A$, which has in some sense the minimal distance from \hat{b} and accept the elements of $M(b^{opt})$ as appropriate approximate solutions.

In this article, we shall use the Tshebyshev distance, i.e. the following distance:

$$\|b - \hat{b}\| \equiv \max_{i \in S} |b_i - \hat{b}_i| \quad (4)$$

The problem, we are going to solve here is thus in the following form:

$$\|b - \hat{b}\| \rightarrow \min \quad \text{subject to} \quad b \in A \quad (5)$$

Since if $b \in A$, it means that there exists x such that $b = R(x)$ so that we can reformulate the problem (5) as follows:

$$\|R(x) - \hat{b}\| \equiv \max_{i \in S} |R_i(x) - \hat{b}_i| \rightarrow \min \quad \text{subject to} \quad h \leq x \leq H \quad (6)$$

The reformulation (6) shows that we minimize a continuous function of x on a compact set, so that there exists always at least one optimal solution x^{opt} of the system (6); thus if we set $b^{opt} \equiv R(x^{opt})$, we will obtain an optimal solution of the problem (5).

Let us define the set $M(t)$ for any $t : t \in [0, \infty)$ as follows:

$$M(t) \equiv \{x \mid h \leq x \leq H \ \& \ \|R(x) - \hat{b}\| \leq t\}. \quad (7)$$

The set $M(t)$ is nonempty if and only if the following system of inequalities:

$$R_i(x) \leq \hat{b}_i + t, \ i \in S \ \& \ R_i(x) \geq \hat{b}_i - t, \ i \in S \ \& \ h \leq x \leq H, \quad (8)$$

is soluble with respect to x ; note that the set $M(t)$ is the set of all solutions x of (8). We can replace our original problems (5), (6) by the following problem:

$$t \rightarrow \min \quad \text{subject to} \quad M(t) \neq \emptyset. \quad (9)$$

We shall show in the sequel that there exists always the optimal solution $t^{opt} \geq 0$ of the problem (9) and also we will derive a direct numerical procedure for determining t^{opt} . If x^{opt} is an any element of $M(t^{opt})$, then $\|R(x^{opt}) - \hat{b}\| \leq t^{opt}$; since the strict inequality can't hold, the equality must occur i.e. $b^{opt} \equiv R(x^{opt}) \in A$ is the optimal solution of the problem (5), and the vector x^{opt} can be accepted as an approximate solution of the system (2) in the case the $\hat{b} \notin A$, since for any solution x of the system (2), we have $\|R(x) - \hat{b}\| \geq t^{opt}$.

In the next section we investigate some properties of the set $M(t)$ where $t \in [0, \infty)$. This will enable us to derive the direct solution method for the system (5).

3 Properties of $M(t)$

We shall introduce the following notations $\forall i : i \in S, \forall j : j \in N, t \in [0, \infty)$:

$$V_{ij}(t) = \{x_j \mid h_j \leq x_j \leq H_j \quad \text{and} \quad a_{ij} \wedge r_{ij}(x_j) \leq \hat{b}_i + t\}$$

$$V_j(t) = \bigcap_{i \in S} V_{ij}(t) \quad \& \quad W_{ij}(t) = \{x_j \mid h_j \leq x_j \leq H_j \quad \text{and} \quad a_{ij} \wedge r_{ij}(x_j) \geq \hat{b}_i - t\}.$$

For the illustration of these sets see the Appendix. The following theorem gives the necessary and sufficient conditions for $M(t) \neq \emptyset$.

Theorem 3.1.

$$M(t) \neq \emptyset \Leftrightarrow \left\{ \begin{array}{l} 1) \quad V_j(t) \neq \emptyset, \quad \forall j \in N \\ 2) \quad \forall i \in S \exists j(i) \in N \text{ such that } W_{ij(i)}(t) \cap V_{j(i)}(t) \neq \emptyset. \end{array} \right\}$$

Proof. Define the interval $I_j = [h_j, H_j]$ and introduce the following notations:

$$a1_{ij} = a_{ij} - (\hat{b}_i + t), \quad a2_{ij} = a_{ij} - (\hat{b}_i - t) \quad \& \quad r1_{ij}(x_j) = r_{ij}(x_j) - (\hat{b}_i + t), \\ r2_{ij}(x_j) = r_{ij}(x_j) - (\hat{b}_i - t)$$

Sufficiency.

Assume that we have a point $x \equiv (x_1, \dots, x_n)$ which satisfies the right hand side of the \Leftrightarrow -relation.

$$\begin{aligned} x_j \in V_j(t), \quad \forall j \in N &\rightarrow x_j \in \bigcap_{i \in S} V_{ij}(t), \quad \forall j \in N \\ &\rightarrow (a1_{ij} \wedge r1_{ij}(x_j) \leq 0, \quad \forall i \in S; x_j \in I_j), \quad \forall j \in N \\ &\rightarrow \max_{j \in N} (a1_{ij} \wedge r1_{ij}(x_j) \leq 0), \quad \forall i \in S; \forall x_j \in I_j, \\ x_{j(i)} \in W_{ij(i)}(t) &\rightarrow a2_{ij(i)} \wedge r2_{ij(i)}(x_j) \geq 0; x_{j(i)} \in I_{j(i)} \\ &\rightarrow \max_{j \in N} (a2_{ij} \wedge r2_{ij}(x_j) \geq 0; x_{j(i)} \in I_{j(i)}). \end{aligned}$$

Then we can deduce that $x \in M(t)$.

Necessity.

$$\begin{aligned}
x \in M(t) &\rightarrow x_j \in I_j, \forall j \in N \ \& \ (\max_{j \in N} a_{1ij} \wedge r_{1ij}(x_j) \leq 0 \ \forall i \in S) \\
&\quad (\max_{j \in N} a_{2ij} \wedge r_{2ij}(x_j) \geq 0 \ \forall i \in S) \\
&\rightarrow (x_j \in I_j, \text{ and } a_{1ij} \wedge r_{1ij}(x_j) \leq 0 \ \forall i \in S), \forall j \in N \\
&\quad \& \ \forall i \in S \ \exists j(i) \in N \text{ such that } a_{2ij(i)} \wedge r_{2ij(i)}(x_j) \geq 0; x_{j(i)} \in I_{j(i)} \\
&\rightarrow 1) V_j(t) \neq \emptyset, \forall j \in N \\
&\quad 2) \forall i \in S \ \exists j(i) \in N \text{ such that } W_{ij(i)}(t) \cap V_{j(i)}(t) \neq \emptyset.
\end{aligned}$$

Thus the proof of the theorem is complete. \square

4 Properties of $V(t)$ & $W(t)$

We shall investigate here the conditions for $V_{ij}(t) \neq \emptyset$, $V_j(t) \neq \emptyset$, $W_{ij}(t) \neq \emptyset$. Define the following variables:

$$\begin{aligned}
\eta^{ij} &\equiv \min (\max \{a_{ij} - \hat{b}_i, 0\}, \max \{r_{ij}(h_j) - \hat{b}_i, 0\}), \quad \eta^j \equiv \max_{i \in S} \eta^{ij}, \\
\eta &\equiv \max_{j \in N} \eta^j \quad \text{and} \quad \tau^{ij} \equiv \max (0, \hat{b}_i - a_{ij}, \hat{b}_i - r_{ij}(H_j)). \quad (10)
\end{aligned}$$

For the illustration of these variables see the appendix.

Theorem 4.1. For each $j \in N$, $\exists \eta^{ij} \geq 0$ such that $V_{ij}(t) \neq \emptyset \Leftrightarrow t \geq \eta^{ij}$.

Proof.

$$\begin{aligned}
V_{ij}(t) = \emptyset &\Leftrightarrow a_{ij} \wedge r_{ij}(x_j) > \hat{b}_i + t; \quad \forall x_j \in I_j \\
&\Leftrightarrow \hat{b}_i + t < r_{ij}(x_j) < a_{ij} \text{ or } \hat{b}_i + t < a_{ij} < r_{ij}(x_j); \text{ it is further } t \geq 0 \\
&\Leftrightarrow t \min (\max \{r_{ij}(h_j) - \hat{b}_i, 0\}, \max \{a_{ij} - \hat{b}_i, 0\}) = \eta^{ij}
\end{aligned}$$

where η^{ij} is given by (1); this completes the proof of the theorem. \square

Theorem 4.2. For each $j \in N$, $V_j(t) \neq \emptyset \Leftrightarrow t \geq \eta^j$; where η^j is given by (10).

Proof. $V_j(t) = \emptyset$ equivalent to the fact $V_{i_0j}(t) = \emptyset$ for some $i_0 \in S$; since V_{i_0j} are nested¹⁾ sets for fixed $j_0 \in N$; which means that $t < \eta^{i_0j} \leq \eta^j$ which is the maximum of η^{ij} on S . \square

Corollary 4.1. For each $j \in N$, $V_j(t) \neq \emptyset \Leftrightarrow t \geq \eta$; where η is given by (10).

1) Since for each $j, 1 \leq j \leq n$, there exists a permutation $\{i, \dots, i_m\}$ such that $V_{i_1j} \subset V_{i_2j} \subset \dots \subset V_{i_mj}$ because of the fact that $a_{ij} \wedge r_{ij}(x_j)$ are nondecreasing in x_j .

Proof. The proof is obviously derived from theorem 4.2.

Corollary 4.2. $M(t) \neq \emptyset \Leftrightarrow t \geq \eta$.

Proof. The proof is obviously derived from theorem 3.1, theorem 4.2 and corollary 4.1, where η is given by (10).

Theorem 4.3. For each $i \in S, j \in N; \exists \tau^{ij}$ such that $W_{ij}(t) \neq \emptyset \Leftrightarrow t \geq \tau^{ij}$.

Proof.

$$\begin{aligned} W_{ij}(t) = \emptyset &\Leftrightarrow a_{ij} \wedge r_{ij}(x_j) < \hat{b}_i - t; \quad \forall x_j \in I_j \\ &\Leftrightarrow a_{ij} < r_{ij}(x_j) < \hat{b}_i - t \text{ or } r_{ij}(x_j) < a_{ij} < \hat{b}_i - t; \text{ it is further } t \geq 0 \\ &\Leftrightarrow t < -\min(0, a_{ij} - \hat{b}_i, r_{ij}(H_j) - \hat{b}_i) \\ &\Leftrightarrow t < \max(0, \hat{b}_i - a_{ij}, \hat{b}_i) r_{ij}(H_j) = \tau^{ij} \end{aligned}$$

where τ^{ij} is given by (1); this completes the proof of the theorem. \square

Corollary 4.3. For each $j \in N, i \in S; \exists \tau^{ij} \geq 0$ such that

$$\forall t : t < \max_{i \in S} \min_{j \in N} \max_{i \in S} (\eta, \tau^{ij}) \Rightarrow M(t) \neq \emptyset.$$

Proof. It is clear from corollary 4.1, 4.2 and theorem 3.1, where τ^{ij} is given by (10).

Let us define the following sets: $P_{ikj}(t) = W_{ij}(t) \cap V_{kj}(t); \forall i, k \in S, j \in N$. To investigate the necessary and sufficient conditions for $P_{ikj}(t) \neq \emptyset$, assume that the variable η_{ikj} satisfies the following equation $r_{kj}^{-1}(\hat{b}_k + \eta_{ikj}) = r_{ij}^{-1}(\hat{b}_i - \eta_{ikj})$ for some, $i, k \in S, j \in N$ and define the variables ξ_{ikj}, ζ_{ikj} and γ_{ikj} for some $i, k \in S$ and $j \in N$ as follows:

$$\xi_{ikj} = \begin{cases} \eta_{ikj} & \text{if } \max(\tau^{ij}, \eta^{ij}) < \eta_{ikj} < \min(a_{kj} - \hat{b}_k, \hat{b}_i - r_{ij}(h_j)) \\ a_{kj} - \hat{b}_k & \text{if } r_{kj}^{-1}(\hat{b}_k + t) \leq r_{ij}^{-1}(\hat{b}_i - t) < H_j \\ \max(\tau^{ij}, \eta^{ij}) & \text{otherwise} \end{cases} \quad (11)$$

$$\zeta_{ikj} = \begin{cases} \eta_{ikj} & \text{if } \max(\tau^{ij}, \eta^{ij}) < \eta_{ikj} < \min(r_{kj}(H_j) - \hat{b}_k, \hat{b}_i - r_{ij}(h_j)) \\ \max(\tau^{ij}, \eta^{ij}) & \text{otherwise} \end{cases} \quad (12)$$

$$\gamma_{ikj} = \max(\tau^{ij}, \eta^{ij}) \quad (13)$$

For the illustration of these variables see also the appendix.

Concerning the definition of the sets $V_{kj}(t)$ & $W_{ij}(t)$, if we assume that

$$r_{ij}(h_j) \leq a_{ij} \leq r_{ij}(H_j) \quad \text{for all } i \in S, j \in N$$

then it is easy to recognize the following remarks:

Remark 1.

If the two sets $[r_{kj}(h_j) - \hat{b}_k, a_{kj} - \hat{b}_k]$ & $[\hat{b}_i - a_{ij}, \hat{b}_i - r_{ij}(h_j)]$ have an empty intersection, then we can deduce the following:

- if $a_{kj} - \hat{b}_k < \hat{b}_i - a_{ij} = \tau^{ij} = \max(\tau^{ij}, \eta^{ij})$, for some $k \in S$, then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < \hat{b}_i - a_{ij}$; and if $t \geq \hat{b}_i - a_{ij}$ the intersection equals $W_{ij}(t)$.
- if $\hat{b}_i - r_{ij}(h_j) < r_{kj}(h_j) - \hat{b}_k = \eta^{ij} = \max(\tau^{ij}, \eta^{ij})$, for some $k \in S$, then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < r_{kj}(h_j) - \hat{b}_k$; and if $t \geq r_{kj}(h_j) - \hat{b}_k$ the intersection equals $V_{kj}(t)$.

Remark 2.

If the two sets $[r_{kj}(h_j) - \hat{b}_k, a_{kj} - \hat{b}_k]$ & $[\hat{b}_i - a_{ij}, \hat{b}_i - r_{ij}(h_j)]$ have a single point in their intersection, then we can deduce the following:

- if the point of intersection is $x = \hat{b}_i - r_{ij}(h_j) = r_{kj}(h_j) - \hat{b}_k = \eta^{ij} = \max(\tau^{ij}, \eta^{ij})$, for some $k \in S$, then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < x$; and if $t \geq x$ the intersection equals $V_{kj}(t)$.
- if the point of intersection is $x = a_{kj} - \hat{b}_k = \hat{b}_i - a_{ij}$, for some $k \in S$, then we have the following two cases:
 $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < x$, given that $r_{ij}^{-1}(\hat{b}_i - t) \leq r_{kj}^{-1}(\hat{b}_k + t) < H_j$; and if $t \geq x$ the intersection equals $W_{ij}(t)$; $x = a_{kj} - \hat{b}_k$.
 $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < x$, given that $r_{kj}^{-1}(\hat{b}_k + t) \leq r_{ij}^{-1}(\hat{b}_i - t) < H_j$; and if $t \geq x$ the intersection equals $V_{kj}(t)$; $x = \hat{b}_i - a_{ij} = \tau^{ij} = \max(\tau^{ij}, \eta^{ij})$.

Remark 3.

Let

$$\begin{aligned} z_1 &= r_{kj}(h_j) - \hat{b}_k, & y_1 &= a_{kj} - \hat{b}_k \\ z_2 &= \hat{b}_i - a_{ij}, & y_2 &= \hat{b}_i - r_{ij}(h_j). \end{aligned}$$

Assuming that $[z, y] = [z_1, y_1] \cap [z_2, y_2]$ one can find the following cases:

- if $[z, y] = [z_1, y_1]$, then there exists some t_0 such that

$$\tau^{ij} < \eta^{ij} = z_1 < t_0 < y_1 < y_2 \quad \text{and} \quad r_{kj}^{-1}(\hat{b}_k + t_0) = r_{ij}^{-1}(\hat{b}_i - t_0);$$

i.e. $\max(\tau^{ij}, \eta^{ij}) < t_0 < \min(y_1, y_2)$,

then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t \geq t_0$ the intersection is nonempty, (similarly the case $[z, y] = [z_2, y_2]$).

- if $[z, y] = [z_0, y_1]$, $z_0 = z_1 = z_2$ then there exists some t_0 such that

$$\tau^{ij} = \eta^{ij} = z_0 < t_0 < y_1 < y_2 \quad \text{and} \quad r_{kj}^{-1}(\hat{b}_k + t_0) = r_{ij}^{-1}(\hat{b}_i - t_0);$$

i.e. $\max(\tau^{ij}, \eta^{ij}) < t_0 < \min(y_1, y_2)$,

then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < t_0$; and if $t \geq t_0$ the intersection is nonempty, (similarly the case $[z, y] = [z_0, y_2]$).

- if $[z, y] = [z_1, y_0]$, $y_0 = y_1 = y_2$ then there exists some t_0 such that

$$\tau^{ij} < \eta^{ij} = z_1 < t_0 < y_0 \quad \text{and} \quad r_{kj}^{-1}(\hat{b}_k + t_0) = r_{ij}^{-1}(\hat{b}_i - t_0);$$

i.e. $\max(\tau^{ij}, \eta^{ij}) < t_0 < \min(y_1, y_2)$,

then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < t_0$; and if $t \geq t_0$ the intersection is nonempty, (similarly the case $[z, y] = [z_2, y_0]$).

- if $[z, y] = [z_1, y_2]$, then there exists some t_0 such that

$$\tau^{ij} < \eta^{ij} = z_1 < t_0 < y_2 < y_1 \quad \text{and} \quad r_{kj}^{-1}(\hat{b}_k + t_0) = r_{ij}^{-1}(\hat{b}_i - t_0);$$

i.e. $\max(\tau^{ij}, \eta^{ij}) < t_0 < \min(y_1, y_2)$,

then $V_{kj}(t) \cap W_{ij}(t) = \emptyset$ if $t < t_0$; and if $t \geq t_0$ the intersection is nonempty, (similarly the case $[z, y] = [z_2, y_1]$).

Theorem 4.4. *Let $i, k \in S, j \in N; r_{ij}(h_j) \leq a_{ij} \leq r_{ij}(H_j)$ for all $i \in S, j \in N$; then $\exists \xi_{ikj} \geq 0$ such that $P_{ikj}(t) = \emptyset \Leftrightarrow t \geq \xi_{ikj}$.*

Proof. The proof is obviously derived from the above remarks and from the definition of ξ_{ikj} which is given in (11).

Theorem 4.5. *Let $i, k \in S, j \in N; r_{ij}(h_j) \leq r_{ij}(H_j) \leq a_{ij}$ for all $i \in S, j \in N$; then $\exists \zeta_{ikj} \geq 0$ such that $P_{ijk}(t) \neq \emptyset \Leftrightarrow t \geq \zeta_{ikj}$.*

Proof. The proof is obviously derived from the above remarks and from the definition of ζ_{ikj} which is given in (12).

Theorem 4.6. *Let $i, k \in S, j \in N; a_{ij} \leq r_{ij}(h_j) \leq r_{ij}(H_j)$ for all $i \in S, j \in N$; then $\exists \gamma_{ikj} \geq 0$ such that $P_{ikj}(t) \neq \emptyset \Leftrightarrow t \geq \gamma_{ikj}$.*

Proof. The proof is obviously derived from the above remarks and from the definition of γ_{ikj} which is given in (13).

To generalize the above three theorems we introduce the following lemmas and remarks.

Lemma 4.1. *Let $i, k \in S, j \in N$; then $\exists \delta^{ij} \geq 0$ such that $V_j(t) \cap W_{ij}(t) \neq \emptyset \Rightarrow t \geq \delta^{ij}$.*

Proof. Let $\delta^{ij} = \max(\tau^{ij}, \eta^{ij})$; assume that for some fixed $i \in S, j \in N$ (say i_0, j_0); $n^{j_0} \geq \tau^{i_0 j_0}$; then $t < \delta^{i_0 j_0} \Rightarrow t < \eta^{j_0} \Rightarrow V_{j_0}(t) \cap W_{i_0 j_0}(t) = \emptyset$ (th. 4.2). Similarly we can treat the other case, and then the proof is complete. \square

Lemma 4.2. *Let $i, k \in S, j \in N$; then $\exists \beta \geq 0$ such that $V_j(t) \cap W_{ij}(t) \neq \emptyset \Rightarrow t \geq \beta$.*

Proof. Let $\beta = \max_{i \in S} \min_{j \in N} \delta^{ij}$.

Assume that $t < \beta \Rightarrow$ for some fixed $i \in S, j \in N$ (say i_0, j_0) we have $t < \delta^{i_0 j_0}$

$$\rightarrow t < \max(\tau^{i_0 j_0}, \eta^{j_0})$$

$$\Rightarrow V_{j_0}(t) \cap W_{i_0 j_0}(t) = \emptyset \text{ (th. 4.1),}$$

and then the proof is complete. \square

Remarks.

- From lemma 4.1 and lemma 4.2 we have $V_j(t) \neq \emptyset$ and $W_{ij}(t) \neq \emptyset$ if $t \geq \beta$ i.e. $t \in [\beta, \infty)$.
- If $t \geq \max(\tau^{ij}, \eta^j)$, then $V_j(t) \neq \emptyset$ and $W_{ij}(t) \neq \emptyset$.
- From th. 4.1 and th. 4.2 we have:
 $V_j(t) \neq \emptyset$ and $W_{ij}(t) \neq \emptyset$ if $t < b_i - r_{ij}(h_j)$ and $t \leq \max(\tau^{ij}, \eta^j)$ i.e. $t \in [\max(\tau^{ij}, \eta^j), b_i - r_{ij}(h_j))$.

If we redefine ξ_{ikj} , ζ_{ikj} and γ_{ikj} by replacing η^{ij} by η^j and then by η in (11), (12), (13), then from theorems 4.1 & 4.2, lemmas 4.1 & 4.2 and also from the above remarks, we can prove again the generalized form of theorems 4.4 & 4.5 & 4.6 which obtained by the new formulas of ξ_{ikj} , ζ_{ikj} and γ_{ikj} .

Now let us define the following maximum variables:

$$\xi^{ij} = \max_{k \in S} \xi_{ikj}, \quad \zeta^{ij} = \max_{k \in S} \zeta_{ikj} \quad \text{and} \quad \gamma^{ij} = \max_{k \in S} \gamma_{ikj}. \quad (14)$$

The following three theorems give another sufficient and necessary conditions for

$$V_j(t) \cap W_{ij}(t) \neq \emptyset.$$

Theorem 4.7. *Let $i \in S, j \in N; r_{ij}(h_j) \leq a_{ij} \leq r_{ij}(H_j)$ for all $i \in S, j \in N$; then $\exists \xi^{ij} \geq 0$ such that $V_j(t) \cap W_{ij}(t) \neq \emptyset \Leftrightarrow t \geq \xi^{ij}$; ξ^{ij} is given by (14).*

Proof. The assertion follows immediately from theorem 4.4 and the definition of $V_j(t)$.

Theorem 4.8. *Let $i \in S, j \in N; r_{ij}(h_j) \leq r_{ij}(H_j) \leq a_{ij}$ for all $i \in S, j \in N$; then $\exists \zeta^{ij} \geq 0$ such that $V_j(t) \cap W_{ij}(t) \neq \emptyset \Leftrightarrow t \geq \zeta^{ij}$; ζ^{ij} is given by (14).*

Proof. The assertion follows immediately from theorem 4.5 and the definition of $V_j(t)$.

Theorem 4.9. *Let $i \in S, j \in N; a_{ij} \leq r_{ij}(h_j) \leq r_{ij}(H_j)$ for all $i \in S, j \in N$; then $\exists \gamma^{ij} \geq 0$ such that $V_j(t) \cap W_{ij}(t) \neq \emptyset \Leftrightarrow t \geq \gamma^{ij}$; γ^{ij} is given by (14).*

Proof. The assertion follows immediately from theorem 4.6 and the definition of $V_j(t)$.

From the above results we conclude that there exist some values, say t^j & T^{ij} for which the relations $V_j(t) \neq \emptyset \Leftrightarrow t \geq t^j$ and $W_{ij}(t) \cap V_j(t) \neq \emptyset \Leftrightarrow t \geq T^{ij}$ hold $\forall i : i \in S, \forall j : j \in N$; where T^{ij} is equal to one of the values ξ^{ij}, ζ^{ij} or γ^{ij} according to which of the conditions from Theorems 4.7, 4.8 and 4.9 are satisfied and t^j is the same as η^j which defined in (10). Then the optimal value of $t(t^{opt})$ is calculated according to the following formula:

$$t^{opt} = \max \left(\max_{j \in N} t^j, \max_{i \in S} \min_{j \in N} T^{ij} \right). \quad (15)$$

Consequently we can deduce that, the optimal value of $t(t^{opt})$ is calculated according to the following theorem:

Theorem 4.10. *If t is the solution of problem (9), then t holds one of the following relations:*

$$\begin{aligned} \text{If } r_{ij}(h_j) \leq a_{ij} \leq r_{ij}(H_j) \text{ for all } i \in S, j \in N; \text{ then} \\ t \geq t^{opt} = \max_{i \in S} \min_{j \in N} \xi^{ij}. \end{aligned} \quad (16)$$

$$\begin{aligned} \text{If } r_{ij}(h_j) \leq r_{ij}(H_j) \leq a_{ij} \text{ for all } i \in S, j \in N; \text{ then} \\ t \geq t^{opt} = \max_{i \in S} \min_{j \in N} \zeta^{ij}. \end{aligned} \quad (17)$$

$$\begin{aligned} \text{If } a_{ij} \leq r_{ij}(h_j) \leq r_{ij}(H_j) \text{ for all } i \in S, j \in N; \text{ then} \\ t \geq t^{opt} = \max_{i \in S} \min_{j \in N} \gamma^{ij}. \end{aligned} \quad (18)$$

Where ξ^{ij} , ζ^{ij} and γ^{ij} are given in (14).

Proof. In our proof we will concentrate on the first case. Let $\xi^{iojo} = \max_{i \in S} \min_{j \in N} \xi^{ij}$ and assume that $t < \xi^{iojo}$, then from theorem 4.7 we can deduce that:

$$W_{ioj}(t) \cap V_j(t) = \emptyset; \quad \forall j \in N,$$

hence, according to theorem 3.1; $M(t) = \emptyset$; this complete the proof of the theorem. \square

5 Algorithm for Calculating t^{opt}

Step 1:

Find η^{ij} , η^j , η and τ^{ij} from relations (10), for each $i \in S$ and each $j \in N$.

Step 2:

Calculate η_{ikj} from the equation

$$r_{kj}^{-1}(\hat{b}_k + \eta_{ikj}) = r_{kj}^{-1}(\hat{b}_k - \eta_{ikj})$$

for each i , $i \in S$ and each $j \in N$.

Step 3:

Find ξ_{ikj} , ζ_{ikj} or γ_{ikj} from relations (11), (12), (13) for each $i, k \in S$ and each $j \in N$.

Step 4:

Find ξ^{ij} , ζ^{ij} or γ^{ij} from relations (14) for each $i \in S$ and $j \in N$.

Step 5:

Find t^{opt} from relations (16), (17), (18).

Example.

Here we want to solve the following problem

$$R_i \equiv \max_{j \in N} (a_{ij} \wedge r_{ij}(x_j)) = b_i \quad \forall i \in S$$

and

$$h_j \leq x_j \leq H_j \quad \forall j \in N$$

where

$$N = \{1, 2, 3, 4\}; \quad S = \{1, 2, 3\}; \quad x = [x_1 x_2 x_3 x_4]^T; \\ b = [5 \ 7 \ 3]^T; \quad h = [1 \ 2 \ 0 \ 1]^T; \quad H = [5 \ 6 \ 4 \ 3]^T;$$

$$\begin{bmatrix} r_{11}(x_1) & r_{12}(x_2) & r_{13}(x_3) & r_{14}(x_4) \\ r_{21}(x_1) & r_{22}(x_2) & r_{23}(x_3) & r_{24}(x_4) \\ r_{31}(x_1) & r_{32}(x_2) & r_{33}(x_3) & r_{34}(x_4) \end{bmatrix} \equiv \begin{bmatrix} 4x_1 & 7x_2 & 6x_3 & x_4 + 1 \\ 2x_1 & x_2 & 6x_3 + 1 & 2x_4 \\ x_4 + 1 & x_2 - 1 & 2x_3 - 1 & x_4 - 3 \end{bmatrix}$$

and

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{bmatrix} = \begin{bmatrix} 7 & 15 & 4 & 2 \\ 3 & 4 & 1 & 4 \\ 3 & 3 & 2 & -1 \end{bmatrix}$$

Note that this problem has no solution in general.

It is clear that $r_{ij}(h_j) \leq a_{ij} \leq r_{ij}(H_j) \forall i \in S, j \in N$. Using the relations (10) we can deduce that

$$\begin{bmatrix} \eta^{11} & \eta^{12} & \eta^{13} & \eta^{14} \\ \eta^{21} & \eta^{22} & \eta^{23} & \eta^{24} \\ \eta^{31} & \eta^{32} & \eta^{33} & \eta^{34} \end{bmatrix} = \begin{bmatrix} 0 & 9 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

this gives that

$$[\eta^1 \ \eta^2 \ \eta^3 \ \eta^4] = [0 \ 9 \ 0 \ 0]$$

which implies that $\eta \leq 9$.

Also we get from (10), the following

$$\begin{bmatrix} \tau^{11} & \tau^{12} & \tau^{13} & \tau^{14} \\ \tau^{21} & \tau^{22} & \tau^{23} & \tau^{24} \\ \tau^{31} & \tau^{32} & \tau^{33} & \tau^{34} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 & 3 \\ 4 & 3 & 6 & 3 \\ 0 & 0 & 1 & 4 \end{bmatrix}$$

From the equality

$$r_{kj}^{-1}(b_k + \eta_{ikj}) = r_{ij}^{-1}(b_i - \eta_{ikj})$$

We can calculate η_{ikj} for each $k \in S; i \in S$ and $j \in N$, then the application of the above relation will give us the following three matrices:

at $k = 1$

$$\begin{bmatrix} \eta_{111} & \eta_{112} & \eta_{113} & \eta_{114} \\ \eta_{211} & \eta_{212} & \eta_{213} & \eta_{214} \\ \eta_{311} & \eta_{312} & \eta_{313} & \eta_{314} \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 3 & \frac{11}{2} & \frac{31}{7} & \frac{-1}{3} \\ \frac{3}{5} & \frac{23}{8} & \frac{7}{4} & 1 \end{bmatrix}$$

at $k = 2$

$$\begin{bmatrix} \eta_{121} & \eta_{122} & \eta_{123} & \eta_{124} \\ \eta_{221} & \eta_{222} & \eta_{223} & \eta_{224} \\ \eta_{321} & \eta_{322} & \eta_{323} & \eta_{324} \end{bmatrix} = \begin{bmatrix} -3 & \frac{-11}{2} & \frac{-31}{7} & \frac{1}{3} \\ 0 & 0 & 0 & 0 \\ -1 & \frac{-3}{2} & \frac{-8}{3} & \frac{5}{3} \end{bmatrix}$$

at $k = 3$

$$\begin{bmatrix} \eta_{131} & \eta_{132} & \eta_{133} & \eta_{134} \\ \eta_{231} & \eta_{232} & \eta_{233} & \eta_{234} \\ \eta_{331} & \eta_{332} & \eta_{333} & \eta_{334} \end{bmatrix} = \begin{bmatrix} \frac{-3}{5} & \frac{-23}{8} & \frac{-7}{4} & -1 \\ 1 & \frac{3}{2} & \frac{8}{3} & \frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Now from equation (11) and the values of the above parameters, we can obtain the following matrices

at $k = 1$

$$\begin{bmatrix} \xi_{111} & \xi_{112} & \xi_{113} & \xi_{114} \\ \xi_{211} & \xi_{212} & \xi_{213} & \xi_{214} \\ \xi_{311} & \xi_{312} & \xi_{313} & \xi_{314} \end{bmatrix} \equiv \begin{bmatrix} 0 & 9 & 1 & 3 \\ 2 \text{ if } 0 \leq t \leq 3 & \begin{cases} 10 \text{ if } 1 < t \leq \frac{11}{2} \\ 3 \text{ otherwise} \end{cases} & \begin{cases} -1 \text{ if } 2 < t \leq \frac{31}{7} \\ 6 \text{ otherwise} \end{cases} & \begin{bmatrix} 3 \\ 3 \\ 4 \end{bmatrix} \\ 3/5 & \begin{cases} 10 \text{ if } 0 \leq t \leq \frac{23}{8} \\ 0 \text{ otherwise} \end{cases} & \begin{cases} -1 \text{ if } 0 < t \leq \frac{7}{4} \\ 1 \text{ otherwise} \end{cases} & 4 \end{bmatrix}$$

at $k = 2$

$$\begin{bmatrix} \xi_{121} & \xi_{122} & \xi_{123} & \xi_{124} \\ \xi_{221} & \xi_{222} & \xi_{223} & \xi_{224} \\ \xi_{321} & \xi_{322} & \xi_{323} & \xi_{324} \end{bmatrix} = \begin{bmatrix} 0 & 9 & 1 & 3 \\ 4 & 3 & 6 & 3 \\ 0 & 0 & 1 & 4 \end{bmatrix}$$

at $k = 3$

$$\begin{bmatrix} \xi_{131} & \xi_{132} & \xi_{133} & \xi_{134} \\ \xi_{231} & \xi_{232} & \xi_{233} & \xi_{234} \\ \xi_{331} & \xi_{332} & \xi_{333} & \xi_{334} \end{bmatrix} \equiv \begin{bmatrix} 0 & 9 & 1 & 3 \\ 0 \text{ if } 0 \leq t < 1 & \begin{cases} 0 \text{ if } 1 < t \leq \frac{3}{2} \\ 3 \text{ otherwise} \end{cases} & \begin{cases} -1 \text{ if } 2 < t \leq \frac{10}{3} \\ 6 \text{ otherwise} \end{cases} & \begin{bmatrix} 3 \\ 3 \\ 4 \end{bmatrix} \\ 0 & 0 & 1 & 4 \end{bmatrix}$$

From the definition of ξ^j given in (14), it is easy to obtain the following

$$\begin{bmatrix} \xi^{11} & \xi^{12} & \xi^{13} & \xi^{14} \\ \xi^{21} & \xi^{22} & \xi^{23} & \xi^{24} \\ \xi^{31} & \xi^{32} & \xi^{33} & \xi^{34} \end{bmatrix} \equiv \begin{bmatrix} 0 & 9 & 1 & 3 \\ 4 & \begin{cases} 10 \text{ if } 1 < t \leq \frac{11}{2} \\ 3 \text{ otherwise} \end{cases} & \begin{cases} -1 & 1 & 3 \\ 6 & 3 \end{cases} \\ 3/5 & \begin{cases} 10 \text{ if } 0 < t \leq \frac{23}{8} \\ 0 \text{ otherwise} \end{cases} & \begin{cases} 1 & 4 \end{cases} \end{bmatrix}$$

Hence t^{opt} which given by equation (15) will be

$$t^{opt} = \max(9, 3) = 9$$

Take any point (say x^*) from the set

$$M(b) = \{x: h \leq x \leq H, \|R(x) - b\| \leq t, t \geq 9\},$$

then x^* will be accepted as an approximate solution of our problem.

In the original case $V_{12} = \emptyset$ {since $15 \wedge 7x_2 > 5, x_2 \in [2, 6]$ },

$$V_{12} = \emptyset \rightarrow V_2 = \emptyset \rightarrow M = \emptyset$$

i.e. there is no solution for the original problem.

In the modified case, if we take $t = 9$, then we try to solve the following problem

$$\begin{aligned} -4 &\leq \max 7 \wedge 4x_1, 15 \wedge 7x_2, 4 \wedge 6x_3, 2 \wedge (x_4 + 1) \leq 14 \\ -2 &\leq \max 3 \wedge 2x_1, 4 \wedge x_2, 1 \wedge (x_3 + 1), 4 \wedge 2x_4 \leq 16 \\ -6 &\leq \max 3 \wedge (x_1 + 1), 3 \wedge (x_2 - 1), 2 \wedge (2x_3 - 1), -1 \wedge (x_4 - 3) \leq 12 \\ 1 &\leq x_1 \leq 5, 2 \leq x_2 \leq 6, 0 \leq x_3 \leq 4, 1 \leq x_4 \leq 3. \end{aligned}$$

Then

$$V_1 = [1, 5] \quad V_2 = \{2\} \quad V_3 = [0, 4] \quad V_4 = [1, 3],$$

and

$$V_1 \cap W_{11} \neq \emptyset, \quad V_2 \cap W_{22} \neq \emptyset, \quad V_3 \cap W_{33} \neq \emptyset$$

where

$$W_{11} = [1, 5] \quad W_{22} = [2, 6] \quad W_{33} = [0, 4]$$

then choose any x such that

$$x = (x_1, 2, x_3, x_4)$$

where

$$x_1 \in [1, 5] \quad x_3 \in [0, 4] \quad x_4 \in [1, 3]$$

will be an approximate solution for the original problem.

References

- [1] CECHLÁROVÁ K., CUNNINGHAME-GREEN R. A., *Residuation in Fuzzy Algebra and some Applications*, Preprint No. 93/22, The University of Birmingham.
- [2] CUNNINGHAME-GREEN, R. A., *Minimax Algebra, Lecture Notes in Economics and Mathematical systems*, Springer Verlag, 1979, 166.
- [3] JAJOU A., ZIMMERMANN K., *Max-Separable Optimization Problems with Parameters in the Right-Hand Sides of the Constraints*, 1985.
- [4] PEDRYCZ W., *Inverse Problem in Fuzzy Relation Equations*, *Fuzzy Sets and Systems* 36 (1990), pp. 277–291.
- [5] ZIMMERMANN K., *On Some Extremal Optimization Problems*. *Ekonomicko-matematický obzor*, 1979, No. 4.

- [6] ZIMMERMANN K., *Solution of Some Optimization Problems on Extremal Algebra*, Methods in OR, Studies in Math. Programming (Ed. A. Prékopa) Akademiai Kiadó, Budapest 1980, pp. 179–186.
- [7] ZIMMERMANN K., *The explicit solution of max-separable optimization problem*, Ekonomicko-matematický obzor, 1982, No. 4.
- [8] ZIMMERMANN K., *On max-separable optimization problems*. Annals of Discrete Mathematics 19, 1984, North Holland.