

Ladislav Lukšan; Jan Vlček

A hybrid method for nonlinear least squares that uses quasi-Newton updates applied to an approximation of the Jacobian matrix

In: Jan Chleboun and Pavel Kůs and Petr Příkryl and Miroslav Rozložník and Karel Segeth and Jakub Šístek and Tomáš Vejchodský (eds.): *Programs and Algorithms of Numerical Mathematics, Proceedings of Seminar*. Hejnice, June 24-29, 2018. Institute of Mathematics CAS, Prague, 2019. pp. 99–106.

Persistent URL: <http://dml.cz/dmlcz/703066>

Terms of use:

© Institute of Mathematics CAS, 2019

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://dml.cz>

A HYBRID METHOD FOR NONLINEAR LEAST SQUARES THAT USES QUASI-NEWTON UPDATES APPLIED TO AN APPROXIMATION OF THE JACOBIAN MATRIX

Ladislav Lukšan, Jan Vlček

Institute of Computer Science, The Czech Academy of Sciences
Pod Vodárenskou věží 2, 182 07 Prague 8, Czech Republic
luksan@cs.cas.cz, vlcek@cs.cas.cz

Abstract: In this contribution, we propose a new hybrid method for minimization of nonlinear least squares. This method is based on quasi-Newton updates, applied to an approximation A of the Jacobian matrix J , such that $A^T f = J^T f$. This property allows us to solve a linear least squares problem, minimizing $\|Ad + f\|$ instead of solving the normal equation $A^T Ad + J^T f = 0$, where $d \in R^n$ is the required direction vector. Computational experiments confirm the efficiency of the new method.

Keywords: Nonlinear least squares, hybrid methods, trust-region methods, quasi-Newton methods, numerical algorithms, numerical experiments.

MSC: 65K10, 65F30

1. Introduction

Consider the objective function

$$F(x) = \frac{1}{2} f^T(x) f(x) = \frac{1}{2} \sum_{k=1}^m f_k^2(x), \quad (1)$$

where $f: R^n \rightarrow R^m$ is a twice continuously differentiable mapping with elements $f_k(x)$, $1 \leq k \leq m$. Let $J(x)$ be its Jacobian matrix with elements $J_{kl}(x) = \partial f_k(x) / \partial x_l$, where $1 \leq k \leq m$ and $1 \leq l \leq n$. Then the gradient and the Hessian matrix of function (1) have the form

$$g(x) = J^T(x) f(x) = \sum_{k=1}^m f_k(x) g_k(x), \quad (2)$$

$$G(x) = J^T(x) J(x) + C(x) = \sum_{k=1}^m g_k(x) g_k^T(x) + \sum_{k=1}^m f_k(x) G_k(x), \quad (3)$$

where $g_k(x)$ and $G_k(x)$ are gradients and Hessian matrices of functions $f_k(x)$, $1 \leq k \leq m$. The most known methods for minimization of the objective function (1) are trust-region realizations of the Gauss-Newton method, which are iterative and their iterations have the form

$$x_{i+1} = x_i, \quad \Delta_{i+1} < \Delta_i \quad \text{if} \quad \frac{F(x_i + d_i) - F(x_i)}{Q_i(d_i)} < \underline{\rho}, \quad (4)$$

$$x_{i+1} = x_i + d_i, \quad \Delta_{i+1} \geq \Delta_i \quad \text{if} \quad \frac{F(x_i + d_i) - F(x_i)}{Q_i(d_i)} \geq \underline{\rho}, \quad (5)$$

where $0 < \underline{\rho} < 1$, $Q_i(d) = g(x_i)^T d + (1/2)d^T B_i d$, $B_i = J(x_i)^T J(x_i)$ and d_i is an approximate minimum of the quadratic function $Q_i(d)$ on the trust region defined by constraint $\|d\| \leq \Delta_i$ [2], [8]. Let $x^* \in R^n$ be a minimum of function (1). The Gauss-Newton method works well if $F(x^*)$ is small (if $F(x^*) = 0$, the rate of convergence is superlinear), but the convergence can be slow if $F(x^*)$ is large. Thus hybrid methods, which are combinations of the Gauss-Newton method and variable metric methods, are advantageously used. In the subsequent text, we use the notation $F_i = F(x_i)$, $g_i = g(x_i)$, $G_i = G(x_i)$, etc, and $F^* = F(x^*)$, $g^* = g(x^*)$, $G^* = G(x^*)$, etc. Sometimes index i is omitted and index $i+1$ is replaced by the symbol $+$. More details concerning methods described in this contribution can be found in [6].

2. Hybrid methods

Hybrid methods are based on the fact that $(F_i - F_{i+1})/F_i \rightarrow 1$, if $F_i \rightarrow F^* = 0$ Q -superlinearly, and $(F_i - F_{i+1})/F_i \rightarrow 0$, if $F_i \rightarrow F^* > 0$. This fact forms the basis for a simple hybrid method in [1]: Let $B_1 = J_1^T J_1$. If $(F_i - F_{i+1})/F_i \geq \underline{\varrho}$, we set $B_{i+1} = J_{i+1}^T J_{i+1}$. If $(F_i - F_{i+1})/F_i < \underline{\varrho}$, we set

$$B_{i+1} = \frac{1}{\gamma_i} (B_i + [y_i, B_i s_i] M_i^B [y_i, B_i s_i]^T), \quad (6)$$

where $s_i = x_{i+1} - x_i$, $y_i = g_{i+1} - g_i = J_{i+1}^T f_{i+1} - J_i^T f_i$, $\gamma_i > 0$ and the matrix $M_i^B \in R^{2 \times 2}$ is chosen in such a way that the quasi-Newton condition $B_{i+1} s_i = y_i$ is satisfied [7]. This simple hybrid method switches between the Gauss-Newton method and a selected variable metric method (defined by matrix M_i^B). More complicated hybrid methods are based on structured variable metric updates [4]: Let $C_1 = 0$ and $B_i = J_i^T J_i + C_i$. If $(F_i - F_{i+1})/F_i \geq \underline{\varrho}$, we set $C_{i+1} = 0$. If $(F_i - F_{i+1})/F_i < \underline{\varrho}$, we set

$$C_{i+1} = \frac{1}{\gamma_i} (C_i + [z_i, C_i s_i] M_i^C [z_i, C_i s_i]^T), \quad (7)$$

where $s_i = x_{i+1} - x_i$, $z_i = J_{i+1}^T f_{i+1} - J_i^T f_{i+1}$, $\gamma_i > 0$ and the matrix $M_i^C \in R^{2 \times 2}$ is chosen in such a way that the quasi-Newton condition $C_{i+1} s_i = z_i$ is satisfied, see [7].

If matrix B_i is ill-conditioned, then a more advantageous way is to use a full rank approximation A_i of the Jacobian matrix J_i and replace the solution of the normal

equation $d_i = -B_i^{-1}g_i$, where $B_i = A_i^T A_i$, by the solution of the linear least-squares problem $d_i = -A_i^\dagger f_i$. This approach is used in [13], where matrix A_i is expressed as a sum $A_i = J_i + L_i$ and matrix L_i is updated to satisfy the quasi-Newton condition $(J_{i+1} + L_{i+1})^T (J_{i+1} + L_{i+1}) s_i = y_i$. This approach is not quite rigorous, since usually $A_i^\dagger f_i \neq B_i^{-1}g_i$. The equality $A_i^\dagger f_i = B_i^{-1}g_i$ is satisfied only if $A_i^T f_i = g_i = J_i^T f_i$. For this purpose, the additional condition $L_{i+1} f_{i+1} = 0$ was added to the above quasi-Newton condition in [11]. In this contribution, we confine our attention to the simple hybrid method of the form (6). By a variational principle, we derive the update, which satisfies the quasi-Newton condition $A_{i+1}^T A_{i+1} s_i = y_i$ together with the condition $A_{i+1}^T f_{i+1} = g_{i+1} = J_{i+1}^T f_{i+1}$.

3. New hybrid method

Let $B = A^T A$, where $A = J$ if the Gauss-Newton step is accepted, so $B = J^T J$ holds. To use the variational principle, we write the standard quasi-Newton condition $B_+ s = A_+^T A_+ s = y$ in the form

$$\sqrt{\gamma} A_+ s = \tilde{z}, \quad \sqrt{\gamma} A_+^T \tilde{z} = \gamma y, \quad \tilde{z}^T \tilde{z} = \gamma s^T y, \quad (8)$$

where $\tilde{z} \in R^m$ is a free vector parameter. Notice that the last equality, which is a consequence of the first two equalities, is the only restriction on the choice of \tilde{z} .

Theorem 1. *Let W be a symmetric positive definite matrix. Then the Frobenius norm $\|W^{-1/2}(\sqrt{\gamma} A_+ - A)^T\|_F$ is minimal on the set of all matrices satisfying quasi-Newton condition (8) if and only if*

$$\sqrt{\gamma} A_+^T = A^T - \frac{W s}{s^T W s} \tilde{s}^T + \left(\gamma y - z + s^T z \frac{W s}{s^T W s} \right) \frac{\tilde{z}^T}{\tilde{z}^T \tilde{z}}, \quad \tilde{z}^T \tilde{z} = \gamma s^T y. \quad (9)$$

where $\tilde{s} = A s$ and $z = A^T \tilde{z}$.

Proof. This proof is similar to the proof of Theorem 3.1 proposed in [12]. Denote $X = \sqrt{\gamma} A_+^T$. Necessity will be proven using the Lagrangian function

$$\begin{aligned} L &= \frac{1}{2} \|W^{-1/2} (X - A^T)\|_F^2 + \tilde{u}^T (X^T s - \tilde{z}) + v^T (X \tilde{z} - \gamma y) \\ &= \sum_{i=1}^m \left[\frac{1}{2} (\xi_i - d_i)^T W^{-1} (\xi_i - a_i) + \tilde{u}_i s^T \xi_i + \tilde{z}_i v^T \xi_i \right] - \tilde{u}^T \tilde{z} - \gamma v^T s, \end{aligned}$$

where $A^T = [a_1, \dots, a_m]$ and $X = [\xi_1, \dots, \xi_m]$. Differentiating the Lagrangian function we obtain

$$\frac{\partial L}{\partial \xi_i} = W^{-1} (\xi_i - a_i) + \tilde{u}_i s + \tilde{z}_i v.$$

Therefore, the conditions for stationarity of the Lagrangian function have the form $W^{-1}(\xi_i - a_i) + \tilde{u}_i s + \tilde{z}_i v = 0$, $1 \leq i \leq m$, or

$$X - A^T = -W s \tilde{u}^T - W v \tilde{z}^T.$$

Using the first condition from (8) we obtain

$$X^T s = As - s^T W s \tilde{u} - v^T W s \tilde{z} = \tilde{z} \quad \Rightarrow \quad \tilde{u} = \frac{1}{s^T W s} (As - (1 + v^T W s) \tilde{z}),$$

which after substitution to the previous equality gives

$$X - A^T = -\frac{W s}{s^T W s} \tilde{s}^T + w \tilde{z}^T,$$

where $w \in R^n$ is an unknown vector (determined uniquely by vector v). Using the second condition from (8) we obtain

$$X \tilde{z} = A^T \tilde{z} - s^T A^T \tilde{z} \frac{W s}{s^T W s} + \tilde{z}^T \tilde{z} w = \gamma y \quad \Rightarrow \quad w = \frac{1}{\tilde{z}^T \tilde{z}} \left(\gamma y - z + s^T z \frac{W s}{s^T W s} \right),$$

which after substitution to the previous equality (with using relation $X = \sqrt{\gamma} A_+^T$) gives (9). Sufficiency follows from the convexity of the Frobenius norm. \square

Update (9) contains two vector parameters $W s / s^T W s$ and \tilde{z} . These parameters should be chosen in such a way to guarantee condition $A_+^T f_+ = g_+$.

Lemma 1. *Equalities*

$$\sqrt{\gamma} A_+ s = \tilde{z}, \quad \sqrt{\gamma} A_+^T \tilde{z} = \gamma y, \quad A_+^T f_+ = g_+ \quad (10)$$

can be satisfied simultaneously only if

$$f_+^T f_+ s^T y \geq (s^T g_+)^2. \quad (11)$$

Proof. From the first two equalities in (10), the relation $\tilde{z}^T \tilde{z} = \gamma s^T y$ follows, which determines the norm of vector \tilde{z} . The first and the third equalities imply $f_+^T \tilde{z} = \sqrt{\gamma} f_+^T A_+ s = \sqrt{\gamma} s^T g_+$. Since the distance of the hyperplane $f_+^T \tilde{z} = \sqrt{\gamma} s^T g_+$ from the origin is equal to $\sqrt{\gamma} |s^T g_+| / \|f_+\|$, the norm of vector \tilde{z} cannot be smaller than this number, which together with equality $\|\tilde{z}\| = \sqrt{\gamma s^T y}$ gives $\sqrt{\gamma} |s^T g_+| / \|f_+\| \leq \sqrt{\gamma s^T y}$, or $f_+^T f_+ s^T y \geq (s^T g_+)^2$. \square

Remark 1. If perfect line search is used, then $s_i^T g_{i+1} = 0$ holds in every iteration, so $s_i^T y_i = s_i^T g_{i+1} - s_i^T g_i = -s_i^T g_i > 0$, and condition (11) is always satisfied. If the strong Wolfe condition is used (see [10]), then $|s_i^T g_{i+1}| \leq \varepsilon_2 |s_i^T g_i|$ holds in every iteration, so $s_i^T y_i = s_i^T g_{i+1} - s_i^T g_i \geq (1 - \varepsilon_2) |s_i^T g_i|$, and condition (11) is satisfied whenever

$$f_{i+1}^T f_{i+1} \geq \frac{\varepsilon_2^2}{1 - \varepsilon_2} |s_i^T g_i|. \quad (12)$$

If $x_i \rightarrow x^*$ (so $g_i \rightarrow 0$ and $s_i \rightarrow 0$) and $F(x^*) > 0$, there exists an index $k \in N$ such that condition (12) (and therefore also condition (11)) is satisfied $\forall i \geq k$. Moreover, in our numerical experiments with Algorithm 1, the condition (11) was always satisfied, if $F_i - F_{i+1} \leq \vartheta F_i$ with $\vartheta = 0.0005$.

Theorem 2. Let vectors f_+ and As be linearly independent and assume the inequality (11) holds. If we use vectors

$$\tilde{z} = \sqrt{\gamma}(\lambda_1 f_+ + \lambda_2 As), \quad (13)$$

where

$$\lambda_2^2 = \frac{s^T y f_+^T f_+ - (s^T g_+)^2}{f_+^T f_+ s^T A^T A s - (s^T A^T f_+)^2}, \quad \lambda_1 = \frac{s^T g_+ - \lambda_2 s^T A^T f_+}{f_+^T f_+}, \quad (14)$$

and

$$\frac{Ws}{s^T W s} = \frac{\gamma s^T y (A^T f_+ - \sqrt{\gamma} g_+) + \sqrt{\gamma} s^T g_+ (\gamma y - A^T \tilde{z})}{\gamma s^T y s^T A^T f_+ - \sqrt{\gamma} s^T g_+ s^T A^T \tilde{z}}, \quad (15)$$

in formula (9), then equalities (10) hold.

Proof. Vector \tilde{z} has to satisfy equalities $f_+^T \tilde{z} = \sqrt{\gamma} s^T g_+$ and $\tilde{z}^T \tilde{z} = \gamma s^T y$. Setting $\tilde{z} = \sqrt{\gamma}(\lambda_1 f_+ + \lambda_2 As)$, we obtain the system of equations

$$\begin{aligned} \lambda_1 f_+^T f_+ + \lambda_2 s^T A^T f_+ &= \sqrt{\gamma} s^T g_+, \\ \lambda_1^2 f_+^T f_+ + 2\lambda_1 \lambda_2 s^T A^T f_+ + \lambda_2^2 s^T A^T A s &= \gamma s^T y \end{aligned}$$

for unknowns λ_1 and λ_2 . Since the vectors f_+ and As are linearly independent, these equations have the unique solution given by (14). Update (9) satisfies the first two equalities in (10) (Theorem 1). Using the third equality, we obtain

$$\begin{aligned} \sqrt{\gamma} g_+ &= A^T f_+ - \frac{Ws}{s^T W s} s^T A^T f_+ + \left(\gamma y - A^T \tilde{z} + s^T A^T \tilde{z} \frac{Ws}{s^T W s} \right) \frac{\tilde{z}^T f_+}{\tilde{z}^T \tilde{z}} \\ &= A^T f_+ - \left(s^T A^T f_+ - s^T A^T \tilde{z} \frac{\sqrt{\gamma} s^T g_+}{\gamma s^T y} \right) w + (\gamma y - A^T \tilde{z}) \frac{\sqrt{\gamma} s^T g_+}{\gamma s^T y}, \end{aligned}$$

where $w = Ws/s^T W s$. This relation implies that

$$w = \lambda \left(A^T f_+ - \sqrt{\gamma} g_+ + (\gamma y - A^T \tilde{z}) \frac{\sqrt{\gamma} s^T g_+}{\gamma s^T y} \right), \quad (16)$$

and since $s^T w = s^T W s / s^T W s = 1$, one can write

$$\lambda (\gamma s^T y s^T A^T f_+ - \sqrt{\gamma} s^T g_+ s^T A^T \tilde{z}) = \gamma s^T y.$$

Substituting this value λ into (16), we obtain (15). \square

The above considerations are summarized in the following algorithm.

Algorithm 1

Data: Trust-region parameters [8], update parameter $\underline{\vartheta} = 0.0005$, termination parameters $\underline{\varepsilon} = 10^{-15}$, $\bar{\varepsilon} = 10^{-5}$.

Step 1: Initiation. Choose starting point $x_1 \in R^n$ and initial trust-region radius $\Delta_1 > 0$. Compute $f_1 = f(x_1)$, $J_1 = J(x_1)$, $F_1 = (1/2)f_1^T f_1$, $g_1 = J_1^T f_1$. Set $A_1 = J_1$ and $i = 1$

Step 2: Termination. If $F_i \leq \underline{\varepsilon}$ or $\|g_i\| \leq \bar{\varepsilon}$, then terminate the computation.

Step 3: Direction determination. Determine direction vector d_i using a trust-region strategy (see [8]). Compute $f(x_i + d_i)$, $F(x_i + d_i) = (1/2)f(x_i + d_i)^T f(x_i + d_i)$. Determine x_{i+1} and Δ_{i+1} by (4)–(5).

Step 4: Decision. If $x_{i+1} = x_i$, go to Step 2. If $x_{i+1} \neq x_i$, set $f_{i+1} = f(x_i + d_i)$, $F_{i+1} = F(x_i + d_i)$ and compute $J_{i+1} = J(x_i + d_i)$, $g_{i+1} = g(x_i + d_i)$.

Step 5: Update. If $(F_i - F_{i+1})/F_i \geq \underline{\vartheta}$, set $A_{i+1} = J_{i+1}$. If $(F_i - F_{i+1})/F_i < \underline{\vartheta}$, compute matrix A_{i+1} by (9) with (13)–(15).

Step 6: Increase i by 1 and go to Step 2.

4. Computational experiments

Methods for nonlinear least-squares were tested by using 80 problems with 200 variables taken from the collection TEST24 contained in the software system for universal functional optimization UFO [9]. Table 1 contains results obtained by the following methods:

- GN - Gauss-Newton's method,
- HN - New hybrid method (Algorithm 1),
- HS - Structured hybrid method proposed in [11],
- QN - New quasi-Newton method proposed in [8],
- QB - Broyden's good quasi-Newton method, see [8].

These methods were implemented as dog-leg trust-region methods (see [8]). Individual methods were realized in two different ways. First the rectangular matrix A was updated and the updated matrix was decomposed as the product $A = QR$ by the standard way. Secondly, the matrices Q and R in the QR decomposition were updated using the algorithm described in [3] (so only the Jacobian matrix was decomposed and the number of QR decompositions was decreased).

Table 1 proposes results obtained by solving 80 problems with 200 variables. Notice that 80 % of these problems have zero residuals, so they are not quite suitable for comparing hybrid methods with the Gauss-Newton method. This table contains the total numbers of iterations NIT, function evaluations NFV, Jacobian (or gradient) evaluations NFJ, matrix decompositions NDC, the total number of failures (number of unsolved problems) F and the total computational time.

	Rectangular matrix update						QR decomposition update					
	NIT	NFV	NFJ	NDC	F	Time	NIT	NFV	NFJ	NDC	F	Time
GN	3376	3698	3454	3256	-	38.31	5867	6421	5946	5698	1	60.82
HN	2477	2730	2556	2372	-	26.53	4046	4389	4126	2603	-	29.29
HS	2477	2716	2557	2395	-	27.61	3319	3665	3399	3354	-	35.45
QN	5473	5988	6435	5413	-	69.37	8064	8566	9102	606	1	22.46
QB	6904	8092	928	6531	-	76.54	8513	9899	1294	1285	1	25.80

Table 1: TEST24 – 80 problems with 200 variables

The results contained in Table 1 imply several conclusions:

- Hybrid methods HN and HS are more robust than Gauss-Newton method GN, since they increase the rate of convergence for large residual problems. The new method seems to be better than structured hybrid method HS, especially if the efficiency is measured by the computational time.
- Quasi-Newton methods [8], developed originally for solving nonlinear equations, are surprisingly efficient, if they are applied to the QR decomposition of the matrix A , especially if the efficiency is measured by the computational time.

For better understanding, the methods that update rectangular matrix (the first part of Table 1) are also compared by using performance profiles proposed in [5]. In Figure 1, value $\rho_M(0)$ is the percentage of the test problems for which method M is the best and value $\rho_M(\tau)$ for τ large enough is the percentage of the problems that method M can solve. Performance profiles show the relative efficiency and reliability of the methods: the higher is the particular curve, the better is the corresponding method.




Figure 1: Test 24 – 80 problems with 200 variables

Notice that the Gauss-Newton method has a better score for $\tau = 0$, since 80 % of problems used have zero residuals.

Acknowledgements

This work was supported by the Institute of Computer Science of the CAS (RVO: 67985807).

References

- [1] Al-Baali, M. and Fletcher, R.: Variational methods for nonlinear least squares. J. Optimizatoin Theory and Applications **36** (1985) 405-421.

- [2] Conn, A.R., Gould, N.I.M. and Toint, P.L.: *Trust-Region Methods* SIAM, Philadelphia, 2000.
- [3] Daniel, J.W., Gragg, W.B., Kaufman, L. and Stewart G.W.: Reorthogonalization and Stable Algorithms for Updating the Gram-Schmidt QR Factorization. *Mathematics of Computation* 30, (1976) 772-795.
- [4] Dennis, J.E., Martinez, H.J. and Tapia, R.A.: Convergence theory for the structured BFGS secant method with an application to nonlinear least squares. *J. Optimizaton Theory and Applications* **61** (1989) 161-178.
- [5] Dolan, E.D. and More, J.J.: Benchmarking optimization software with performance profiles. *Mathematical Programming* **91** (2002) 201-213.
- [6] Lukšan, L.: Numerické optimalizační metody. Výzkumná zpráva V-1152. Praha, ICS AS CR, 2015 (<http://www.cs.cas.cz/luksan/lekce4.pdf>).
- [7] Lukšan, L. and Spedicato, E.: Variable metric methods for unconstrained optimization and nonlinear least squares. *Journal of Computational and Applied Mathematics* **124** (2000) 61-93.
- [8] Lukšan, L. and Vlček, J.: New quasi-Newton method for solving systems of nonlinear equations. *Applications of Mathematics* **62** (2017) 121-134. Presented on PANM18.
- [9] Lukšan, L., Tůma, M., Vlček, J., Ramešová, N., Šiška, M., Hartman, J., and Matonoha, C.: UFO 2017. Interactive System for Universal Functional Optimization. Technical Report V-1252. Prague, Institute of Computer Science AS CR, Prague 2017 (<http://www.cs.cas.cz/luksan/ufo.pdf>).
- [10] Nocedal, J. and Wright, S.J.: *Numerical Optimization*. Springer, New York, 2006.
- [11] Sheng, S.B. and Zou, Z.H.: A new secant method for nonlinear least squares problems. *Numerical Mathematics. A Journal of Chinese Universities*, **2** (1993) 125137.)
- [12] Vlček, J. and Lukšan, L.: Shifted limited-memory variable metric methods for large-scale unconstrained minimization. *J. of Computational and Applied Mathematics*, **186** (2006) 365-390.
- [13] Yabe, H. and Takahashi, T.: Factorized quasi-Newton methods for nonlinear least squares problems. *Mathematical Programming* **51** (1991) 75-100.