

Otomar Hájek

Characteristics of modular finite-length lattices

Czechoslovak Mathematical Journal, Vol. 15 (1965), No. 4, 521–525

Persistent URL: <http://dml.cz/dmlcz/100691>

Terms of use:

© Institute of Mathematics AS CR, 1965

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CHARACTERISTICS OF MODULAR FINITE-LENGTH LATTICES

ОТОМАР ХАЈЕК, Praha

(Received December 12, 1963)

Two integer-valued characteristics of modular lattices of finite length are described, and their relation with the defect (as introduced in [1]) exhibited.

This paper is closely connected with [1], and the notation, terminology and definitions of [1] are assumed (the results of [1] will be referred to directly). In particular, m. l. f. l. means modular lattice of finite length.

We shall introduce, in definitions 1 and 2, two integer-valued characteristics of m. l. f. l., with apparently intuitive meaning.

Definition 1. Let L be a lattice not $\mathbf{1}$. The discriminator $d(L)$ of L is the least cardinal n such that for any $x \neq y$ in L there is a lattice M with $l_M \leq n$ and a homomorphism $h : L \rightarrow M$ such that $hx \neq hy$.

We have immediately the following elementary properties: $d(L)$ is not changed if we also require h to be onto. Also $d(L) \leq l_L$, so that $d(L)$ is finite if L has finite length. If L_1 is a sublattice or a factor lattice of L (i.e. $L_1 = L/\theta$ for some congruence relation θ on L) then $d(L_1) \leq d(L)$. Finally,

$$(1) \quad L \text{ simple implies } d(L) = l_L.$$

Lemma 1. *If L is a m. l. f. l. then $d(L)$ in definition 1 is not changed if we require further that M be simple.*

Proof. Let $h : L \rightarrow M$ be a homomorphism onto; then $M \approx L/\theta$ for the congruence relation θ defined by $x \equiv y (\theta)$ iff $h(x) = h(y)$. Since the congruence lattice Θ_L of a m. l. f. l. is a Boolean algebra, θ is the intersection of all dual atoms $\beta \geq \theta$. Thus, if $hx \neq hy$, then $x \not\equiv y (\theta)$ and hence $x \not\equiv y (\beta)$ for some dual atom β . But then $S = L/\beta$ is simple, the natural homomorphism $L \rightarrow S$ does not identify x, y , and $l(L/\beta) \leq l(L/\theta) = l_M$. This completes the proof.

For m. l. f. l. L , let B be the set of all dual atoms β in Θ_L (i. e. the set of all congruence relations θ such that L/θ is simple nontrivial). Then we have from lemma 1

$$(2) \quad d(L) = \max_B l(L/\beta)$$

and thus, since $\delta_M = l_M - 2$ for simple m. l. f. l.,

$$(3) \quad d(L) = 2 + \max_B \delta(L/\beta).$$

From (1) and (2) we then conclude

$$(4) \quad d(L) = \max_B d(L/\beta).$$

Lemma 2. *Let L be a m. l. f. l. If $L \leq \mathbf{P}L_a$ then*

$$(5) \quad d(L) = \max d(L_a).$$

L is simple if and only if $d(L) = l_L$.

Proof. Let $L_a = L/\theta_a$. Then $d(L_a) = d(L/\theta_a) \leq d(L)$, so that

$$(6) \quad \max d(L_a) \leq d(L).$$

Since the decomposition is subdirect, we have $\bigwedge \theta_a = O$, so that each dual atom $\beta \geq \theta_a$ for some θ_a , and then

$$l(L/\beta) \leq \max \{l(L/\beta) : \theta_a \leq \beta \in B\} = d(L_a)$$

on applying (2). Thus $l(L/\beta) \leq \max d(L_a)$, and (2) again yields $d(L) \leq \max d(L_a)$. With (6), this proves (5). As for the second statement of lemma 2, we already have (1). If L is not simple, then it is an exact subdirect product of more than one simple nontrivial m. l. f. l. (cf. [1], lemmas 9, 13, and (12)): $L \leq \mathbf{P}M_j$ with

$$l_L - 1 = \sum (l(M_j) - 1), \quad l(M_j) \geq 2,$$

and hence (5 and 1)

$$d(L) = \max d(M_j) = \max l(M_j) = l(M_k) < l_L,$$

so that $d(L) = l_L$ is excluded. This completes the proof of lemma 2.

Definition 2. The trivialiser $t(L)$ of a finite-length lattice $L (\neq \mathbf{1})$ is the maximal integer n such that every homomorphism $h : L \rightarrow M$ with $l_M \leq n$ is constant.

Obviously $1 \leq t(L) < l(L)$; also

$$(7) \quad L \text{ simple implies } t(L) = l(L) - 1.$$

Lemma 3. *In definition 2, if L is a m. l. f. l. then we may add the requirement that M be simple.*

Proof. Definition 2 may also be formulated thus (h denotes a homomorphism): $t(L) \geq n$ iff $h : L \rightarrow M$ is constant whenever $l_M \leq n$. Now define: $t_1(L) \geq n$ iff $h : L \rightarrow M$ is constant whenever $l_M \leq n$ and M is simple.

Then obviously $t(L) \leq t_1(L)$. Now assume $t(L) + 1 \leq t_1(L)$, and aim at a contradiction. There is a nonconstant $h : L \rightarrow M$ with $l_M \leq t(L) + 1$; necessarily though,

$l_M = t(L) + 1$. Since h is nonconstant and $l_M \leq t_1(L)$, M cannot be simple. Then, as in the proof of lemma 2,

$$M \leq \mathbf{P}M_j, \quad M_j \text{ simple not } \mathbf{1},$$

$$l(M_j) - 1 \leq l_M - 1 = t(L) < t_1(L).$$

Thus $l(M_j) \leq t_1(L)$; by definition, every $\eta_j h : L \rightarrow M_j$ is constant (η_j is the natural projection $M \rightarrow M_j$). But then h itself is constant; this contradiction proves $t(L) \geq t_1(L)$ and our lemma.

Using lemma 3, we may conclude immediately that

$$(8) \quad t(L) = -1 + \min_B l(L/\beta)$$

where B is the set of all dual atoms β of Θ_L . Using (7),

$$(9) \quad t(L) = \min_B t(L/\beta) = 1 + \min_B \delta(L/\beta).$$

From (3) and (9) there follows

$$t(L) < d(L).$$

Lemma 4. *If L is a m. l. f. l. and $L \leq \mathbf{P}L_a$ then $t(L) = \min t(L_a)$.*

Proof. Essentially, the proof is similar to that of lemma 2. Let $L_a = L/\theta_a, \bigwedge \theta_a = 0$. By (8), $t(L) = \min_{\beta \in B} (-1 + l(L/\beta)) = \min_a \min_{\beta \geq \theta_a} (-1 + l(L/\beta)) = \min_a t(L_a)$.

Lemma 5. *Let L be a m. l. f. l. Then $d(L) = t(L) + 1$ if and only if L is a subdirect product of simple lattices all with length $d(L)$ (i.e. defect $d(L) - 2$).*

The proof follows from lemmas 2, 4 and formulae (2), (8).

The representation theorems of [1], theorems 4 and 5, may now be completed by the following (D and M have the same meaning as in theorem 4, l.c.).

Theorem. *Let L be a m. l. f. l. Then there is a unique subdirect decomposition $L \leq \mathbf{P}L_j$ ($0 \leq j \leq l_L - 1$) such that, for each j , either $L_j = \mathbf{1}$ or*

$$(10) \quad d(L_j) - 2 = t(L_j) - 1 = j.$$

Furthermore the decomposition is then exact, $L_0 = D$ is finite distributive, every L_j is exactly decomposable into say n_j simple lattices with defects j ($n_j = 0$ iff $L_j = \mathbf{1}$);

$$\begin{aligned} \delta(L_j) &= n_j \cdot j, & l(L_j) &= 1 + n_j(j + 1), & \lambda(L_j) &= 1 + n_j; \\ \delta_L &= \sum n_j \cdot j, & l_L &= 1 + \sum n_j(j + 1), & \lambda_L &= 1 + \sum n_j; \\ l_M &= 1 + \sum_{j \geq 1} n_j(j + 1), & \lambda_M &= 1 + \sum_{j \geq 1} n_j. \end{aligned}$$

Proof. First prove existence. If B is the set of all dual atoms of Θ_L , set

$$B_j = \{\beta \in B : \delta(L/\beta) = j\}, \quad \theta_j = \bigwedge \{\beta : \beta \in B_j\}, \quad L_j = L/\theta_j.$$

Then either B_j is empty and $L_j = \mathbf{1}$, or (10) holds (lemma 5: L_j is an exact subdirect product of simple L/β with $\beta \in B_j$). Obviously $B = \bigcup B_j$ is a disjoint decomposition, so that L decomposes subdirectly into the L_j and this decomposition is exact.

As mentioned, L_j is an exact subdirect product of simple lattices, say M_{ij} , with the same defect j . By exactness, then, $\delta(L_j) = \sum_i \delta(M_{ij}) = n_j \cdot j$ with n_j integral (and $n_j = 0$ iff $L_j = \mathbf{1}$). Then n_j is the number of simple exact factors of L_j ; thus (cf. [1], lemma 9) $\lambda_L - 1 = \sum n_j$. Also $l_L - 1 = \delta_L + \lambda_L - 1 = \sum n_j \cdot j + \sum n_j$ by the previous result. This proves the formulae for λ_L, l_L ; those for λ_M, l_M are similar.

Finally, consider unicity. Thus, let $L \leq \mathbf{PL}/\tau_j$ with $\bigwedge \tau_j = O$, and either $\tau_j = I$ or

$$d(L/\tau_j) - 2 = t(L/\tau_j) - 1 = j.$$

Take any $\tau_j \neq I$; let $\beta \geq \tau_j$ be any dual atom. Then by (2),

$$l(L/\beta) \leq \max_{\beta \geq \tau_j} l(L/\beta) = d(L/\tau_j) = j + 2$$

and by (8),

$$-1 + l(L/\beta) \geq \min_{\beta \geq \tau_j} (-1 + l(L/\beta)) = t(L/\tau_j) = j + 1.$$

Thus $l(L/\beta) = j + 2$, and since L/β is simple, $\delta(L/\beta) = j$. Summarising, if $\beta \leq \tau_j$ for a dual atom β , then $\delta(L/\beta) = j$, and hence, by construction, $\beta \in B_j$, $\beta \geq \theta_j$. This proves that $\tau_j \geq \theta_j$ for each j ; from $\bigwedge \tau_j = O$ it then follows that $\tau_j = \theta_j$ for all j . This completes the proof of the theorem.

The two following corollaries may be proved as in [1], corollary to theorem 4, corollary 2 to theorem 5.

Corollary 1. *In the theorem, if L is complemented, then the subdirect decompositions (of L and of L_j) are direct.*

Corollary 2. *Let L, L_a be m. l. f. l. Let L_j and L_{aj} be the factors of the decompositions described in the theorem, of L and L_a , respectively. Then if L is a subdirect (exact, direct) product of L_j 's, then L_j is a subdirect (exact, direct) product of the L_{aj} 's.*

Next, consider some elementary consequences of the formula $l_L - 1 = \sum_0^{l_L-1} n_j(j + 1)$. (Write l in place of l_L .) Since all summands are nonnegative, we have immediately that

$$n_{l-1} = 0, \quad \text{i.e.} \quad L_{l-1} = \mathbf{1}.$$

Now, assume that some L_j is non-simple (L is not "square-free"), i.e. that some $n_j \geq 2$. Then $l - 1 \geq 2(j + 1)$, $j \leq \frac{1}{2}(l - 1) - 1$, and we conclude

$$\text{for } j > \frac{1}{2}(l - 1) - 1, \quad L_j \text{ is simple.}$$

If (and only if) $n_{l-2} \geq 1$ then necessarily all other $n_j = 0$ and $n_{l-2} = 1$, i.e.

$$L_{l-2} \neq \mathbf{1} \text{ implies } L = L_{l-2} \text{ is simple.}$$

Now assume that this is not the case. Then $n_{l-2} = 0$, $l \geq 3$, and if $n_{l-3} \neq 0$ then we have, principally, several cases:

(i) $n_{l-3} \geq 3$ implies $l - 1 \geq 3(l - 2)$ and $l \leq 2$, a contradiction;

(ii) $n_{l-3} = 2$ implies $l - 1 \geq 2(l - 2)$ and $l \leq 3$, thus $l = 3$; since L was assumed non-simple, the only possibilities are $L = \mathbf{3}$ or $L = \mathbf{2}^2$;

(iii) $n_{l-3} = 1$; then $l - 1 = \sum_0^{l-1} n_j(j + 1) = l - 2 + \sum_0^{l-4} n_j(j + 1)$ and we conclude $n_0 = 1$ and all other $n_j = 0$ (for $0 \neq j \neq l - 3$); thus the only nontrivial factors L_j are

$$L_0 = \mathbf{2}, \text{ simple } L_{l-3}.$$

Thus we have that $n_j = 0$ for $j \geq l - 3$ with the following exceptions: L is simple or a subdirect product of a simple lattice and $\mathbf{2}$.

References

[1] Hájek O.: Representation of finite-length modular lattices, Czech. Math. J. 15 (90) (1965), 503—520.

Резюме

ХАРАКТЕРИСТИКИ ДЕДЕКИНДОВЫХ СТРУКТУР КОНЕЧНОЙ ДЛИНЫ

ОТОМАР ГАЕК (Otomar Hájek) Прага

В настоящем продолжении работы [1] определяются две численные характеристики дедекиндовых структур L конечной длины (д. с. к. д.) — дискриминатор $d(L)$ и тривиализатор $t(L)$. Доказана следующая теорема:

Для всякой д. с. к. д. L существует единственное полупрямое разложение $L \cong \prod_{0 \leq j < l_L} \mathbf{P} L_j$ такое, что или $L = \mathbf{1}$ или $d(L_j) - 2 = t(L_j) - 1 = j$.