

Vlastimil Dlab

A remark to a paper of Gh. Pic

Czechoslovak Mathematical Journal, Vol. 17 (1967), No. 3, 467–468

Persistent URL: <http://dml.cz/dmlcz/100791>

Terms of use:

© Institute of Mathematics AS CR, 1967

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A REMARK TO A PAPER OF GH. PIC

VLASTIMIL DLAB, Canberra

(Received February 2, 1967)

Throughout the paper, G denotes a group and, for a natural n , $\{G_n\}$ – the (fully invariant) subgroup of G generated by all $g \in G$ with $g^n = 1$. The symbol $[n_1, n_2, \dots, n_k]$ is used to denote the least common multiple of the natural numbers n_i ($1 \leq i \leq k$).

In his paper [3], GH. PIC has attempted to “dualize” the results of [1] on the powers $\{G^n\}$ of a group G to the case of the subgroups $\{G_n\}$. The following simple theorem improves the results of [3].

Theorem. *If, for $1 \leq i \leq k$, the subgroups $\{G_{n_i}\} \subseteq G$ possess a property \mathcal{P} of type (*), then $\{G_{[n_1, n_2, \dots, n_k]}\}$ possesses the property \mathcal{P} , as well. Here, a property \mathcal{P} is said to be of type (*) if any group possessing \mathcal{P} is locally nilpotent and if, for any finite set π of primes p , the direct product $\prod_{p \in \pi} G_p$ of p -groups G_p possesses \mathcal{P} if and only if all G_p possess \mathcal{P} .*

Proof. In view of the relation $[n_1, n_2, \dots, n_{k-1}, n_k] = [[n_1, n_2, \dots, n_{k-1}], n_k]$, we can restrict ourselves to $k = 2$. Thus, $\{G_{n_1}\}$ and $\{G_{n_2}\}$ are locally nilpotent normal subgroups of G ; therefore, $\{G_{[n_1, n_2]}\} = \{G_{n_1}\} \{G_{n_2}\}$ is locally nilpotent¹⁾ (see e.g. K. HIRSCH [2]). Being generated by elements of finite order, the subgroups $\{G_{n_1}\}$, $\{G_{n_2}\}$ and $\{G_{r_{n_1, n_2}}\}$ are the direct products of their respective Sylow subgroups, finite in number: $\{G_{[n_1, n_2]}\} = \prod_{p \in \pi} G_p$ with $G_p \neq 1$ and $\{G_{n_i}\} = \prod_{p \in \pi} G_p^{(i)}$ for $i = 1, 2$. Moreover, if p^{t_i} is the highest power of p dividing n_i , then all elements $g \in G$ such that $g^{p^{t_i}} = 1$ generate $G_p^{(i)}$ ($i = 1, 2$). Therefore, since $[n_1, n_2] = p^{\max(t_1, t_2)} n'$ with n' relatively prime to p , $G_p = G_p^{(i)}$ for one of $i = 1, 2$. Thus, all G_p possess \mathcal{P} , and, consequently, $\{G_{[n_1, n_2]}\}$ possesses \mathcal{P} , as required.

Since the properties of being nilpotent, nilpotent regular, abelian, direct product of cyclic groups or cyclic are, evidently, of type (*), we get immediately the validity of statements “dual” to those of [1]; this, in particular, shows that the counter-example of § 3 in [3] is defective.

¹⁾ In the same manner, $\{G^{(n_1, n_2, \dots, n_k)}\}$ is (locally) nilpotent provided that all $\{G^{n_i}\}$ are (locally) nilpotent; this answers the third question of F. SZÁSZ [4].

Also, an immediate consequence of Theorem reads that if every cyclic subgroup of group G has the form $\{G_n\}$ for a suitable n , then G is torsion and locally cyclic, i.e. G is a subgroup of the (multiplicative) group of all complex roots of unity.

Bibliography

- [1] *V. Dlab*: A note on powers of a group, *Acta Sci. Math.*, 25 (1964), 177–178.
- [2] *K. A. Hirsch*: Über lokal-nilpotente Gruppen, *Math. Zeitschr.* 63 (1955), 290–294.
- [3] *Gh. Pic*: Sur un théorème de Vl. Dlab, *Czech. Math. J.*, 16 (91) (1966), 545–551.
- [4] *F. Szász*: Bemerkung zu meiner Arbeit „Über Gruppen, deren sämtliche nichttriviale Potenzen zyklische Untergruppen der Gruppe sind“, *Acta Sci. Math.*, 23 (1962), 64–66.

Author's address: Department of Mathematics, Institute of Advanced Studies, Australian National University, Canberra, A.C.T.